

Synthesis and Ion Sensing Properties of New Colorimetric and Fluorimetric Chemosensors based on Bithienyl-Imidazo-Anthraquinone Chromophores


Rosa M. F. Batista,[§] Elisabete Oliveira,[‡] Susana P. G. Costa,[§]
Carlos Lodeiro,^{‡*} M. Manuela M. Raposo^{§*}

Centro de Química, Universidade do Minho, Campus de Gualtar, 4710-057 Braga, Portugal, REQUIMTE, Departamento de Química, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, 2829-516 Monte de Caparica, Portugal

mfox@quimica.uminho.pt, lodeiro@dq.fct.unl.pt

Received Date (will be automatically inserted after manuscript is accepted)

ABSTRACT



Novel colorimetric receptors for fluoride ion sensing containing anthraquinone as a chromogenic signaling unit and imidazo-2,2'-bithiophene binding sites are reported. Well defined color change was observed upon addition of fluoride ions to acetonitrile solutions of receptors 2. Compounds 2a-c, deprotonated after fluoride ion addition, were studied as metal ion chemosensors in the presence of Zn(II), Hg(II) and Cu(II) in acetonitrile solutions, especially compound 2a which displayed a marked change from pink to yellow-gold colors upon complexation.

The design and synthesis of systems that are capable of sensing various biologically and chemically important anions are currently of major interest because anions play a fundamental role in chemical and biological processes.¹

Among various important anionic analytes, the fluoride ion is one of the most significant due to its role in dental care² and treatment of osteoporosis.³

Colorimetric sensors have attracted much attention for allowing so-called “naked-eye” detection in a straightforward and inexpensive manner, offering qualitative and quantitative information without using expensive equipments. A chemosensor usually consists of three parts, a recognition moiety, a spacer and a signal reporter. Hydrogen-bonded donors such as pyrrole/calixpyrrole, (thio)urea, guanidinium, azophenol, dipyrrolylquinoxalines, indolocarbazoles, (di)amino, amide and (benzo)imidazole usually act as anion binding sites. Anthraquinone derivatives are electron acceptor groups that, electronically connected with recognition sites, have proved suitable receptors for the colorimetric sensing of certain anions (e.g. fluoride).^{1,4-7}

Donor-acceptor substituted thiophene derivatives have been used recently due to their optical properties.⁸ Bithienyl-derivatives of anthra[1,2-*d*]imidazole-6,11-dione which possess intramolecular hydrogen bonding between the NH of the imidazole ring with the neighboring quinone carbonyl group are new candidates to investigate hydrogen-bond formation and proton transfer in the receptor-anion interactions. It should be noted for these kind of sensors that the acidity of the NH proton of the imidazole ring can be tuned by changing the electronic properties (donor or acceptor) of the thiophene substituents.^{7,9} These compounds are very promising for applications in various areas, such as medical and environmental sciences, due to the combination of the two sensing effects, color and fluorescence.

Therefore we propose the synthesis of new imidazo-anthraquinone sensors containing bithiophene moieties. The electronic nature of its substituents was used to tune the selectivity and the photophysical properties of the chemosensors. In this communication we propose a simple synthetic route to the imidazo-bithienyl-anthraquinone derivatives **2** in which 1,2-diaminoanthraquinone and several formyl-bithiophenes are used as precursors for the synthesis of new fluorimetric and/or colorimetric sensors. To our knowledge, this is the first time that the synthesis and characterization of colorimetric/fluorimetric properties of anthraquinone derivatives containing imidazo-bithienyl moieties are reported and evaluated for halide and metal ion detection.


Several 5-formyl-2,2'-bithiophenes **1** with different substituents at 5'-position (hydrogen, ethoxy and cyano) were used in the synthesis of sensors **2**. 5-Formyl-2,2'-bithiophene **1a** was available commercially while the synthesis of 5-ethoxy-5'-formyl-2,2'-bithiophene **1b** was recently reported.^{8a} Aldehyde **1c** was synthesized by a Pd(PPh₃)₄ catalyzed cross-coupling reaction of 5-formyl-2-thiophene boronic acid with 2-bromo-5-cyanothiophene in 61% yield.

The synthesis of sensors **2** is outlined in Scheme 1. In the first step, mixtures of 1,2-diaminoanthraquinone and the appropriate formyl-bithiophenes **1a-c** were refluxed for 12 h in ethanol, with formic acid as catalyst, to yield the corresponding imine intermediates which were used in the second step without isolation or further purification.

The cyclization of the imines to the imidazo-anthraquinone derivatives **2** was achieved with lead tetraacetate in acetic acid at room temperature.¹⁰

Recrystallization of the crude products gave the pure heterocyclic sensors **2** in good to excellent yields (67–92%). Compounds **2** were completely characterized by ¹H and ¹³C NMR, IR spectroscopy and HRMS. In DMSO-*d*₆ at 25°C, the ¹H NMR signal of NH proton of compounds **2a-c** were observed from 13.3 to 13.6 ppm. The signal of NH appearing downfield indicated high acidity and strong hydrogen-bonding ability.

Scheme 1. Synthesis of sensors **2**



Compounds **2a-c** were protonated in an acidic acetonitrile solution of HBF₄ or methanesulphonic acids, which resulted in a blue shift in the band centered at 444 nm (**2a**), 462 nm (**2b**) and 512 nm (**2c**) to 432 nm (**2a**), 450 nm (**2b**) and 417 nm (**2c**). (See Supporting information, figures S1 to S3).

Interaction with basic anions (CN⁻, H₂PO₄⁻, CH₃COO⁻) and halide ions (F⁻, Cl⁻, Br⁻ and I⁻) was investigated using spectrophotometric and spectrofluorimetric titrations, by adding a standard acetonitrile solution of the corresponding tetrabutylammonium salts to an acetonitrile solution of compounds **2a-c** at room temperature. In the case of the basic anions, no change occurred upon interaction with cyanide anions and a small red shift was observed only when a large amount of CH₃COO⁻ or H₂PO₄⁻ was added. In both cases the yellow color of the solution changed to pale orange and no emission was detected. After addition of 100 equivalents of each anion, a plateau was observed. (See Supporting information, figure S4).

In the case of chloride, bromide and iodide ions, no change was observed upon addition of up to 20 molar equivalents. (See figures S5 to S7 Supporting information). However, addition of fluoride ions produced a marked red-shift in the absorption for all systems and

also an increase in the intensity of the emission spectra. Due to the deprotonation of the NH to N⁻ in the imidazole ring in an alkaline medium, the UV-visible absorption band was shifted to a longer wavelength. The deprotonation lowers the steric hindrance between the thiophene unit and the NH group, and enables the formation of a more extended π -conjugated system. As can be observed for compound **2a**, the UV-vis band centered at 444 nm, that can be assigned to the π - π^* transition of the chromophore, disappeared gradually and a new band centered at 503 nm appeared, which was assigned to the charge transfer (CT) band. The yellow colour of the sensor solution turned to light pink at the same time. Two well defined isosbestic points were observed at 389 nm and 470 nm for **2a**, which suggested that the stoichiometry of the **2a**-fluoride interaction was 1:1 (Figure 1). For systems **2b** and **2c** similar results were observed. The initial colour observed for compound **2b** was orange and for **2c** was red. The intensity of these colours resulted in a weaker naked eye effect than that observed for compound **2a** (See figure S13 in the supporting information).




Figure 1. Changes in UV-vis and fluorescence (inset) spectra for compound **2a** (1.75×10^{-5} M) in CH₃CN with addition of CH₃CN solution of [(Bu)₄N]F. Absorptions read at 444 nm and 503 nm.

The selectivity towards fluoride ion in compounds **2a-c** was achieved through the introduction of substituents at 5'-position of the bithienyl moiety with different electronic properties (**2b**, electron-donating ethoxy group, and **2c**, electron-withdrawing cyano group), as the number of fluoride ion equivalents needed to fully deprotonate the system varied from 4 equivalents for **2b**, 3 equivalents for **2a** to 2 equivalents for **2c**. As expected, system **2c** showed the strongest interaction with fluoride ions. Stability constants for the interaction of ligands **2a-c** in the

presence of fluoride ion were calculated using SPECFIT/32 software¹¹ and are summarized in Table 1.

Table 1. Stability constants for ligands **2a-c** in the presence of fluoride ion and some metal cations in acetonitrile

Ligand	Interaction	log <i>K</i>
2a	F ⁻	4.80 ± 5.39E-02
	(2a) ₂ Cu	8.84 ± 1.59E-02
2b	F ⁻	4.68 ± 1.12E-02
	(2b) ₂ Zn	6.73 ± 6.75E-02
2c	F ⁻	5.02 ± 1.15E-02
	(2c) ₂ Cu	9.81 ± 2.09E-02

Recently, Han and co-workers⁷ reported the synthesis and characterization of phenylimidazo-anthraquinone derivatives as selective F⁻ sensors. The highest sensitivity was achieved for the sensor possessing a *p*-nitro group at the phenyl ring (40 equivalents of fluoride ion), which is 20 times higher than the amount needed in the case of our best system **2c**, probably due to the presence of the more efficient electron transfer bithienyl unit.

A great advantage presented by our deprotonated **2a-c** systems is that they are emissive, and at the same time provide a chelating site with recognition effects formed by the deprotonated N from the imidazole ring and the O from the quinone group. Metal ion complexes can be formed with ligands provided with the 9,10-anthraquinone donor-unit with free electron pairs to coordinate to the metal ions.¹² Therefore, we investigated the behavior of compounds **2a-c** in the presence of Zn(II), Cu(II) and Hg(II) cations, after fluoride ion addition. In all cases, a blue shift in the absorption band centered at *ca* 503-505 nm was observed, and the original band, related to the free ligand, centered at 444 nm was restored, suggesting the formation of metal complexes in solution. With increasing addition of metal ions, a plateau was achieved for a 1:2 metal to ligand molar ratio, except for the complex formed by **2a** with Zn(II), where the molar ratio observed was 1:1. (See Figures S15-S19 Supporting information). In Table 1, some stability constants of ligands **2a-c** with Cu(II), Hg(II) and Zn(II) are presented. The strongest interaction was observed for ligand **2a** in the presence of Hg(II). The mononuclear complex formed by **2a** with Zn(II) could be attributed to the great ability of this metal ion to form different complexes with coordination number from 2 to 6.

A CHEQ (Chelation Enhancement of the Quenching) effect in the fluorescence emission for all metals studied, Zn(II), Hg(II) and Cu(II) was observed (Figure 2). This result suggested the involvement of the metal ion with both donor atoms, the O from the quinone ring and the N from the imidazole ring through two units of **2a-c** ligands, in the case of Cu(II) and Hg(II). The quenching effect

could be attributed to an energy transfer quenching of the π^* emissive state through low-lying metal-centered unfilled d-orbitals for Cu(II), and to an intersystem crossing mechanism due to the heavy atom effect for Hg(II).¹³




Figure 2. UV-vis titration of **2a** in CH₃CN with increasing amount of Hg(CF₃SO₃)₂, after addition of [(Bu)₄N]F. (λ_{exc} = 470 nm). Dotted line spectra: free ligand.

In the case of Zn(II), that normally produces a CHEF effect (Chelation Enhancement of Fluorescence),¹⁴ a quenching effect was also seen. Similar effects have been reported for systems bearing a coordinative aromatic nitrogen atom.¹⁵ Previously, two different sensing mechanisms have been described for the anthraquinone chromophore. An inversion of excited luminescence states, $n-\pi^*$ to $\pi-\pi^*$, proposed by Kadararaisamy^{12d} involving the O atom from the quinone ring, and a PET (Photoinduced Electron Transfer) process proposed by Yoshida and co-workers^{12a} involving a N from the imidazole and azacrown rings. In our case involvement of both N and O donor atoms appears to suggest a PET process.

Taking into account the counter-ion used for all metal complexes, the triflate ion (CF₃SO₃⁻) and its non coordinative properties, the hypothesis of 1:2 M:L molar ratio observed is in agreement with the coordinative involvement of these metals (Cu(II) and Hg(II)) with at least four donor-atoms. (See Figure S20 for a schematic representation in supporting information).

In all cases the pink colour of the solution observed upon fluoride addition, changed to yellow or gold upon metal complexation giving an on-off-on, yellow-pink-yellow, reversible colorimetric reaction. (See Figure S14 in supporting information).

In summary, bithienyl-imidazo-anthraquinones **2a-c** were synthesised in good to excellent yields and were evaluated as colorimetric and/or fluorimetric sensors. Selectivity and sensitivity for the fluoride ion was observed for all systems. Moreover, the systems after deprotonation of the imidazole NH upon addition of fluoride ion, proved to be efficient sensors for metal ions such as Zn(II), Hg(II) and Cu(II), suggesting the formation of 1:2 complexes, except for the complex formed by **2a** with Zn(II), with a 1:1 molar ratio.

Acknowledgment We are grateful to the Portuguese Foundation for Science and Technology for financial support through Center of Chemistry (UM) and FCT/UNL REQUIMTE through project POCI/QUI/55519/2004.

Supporting Information Available: Experimental procedures, characterization data including ¹H and ¹³C spectral data of compounds and proton, basic anions, halide and metal ion sensing spectral studies and figures.

§ Centro de Química, Universidade do Minho

‡ REQUIMTE, Universidade Nova de Lisboa

(1) (a) Schmidtchen, F. P.; Berger, M. *Chem. Rev.* **1997**, *97*, 1609-1646. (b) Beer, P. D.; Gale, P. A. *Angew. Chem. Int. Ed. Engl.* **2001**, *40*, 486-516. (c) Gale, P.A., Ed.; Special issue : 35 years of Synthetic Anion Receptor Chemistry. *Coord. Chem. Rev.* **2003**, *240*, 1-2. (d) Bondy, C. R.; Loeb, S.J. *Coord. Chem. Rev.* **2003**, *240*, 77-99. (e) Martínez-Manêz, R.; F. Sancenón; *Chem. Rev.* **2003**, *103*, 4419-4476. (f) Suksai, C.; Tuntulani, T.; *Chem. Soc. Rev.* **2003**, *32*, 192-202. (g) Martínez-Manêz, R.; F. Sancenón; *J. Fluoresc.* **2005**, *15*, 267-285. (h) Martínez-Manêz, R.; F. Sancenón; *Coord. Chem. Rev.* **2006**, *250*, 3081-3093.

(2) (a) Kirk, K. L. *Biochemistry of the Halogens and Inorganic Halides*; Plenum Press: New York, 1991, p 58. (b) Jentsch, T. J. *Curr. Opin. Neurobiol.* **1996**, *6*, 303-310.

(3) (a) Kleerekoper, M. *Endocrinol. Metab. Clin. North Am.* **1998**, *27*, 441. (b) Tomich, J. M.; Wallace, D.; Henderson, K.; Mitchell, K. E.; Radke, G.; Brandt, R.; Ambler, C. A.; Scott, A. J.; Grantham, J.; Sullivan, L.; Iwamoto, T. *Biophys. J.* **1998**, *74*, 256-267.

(4) (a) Jiménez, D.; Martínez-Manêz, R.; Sancenón, F.; Soto, J. *Tetrahedron Lett.* **2002**, *43*, 2823-2825. (b) Jose, D. A.; Kumar, D. K.; Ganguly, B.; Das, Amitava *Org. Lett.* **2004**, *6*, 3445-3448. (c) Wu, F.-y.; Hu, M.-h.; Wu, Y.-m.; Tan, X.-f.; Zhao, Y.-q.; Ji, Z.-j; *Spectrochim. Acta Part A* **2006**, *65*, 633-637.

(5) Miyaji, H.; Sessler, J. L. *Angew. Chem., Int. Ed.* **2001**, *40*, 154-157.

(6) Miyaji, H.; Sato, W.; Sessler, J. L. *Angew. Chem., Int. Ed.* **2000**, *39*, 1777-1780.

(7) Peng, X.; Wu, Y.; Fan, J.; Tian, M.; Han, K. *J. Org. Chem.* **2005**, *70*, 10524-10531.

(8) (a) Raposo, M. M. M.; Kirsch, G. *Tetrahedron* **2003**, *59*, 4891-4899. (b) Raposo, M. M. M.; Fonseca, A. M. C.; Kirsch, G. *Tetrahedron* **2004**, *60*, 4071-4078. (c) Batista, R. M. F.; Costa, S. P. G.; Raposo, M. M. M. *Tetrahedron Lett.* **2004**, *45*, 2825-2828. (d) Costa, S. P. G.; Batista, R. M. F.; Cardoso, P.; Belsley, M.; Raposo, M. M. M. *Eur. J. Org. Chem.* **2006**, *17*, 3938-3946. (e) Raposo, M. M. M.; Sousa, A. M. R. C.; Kirsch, G.; Cardoso, P.; Belsley, M.; Matos Gomes, E.; Fonseca, A. M. C. *Org. Lett.* **2006**, *8*, 3681-3684. (f) Batista, R. M. F.; Costa, S. P. G.; Malheiro, E. L.; Belsley, M.; Raposo, M. M. M. *Tetrahedron* **2007**, *63*, 4258-4265.

(9) (a) Chen, X.; Walthall, D. A.; Brauman, J. I. *J. Am. Chem. Soc.* **2004**, *126*, 12614-12620. (b) Boiocchi, M.; Del Boca, L.; Gómez, D. E.; Fabbri, L.; Licchelli, M.; Monzani, E. *J. Am. Chem. Soc.* **2004**, *126*, 16507-16514. (c) Gómez, D. E.; Fabbri, L.; Licchelli, M.; Monzani, E. *Org. Biomol. Chem.* **2005**, *3*, 1495-1500. (d) Amendola, V.; Boiocchi, M.; Fabbri, L.; Palchetti, A. *Chem. Eur. J.* **2005**, *11*, 120-127. (e) Boiocchi, M.; Del Boca, L.; Gómez, D. E.; Fabbri, L.; Licchelli, M.; Monzani, E. *Chem. Eur. J.* **2005**, *11*, 3097-3104. (f) Gómez, D. E.; Fabbri, L.; Licchelli, M. *J. Org. Chem.* **2005**, *70*, 5717-5720.

(10) Stephens, F. F.; Bower, J. D. *J. Chem. Soc.* **1949**, 2971-2972.

(11) SPECFIT/32 Global Analysis System, v. 3.0 Spectrum Software Associates, Malborough, MA, USA.

(12) (a) Yoshida, K.; Mori, T.; Watanabe, S.; Kawai, H.; Nagamura, T. *J. Chem. Soc. Perkin Trans. 2* **1999**, 393-397. (b) Kampman, B.; Lian, Y.; Klíkel, K. L.; Vecchi, P. A.; Quiring, H. L.; Soh, C. C.; Syes, A. G. *J. Org. Chem.* **2002**, *67*, 3878-3883. (c) Kumar, S. P.; Kaur, S. *Tetrahedron Lett.* **2002**, *43*, 1097-1099. (d) Kadarkaraisamy, M.; Sykes A. G. *Inorg. Chem.* **2006**, *45*, 779-789. (e) Barzegar, M.; Mousavi, M. F.; Khajehsharifi, H.; Shamsipur, M.; Sharghi, S. *IEEE Sensors J.*, **2005**, *5*, 392-397.

(13) (a) Valeur, B. *Molecular Fluorescence. Principles and Applications*; Wiley-VCH: Weinheim, 2002. (b) Rurack, K. *Spect. Acta. A* **2001**, *57*, 2161-2195. (c) Bazzicalupi, C.; Bencini, A.; Bianchi, A.; Borsari, L.; Danesi, A.; Giorgi, C.; Lodeiro, C.; Mariani, P.; Pina, F.; Santarelli, S.; Tamayo, A.; Valtancoli, B. *Dalton Trans.* **2006**, 4000-4010. (d) Tamayo, A.; Lodeiro, C.; Escriche, L.; Casabo, J.; Covelo, B.; González, P. *Inorg. Chem.* **2005**, *44*, 8105-8115. (e) Bazzicalupi, C.; Bencini, A.; Berni, E.; Bianchi, A.; Borsari, L.; Giorgi, C.; Valtancoli, B.; Lodeiro, C.; Lima J. C.; Parola, A. J.; Pina, F. *Dalton Trans.* **2004**, 591-597.

(14) (a) Lodeiro, C.; Parola, A. J.; Pina, F.; Bazzicalupi, C.; Bencini, A.; Bianchi, A.; Giorgi, C.; Masotti, A.; Valtancoli, B. *Inorg. Chem.* **2001**, *40*, 2968-2975. (b) Bazzicalupi, C.; Bencini, A.; Berni, E.; Bianchi, A.; Danesi, A.; Giorgi, C.; Valtancoli, C.; Lodeiro, C.; Lima, J. C.; Pina, F.; Bernardo, M. A. *Inorg. Chem.* **2004**, *43*, 5134-5146. (c) Okamoto, K.; Fukuzumi, S. *J. Am. Chem. Soc.* **2004**, *126*, 13922-13923.

(15) (a) Aucejo, R.; Alarcon, J.; Garcia-España, E.; Llinares, J. M.; Marchin, K. L.; Soriano, C.; Lodeiro, C.; Bernardo, M. A.; Pina, F.; Pina, J.; de Melo, J. S. *Eur. J. Inorg. Chem.* **2005**, *21*, 4301-4308 and references therein. (b) Feng, K.; Hsuy, F.-L.; Bota, K.; Bu, X. R. *Microchem. J.* **2005**, *81*, 23-27.