

19th DRT, Leuven, Bélgica, 14-18 Setembro, 2013

Quantification of synmagmatic flow structures of the Vila Pouca de Aguiar Pluton, NE Portugal

Martin Schwindinger<sup>1</sup>, Mark Peternell<sup>1</sup>, Benedito C. Rodrigues<sup>2</sup>, Jorge Pamplona<sup>3</sup>

<sup>1</sup>Tektonophysik, Johannes-Gutenberg University Mainz, 55099 Mainz, Germany, mschwindinger@web.de

<sup>2</sup>CGUP, Porto University, 4169-007 Porto, Portugal

<sup>3</sup>CIG-R /DCT, Minho University, 4710-057 Braga, Portugal

Methods based on fractal geometry offer the opportunity to quantify complex rock patterns (Kruhl, 2013), which provide information about the pattern forming processes. Mineral distribution patterns of Variscan post-tectonic granites from NW Portugal (Vila Pouca de Aguiar Pluton) were analysed with the MORFA and Map-Counting software (Peternell, 2011). The result of the analysis provides information about pattern inhomogeneity and anisotropy, i.e. magmatic flux directions and mineral equilibrium processes in the crystallizing magma chamber.

The investigated rock is a very homogenous, medium grained biotitic granite, with schlieren structures and mafic enclaves locally occurring. No magmatic foliation or lineation is visible in the rock. For the analysis of rock patterns, high resolution field photographs were taken from six rock surfaces (4.75 - 19 m<sup>2</sup>) and three perpendicular cuts within a quarry. The field photographs were converted to binary images of mineral distribution patterns and analysed with MORFA and Map-Counting. In case of MORFA, 4838 – 20037 single measurements were performed for each rock surface, in case of Map-Counting, 129717 - 533976.

Statistical evaluation of the general very weak pattern anisotropy results in a mean bulk orientation vector that varies in strength, dependent on the orientation of the analysed surface. Therefore, it was possible to determine an extreme weak subhorizontal magmatic foliation (012/15 NW) and a subhorizontal lineation (010/15 NW), which are in good coincidence with earlier anisotropy magnetic susceptibility (AMS) results of Sant'Ovaia & Noronha (2005). With MORFA, also the variation of the lineation can be determined, indicating magmatic flow partitioning scale-dependent into different domains. High variation is observable in the decimetre scale and bulk orientations reveal two perpendicular domains at the scale of several meter. Throughout homogenous results of the Map-Counting analysis with mean  $D_b$ -values of 1.55 indicate equilibrium conditions for mineral crystallization during emplacement of the pluton. Locally, feldspar rich domains show wider  $D_b$ -ranges caused by processes such as magma mixing and extraction.

#### References:

Kruhl, JH, 2013. Fractal-geometry techniques in the quantification of complex rock structures: A special view on scaling regimes, inhomogeneity and anisotropy. *Journal of Structural Geology*, 46:2-21.

Peternell, M, Bitencourt, MF, Kruhl, JH, 2011. Combined quantification of anisotropy and inhomogeneity of magmatic rock fabrics – an outcrop scale analysis recorded in high resolution. *Journal of Structural Geology*, 33:609-623.

Sant'Ovaia, H, Noronha, F, 2005. Classification of Portuguese Hercynian granites based on petrophysical characteristics. *Cadernos Laboratorio Xeolóxico de Laxe*, 30:75-86.