

Poster19

Assessment of ciprofloxacin photocatalysis by-products toxicity with *Vibrio fischeri*

A.R.Silva^{1,2}, S. Teixeira³, P.M. Martins^{1,2}, L.Periera², M. Alves², K.Keuhn³, G. Cuniberti^{3,4,5} and S. Lanceros-Mendez^{1*}

¹Centro/Departamento de Física, Universidade do Minho, 4710-057 Braga, Portugal

²Centro de Engenharia Biológica, Universidade do Minho, 4710-057 Braga, Portugal

³Institute for Materials Science and Max Bergmann Center of Biomaterials, TU Dresden, 01062

⁴Dresden, Germany Dresden Center for Computational Materials Science (DCCMS), TU Dresden, 01062 Dresden, ⁵Center for Advancing Electronics Dresden, TU Dresden, 01062 Dresden, Germany

*Corresponding author: pamartins@fisica.uminho.pt

The presence of pharmaceuticals in water has become a large concern due to the potential negative effects on humans and aquatic ecosystems. From these pharmaceuticals, antibiotics represent a serious problem since their overuse and misuse may lead to adverse environmental effects, in particular, toxicity to microflora and fauna and potential negative effects to humans [1]. Photocatalysis has become attractive to promote the degradation of contaminants in the aquatic environment since it allows their rapid and efficient removal from water, transforming them into by-products [2]. In order to evaluate toxicity of these by-products, several bio tests using bacteria (*Vibrio fischeri*) and algae (*Daphnia spp.*), among others, have been used [3]. In the present work a photocatalytic systems using commercial TiO₂ and ZnO nanoparticles in suspension was used to degrade ciprofloxacin under UV radiation. Samples were withdraw over time in order to monitor degradation and toxicity. The luminescence of the bacteria *Vibrio fischeri* was used to test the toxicity of ciprofloxacin intermediate compounds, produced during the photocatalysis process. If a substance is toxic towards these bacteria, their normal luminescence decreases, as a consequence of a decreasing bacteria viability. Results (Figure 1) indicate that samples without ciprofloxacin degradation (t=0), in contact with bacteria (for 35 min), result in a higher luminescence than with completely degraded ciprofloxacin (t=15min). These results indicate that by products are responsible for low bacteria viability.




Figure 1 – Luminescence of bacteria during 35 minutes of contact with negative control samples and ciprofloxacin degradation samples at t=0 and t=15.

Acknowledgments:

This work was supported by FEDER through the COMPETE Program and by the Portuguese Foundation for Science and Technology (FCT) in the framework of the Strategic Project PEST-C/FIS/UI607/2011 and project PTDC/CTM-NAN/121038/2010. P. M. Martins thanks FCT for grant SFRH/BD/98616/2013.

References:

1. Homem, V. and Santos, L., Journal of Environmental Management, 92 (2011) 2304-2347.
2. Fujishima, A., Rao, T. N., and Tryk, D., Journal of Photochemistry and Photobiology C: 1 (2000) 1–21.
3. M. Heinlaan, A. Ivask, I. Blinova, H.-C. Dubourguier, and A. Kahru, Chemosphere, vol. 71, no. 7, pp. 1308–16, Apr. 2008.