

A note on strong normalization in classical natural deduction

José Espírito Santo
Centro de Matemática
Universidade do Minho
Portugal
jes@math.uminho.pt

In the context of natural deduction for propositional classical logic, with classicality given by the inference rule *reductio ad absurdum*, we investigate the De Morgan translation of disjunction in terms of negation and conjunction. Once the translation is extended to proofs, it obtains a reduction of provability to provability in the disjunction-free subsystem. It is natural to ask whether a reduction is also obtained for, say, strong normalization; that is, whether strong normalization for the disjunction-free system implies the same property for the full system, and whether such lifting of the property can be done along the De Morgan translation. Although natural, these questions are neglected by the literature. We spell out the map of reduction steps induced by the De Morgan translation of proofs. But we need to “optimize” such a map in order to show that a reduction sequence in the full system from a proof determines, in a length-preserving way, a reduction sequence in the disjunction-free system from the De Morgan translation of the proof. In this sense, the above questions have a positive answer.

1 Introduction

In the context of natural deduction for propositional classical logic, with classicality given by the inference rule *reductio ad absurdum* [2], we investigate the De Morgan translation of disjunction in terms of negation and conjunction $A \vee B := \neg(\neg A \wedge \neg B)$. This translation immediately extends to proof-rules, as recalled in Fig. 1, which obtains a reduction of provability to provability in the disjunction-free subsystem.

It is the case that a reduction is also obtained for, say, strong normalization? To be more precise, is it the case that strong normalization for the disjunction-free system implies the same property for the full-system, and that such lifting of the property can be done along the De Morgan translation, that is, through a mapping of reduction sequences? The answer is not clear at all, nevertheless these questions are neglected by the literature. For instance Stålmårck [4] says “(...) *the strong normalization theorem for the restricted version of first order classical N.D. together with the well-known results on the definability of the rules for \vee and \exists in the restricted system does not imply the normalization theorem for the full system*”, and this author moves on to give a direct proof of strong normalization for the full first order system; but the quoted claim is not supported by technical evidence. On the other hand, a claim in the opposite direction, namely that the normalization theorem for the full system does follow from the normalization for the restricted system, may be thought of as the implicit, unproven assumption behind the approach to classical natural deduction in Prawitz [2], where only the restricted system is studied, after being considered “*adequate*”.

In this paper we re-examine these questions, for propositional logic. We spell out the map of reduction steps induced by the De Morgan translation of proofs, and observe that the map does not readily lift strong normalization from the disjunction-free system. We need to “optimize” such map in order to

Figure 1: The de Morgan translation of proofs

$$\begin{array}{c}
\vdots \\
\frac{A_1}{A_1 \vee A_2}
\end{array}
\mapsto
\begin{array}{c}
\frac{[w : \neg A_1 \wedge \neg A_2] \quad \vdots}{\frac{\neg A_i \quad A_i}{\perp}} \\
\frac{\perp}{\neg(\neg A_1 \wedge \neg A_2)} w
\end{array}$$

$$\begin{array}{c}
\vdots \\
\frac{A \vee B \quad \frac{[x:A] \quad [y:B]}{C} \quad x,y}{C}
\end{array}
\mapsto
\begin{array}{c}
\vdots \\
\frac{\neg(\neg A \wedge \neg B) \quad \frac{\frac{[k:\neg C] \quad \frac{\perp}{\neg A} x \quad \frac{[k:\neg C] \quad \frac{\perp}{\neg B} y}{\neg A \wedge \neg B}}{C} k}{\perp}
\end{array}$$

show that a reduction sequence in the full system from a proof determines, in a length-preserving way, a reduction sequence in the disjunction-free system from the De Morgan translation of the proof. In this sense, strong normalization for the full system does indeed reduce to strong normalization for the disjunction-free system along the De Morgan translation. As we provide a proof of strong normalization for the disjunction-free system, this completes a new proof of strong normalization for the full system.

Our natural deduction system is presented as a variant of the λ_Δ -calculus [3], and we employ typed λ -terms throughout to code logical derivations. As to reduction rules, in addition to detour conversion and the commuting conversions pertaining to disjunction, one has conversions related to *reductio ad absurdum inferences* - ρ -conversions, as we will call them. We do not adopt the conversions by Prawitz for the atomization of the conclusion of r.a.a. inferences [2], but rather the conversions of Stålmærck [4] (which go back to Statman [5]), except that we do not impose any constraint on the ρ -conversion for disjunction, in this way following [3].

The paper is organized as follows. In Section 2 we present our natural deduction system and its disjunction-free subsystem. In Section 3 we study the De Morgan translation of proofs, and how it lifts strong normalization from the disjunction-free subsystem to the full system. In Section 4 we prove strong normalization of the disjunction-free subsystem.

2 Background

We present our logical system as a calculus of the λ_Δ family [3]. Types/formulas are given by

$$A, B, C ::= X \mid \perp \mid A \supset B \mid A \wedge B \mid A \vee B$$

We define $\neg A := A \supset \perp$.

Proof terms:

$$\begin{array}{l}
M, N, P, Q ::= x \quad \text{(assumption)} \\
\mid \lambda x^A. M \mid MN \quad \text{(implication)} \\
\mid \langle M, N \rangle \mid \pi_1(M) \mid \pi_2(M) \quad \text{(conjunction)} \\
\mid \text{in}_1(M) \mid \text{in}_2(N) \mid \text{case}(M, x^A. P, y^B. Q) \quad \text{(disjunction)} \\
\mid \Delta k^{-A}. M \quad \text{(reductio ad absurdum)}
\end{array}$$

Figure 2: Typing/inference rules

$$\begin{array}{c}
\frac{}{\Gamma, x : A \vdash x : A} \text{Ass} \\
\\
\frac{\Gamma, x : A \vdash B}{\Gamma \vdash \lambda x^A. M : A \supset B} \supset I \quad \frac{\Gamma \vdash M : A \supset B \quad \Gamma \vdash N : A}{\Gamma \vdash MN : B} \supset E \\
\\
\frac{\Gamma \vdash M : A \quad \Gamma \vdash N : B}{\Gamma \vdash \langle M, N \rangle : A \wedge B} \wedge I \quad \frac{\Gamma \vdash M : A \wedge B}{\Gamma \vdash \pi_1(M) : A} \wedge E1 \quad \frac{\Gamma \vdash M : A \wedge B}{\Gamma \vdash \pi_2(M) : B} \wedge E2 \\
\\
\frac{\Gamma \vdash M : A}{\Gamma \vdash \text{in}_1(M) : A \vee B} \vee I1 \quad \frac{\Gamma \vdash N : A}{\Gamma \vdash \text{in}_2(N) : A \vee B} \vee I2 \\
\\
\frac{\Gamma \vdash M : A \vee B \quad \Gamma, x : A \vdash P : C \quad \Gamma, y : B \vdash Q : C}{\Gamma \vdash \text{case}(M, x^A. P, y^B. Q) : C} \vee E \\
\\
\frac{\Gamma, k : \neg A \vdash M : \perp}{\Gamma \vdash \Delta k^{-A}. M : A} \text{RAA}
\end{array}$$

The type annotation in the bound variable of binders will often be omitted when no confusion arises.

The typing/inference rules are in Fig. 2. Γ denotes a set of *declarations* $x : A$ such that a variable is declared at most one time in Γ .

For the purpose of defining some reduction rules and the translation of proof terms, it is convenient to arrange the syntax of the system in a different way:

$$\begin{array}{ll}
\text{(Terms)} & M, N, P, Q ::= V \mid \mathcal{E}[M] \mid \Delta k^{-A}. M \\
\text{(Values)} & V ::= x \mid \lambda x. M \mid \langle M, N \rangle \mid \text{in}_1(M) \mid \text{in}_2(N) \\
\text{(Elim. contexts)} & \mathcal{E} ::= [-]N \mid \pi_1([-]) \mid \pi_2([-]) \mid \text{case}([-], x^A. P, y^B. Q)
\end{array}$$

A *value* V ranges over terms representing assumptions or introduction inferences. \mathcal{E} stands for an *elimination context*, which is a term representing an elimination inference, but with a “hole” in the position of the main premiss. $\mathcal{E}[M]$ denotes the term resulting from filling the hole of \mathcal{E} with M .

In Fig. 3 one finds the typing rules for contexts. Without surprise [1], these are particular cases of sequent calculus inference rules. In a sequent $\Gamma \mid A \vdash \mathcal{E} : B$, A is the type of the hole of \mathcal{E} and B is the type of the term obtained by filling the hole of \mathcal{E} with a term of type A .

The reduction (or proof transformation) rules are given in Fig. 4. They make use of the following organization of the definition of elimination contexts: $\mathcal{E} ::= \mathcal{E}_{\supset} \mid \mathcal{E}_{\wedge} \mid \mathcal{E}_{\vee}$, where

$$\mathcal{E}_{\supset} ::= [-]N \quad \mathcal{E}_{\wedge} ::= \pi_1([-]) \mid \pi_2([-]) \quad \mathcal{E}_{\vee} ::= \text{case}([-], x^A. P, y^B. Q)$$

For each $R \in \{\beta, \pi, \rho_1, \rho_1^\perp\}$, we put $R := R_{\supset} \cup R_{\wedge} \cup R_{\vee}$. Then we put $\rho := \rho_1 \cup \rho_1^\perp \cup \rho_2$. An equivalent definition of π, ρ_1 and ρ_1^\perp is

$$\begin{array}{lll}
(\pi) & \mathcal{E}[\text{case}(M, x. P, y. Q)] & \rightarrow \text{case}(M, x. \mathcal{E}[P], y. \mathcal{E}[Q]) \\
(\rho_1) & \mathcal{E}[\Delta k. M] & \rightarrow \Delta k'. [\lambda z. k'(\mathcal{E}[z])/k]M \\
(\rho_1^\perp) & \mathcal{E}^\perp[\Delta k. M] & \rightarrow [\lambda z. \mathcal{E}^\perp[z]/k]M
\end{array}$$

Figure 3: Typing for contexts

$$\frac{\Gamma \vdash N : A}{\Gamma | A \supset B \vdash [-]N : B} \quad \frac{}{\Gamma | A_1 \wedge A_2 \vdash \pi_i([-]) : A_i} \quad (i = 1, 2)$$

$$\frac{\Gamma, x : A \vdash P : C \quad \Gamma, y : B \vdash Q : C}{\Gamma | A \vee B \vdash \text{case}([-], x.P, y.Q) : C}$$

$$\frac{\Gamma \vdash M : A \quad \Gamma | A \vdash \mathcal{E} : B}{\Gamma \vdash \mathcal{E}[M] : B}$$

Figure 4: Reduction (or proof transformation) rules

Detour conversion rules:

$$\begin{aligned} (\beta_{\supset}) \quad & (\lambda x.M)N \rightarrow [N/x]M \\ (\beta_{\wedge}) \quad & \pi_i(\langle M_1, M_2 \rangle) \rightarrow M_i \quad (i = 1, 2) \\ (\beta_{\vee}) \quad & \text{case}(\text{in}_i(M), x_1.P_1, x_2.P_2) \rightarrow [M/x_i]P_i \quad (i = 1, 2) \end{aligned}$$

Commutative conversion rule:

$$(\pi_{\circ}) \quad \mathcal{E}_{\circ}[\text{case}(M, x.P, y.Q)] \rightarrow \text{case}(M, x.\mathcal{E}_{\circ}[P], y.\mathcal{E}_{\circ}[Q]) \quad (\circ = \supset, \wedge, \vee)$$

Reductio ad absurdum rules:

$$\begin{aligned} (\rho_{\circ}) \quad & \mathcal{E}_{\circ}[\Delta k.M] \rightarrow \Delta k'. [\lambda z.k'(\mathcal{E}_{\circ}[z])/k]M \quad (\circ = \supset, \wedge, \vee) \\ (\rho_{\circ}^{\perp}) \quad & \mathcal{E}_{\circ}^{\perp}[\Delta k.M] \rightarrow [\lambda z.\mathcal{E}_{\circ}^{\perp}[z]/k]M \quad (\circ = \supset, \wedge, \vee \text{ and } \mathcal{E}_{\circ}^{\perp} : \perp) \\ (\rho_2) \quad & \Delta k.kM \rightarrow M \quad (k \notin M) \end{aligned}$$

The reduction rules preserve types (subject reduction property). The case of ρ_{\supset} is given in Fig. 5, where W , S and $(*)$ are applications of weakening, the typing of substitution, and last rule of Fig. 3, respectively,

Suppose \mathcal{E}_{\circ} in the redex of ρ_{\circ} has (and the reduction happens at) type \perp : in this case, if ρ_{\circ} is applied, the inference $\Delta k'.-$ in the contractum concludes \perp , with $k : \neg \perp$. We separate rules ρ_{\circ}^{\perp} for this case, which avoid such unnecessary uses of RAA. An alternative would have been to adopt another reduction rule

$$(\rho_3) \quad \Delta k^{-\perp}.M \rightarrow [I/k]M \quad (I := \lambda x.x) .$$

Then, ρ_{\circ}^{\perp} would have become a derived rule

$$\begin{aligned} \mathcal{E}_{\circ}^{\perp}[\Delta k.M] & \rightarrow_{\rho_{\circ}} \Delta k'. [\lambda z.k'(\mathcal{E}_{\circ}^{\perp}[z])/k]M \\ & \rightarrow_{\rho_3} [\lambda z.I(\mathcal{E}_{\circ}^{\perp}[z])/k]M \\ & \rightarrow_{\beta_{\supset}^*} [\lambda z.\mathcal{E}_{\circ}^{\perp}[z]/k]M \end{aligned} \quad (1)$$

Still, we do not adopt ρ_3 (in this, we agree with [3]). This is technically convenient, as we point out later.

As we leave out ρ_3 , we might have considered forbidding the formation of $\Delta k.M$ with $k : \neg \perp$, and insisted that rules ρ_{\circ} should only happen at types different from \perp - the latter seems even necessary to

Figure 5: Example of type preservation

$$\begin{array}{c}
\frac{k : \neg(A \supset B), \Gamma \vdash M : \perp}{\Gamma \vdash \Delta k.M : A \supset B} \mathcal{D}_1 \quad \text{RAA} \quad \frac{\Gamma | A \supset B \vdash \mathcal{E}_\supset : B}{\Gamma \vdash \mathcal{E}_\supset[\Delta k.M] : B} \mathcal{D}_2 \quad (*) \\
\downarrow \\
\frac{\frac{\Delta \vdash k' : \neg B}{\Delta \vdash k'(\mathcal{E}_\supset[z]) : \perp} \text{Ass} \quad \frac{\frac{\Delta \vdash z : A \supset B}{\Delta \vdash \mathcal{E}_\supset[z] : B} \text{Ass} \quad \frac{\Gamma | A \supset B \vdash \mathcal{E}_\supset : B}{\Delta | A \supset B \vdash \mathcal{E}_\supset : B} \mathcal{D}_2}{\Delta \vdash k'(\mathcal{E}_\supset[z]) : \perp} \text{W} \quad (*)}{\frac{k' : \neg B, \Gamma \vdash \lambda z.k'(\mathcal{E}_\supset[z]) : \neg(A \supset B)}{k' : \neg B, \Gamma \vdash [\lambda z.k'(\mathcal{E}_\supset[z])/k]M : \perp} \text{S}}{\Gamma \vdash \Delta k'.[\lambda z.k'(\mathcal{E}_\supset[z])/k]M : B} \text{RAA}
\end{array}$$

where $\Delta := z : A \supset B, k' : \neg B, \Gamma$.

guarantee the confluence property. Since we only care about strong normalization, we also refrain from enforcing such restrictions.

Later, we will consider another reduction relation, generated by the rule

$$(\rho_4) \quad k'(\Delta k.M) \rightarrow [k'/k]M .$$

This will be just a technical device to help reasoning. This rule does not belong to the system.

If R is a(n union of) reduction rule(s)¹, then \rightarrow_R denotes R -reduction, the binary relation inductively generated by closing R under all contexts. Then \rightarrow_R^- (resp. \rightarrow_R^+ , \rightarrow_R^*) denotes the reflexive (resp. transitive, reflexive-transitive) closure of \rightarrow_R , whereas the latter's composition with itself $n - 1$ times is denoted \rightarrow_R^n , as usual. $=_R$ is a coarser notion of equality over proof expressions than “syntactic” identity. The later is denoted by $=$ (sometimes by \equiv for emphasis) and coincides with α -equivalence, since we work modulo the name of bound variables.

Definition 1 (Full, disjunction-free, and small systems).

1. The system defined so far is denoted λ_Δ : it is a presentation of propositional classical natural deduction. We also refer to this system as the full system.
2. By $\lambda_\Delta^{-\vee}$ we denote the restriction of λ_Δ obtained by leaving out disjunction. More precisely, in $\lambda_\Delta^{-\vee}$ we omit: type $A \vee B$ and its term constructions and typing rules; and the reduction rules β_\vee , π_\circ , and Δ_\vee . In $\lambda_\Delta^{-\vee}$, $\beta := \beta_\supset \cup \beta_\wedge$, $\rho_1 := \rho_\supset \cup \rho_\wedge$, and $\rho_1^\perp := \rho_\supset^\perp \cup \rho_\wedge^\perp$. $\lambda_\Delta^{-\vee}$ is also called the disjunction-free system.
3. The small system, denoted $\lambda_\Delta^{-\vee\wedge}$, is obtained from $\lambda_\Delta^{-\vee}$ by leaving out conjunction, with implication and absurdity the logical constants remaining. In $\lambda_\Delta^{-\vee\wedge}$, $\beta := \beta_\supset$, and $\rho_1 := \rho_\supset$. Furthermore, ρ_\supset^\perp is omitted, hence there is no ρ^\perp in this system.

¹We will often denote (long) unions by juxtaposition, e.g. $R_1R_2R_3$ instead of $R_1 \cup R_2 \cup R_3$.

Figure 6: The de Morgan translation of proof expressions

$$\begin{aligned}
x^\diamond &= x \\
(\lambda x.M)^\diamond &= \lambda x.M^\diamond \\
\langle M, N \rangle^\diamond &= \langle M^\diamond, N^\diamond \rangle \\
(\text{in}_i(M))^\diamond &= \lambda w.\pi_i(w)M^\diamond && (i = 1, 2) \\
(\mathcal{E}_\circ[M])^\diamond &= \mathcal{E}_\circ^\diamond[M^\diamond] && (\circ = \supset, \wedge) \\
(\text{case}(M, x.P, y.Q))^\diamond &= \begin{cases} M^\diamond \langle \lambda x.P^\diamond, \lambda y.Q^\diamond \rangle & (P, Q : \perp) \\ \Delta k.M^\diamond \langle \lambda x.kP^\diamond, \lambda y.kQ^\diamond \rangle & \text{otherwise} \end{cases} \\
(\Delta k.M)^\diamond &= \Delta k.M^\diamond \\
([_]N)^\diamond &= [_]N^\diamond \\
(\pi_i([_]))^\diamond &= \pi_i([_])
\end{aligned}$$

The full system will be the source of the De Morgan translation, whose target is the disjunction-free system. Strong normalization for our particular disjunction-free system is not a result found off the shelf - it will be proved as a corollary to strong normalization of the small system, the latter being a known result:

Theorem 1. *The small system $\lambda_\Delta^{-\vee \wedge}$ enjoys strong normalization.*

Proof. $\lambda_\Delta^{-\vee \wedge}$ is system $\lambda_\Delta^{\perp, \supset}$ of [3] minus its reduction rule (4), so Theorem 23 of *op. cit.* applies. ■

3 Lifting of strong normalization

3.1 The De Morgan translation

We now present the translation into the disjunction-free sub-system. The *de Morgan translation* consists of the obvious translation of formulas

$$\begin{aligned}
X^\diamond &= X \\
\perp^\diamond &= \perp \\
(A \supset B)^\diamond &= A^\diamond \supset B^\diamond \\
(A \wedge B)^\diamond &= A^\diamond \wedge B^\diamond \\
(A \vee B)^\diamond &= \neg(\neg A^\diamond \wedge \neg B^\diamond)
\end{aligned}$$

together with a translation of proof expressions given in Fig. 6. The translation is defined homomorphically in all cases but those relative to the constructors representing introduction and elimination of disjunction; and, for these, the translation corresponds to the transformations recalled in Fig. 1.

Notice that the target system of this translation is classical, therefore the translation is not a negative/CPS translation.

The case distinction in the definition of $(\text{case}(M, x.P, y.Q))^\diamond$ means that we can only apply the translation of proof expressions to typed expressions, M say, and in that case M^\diamond has the same type as M .

Proposition 1 (Soundness). *The typing rules of Fig. 7 are derivable.*

Figure 7: Typing for the de Morgan translation

$$\frac{\Gamma \vdash M : A}{\Gamma^\diamond \vdash M^\diamond : A^\diamond} \quad \frac{\Gamma | A \vdash \mathcal{E}_\circ : B}{\Gamma^\diamond | A^\diamond \vdash \mathcal{E}_\circ : B^\diamond} \quad \circ = \supset, \wedge$$

The first of the typing rules in Fig. 7 states the logical soundness of the translation: if A is a theorem with proof M in the source system, then A^\diamond is a theorem with proof M^\diamond in the disjunction-free system.

Lemma 1 (Preservation of substitution). $[N^\diamond/x]M^\diamond = ([N/x]M)^\diamond$.

Proof. We need in λ_Δ the obvious concept $[Q/x]\mathcal{E}$:

$$\begin{aligned} [Q/x]([-]N) &= [-][Q/x]N \\ [Q/x](\pi_i([-])) &= \pi_i([-]) \\ [Q/x](\text{case}([-], x_1.P_1, x_2.P_2)) &= \text{case}([-], x_1.[Q/x]P_1, x_2.[Q/x]P_2) \end{aligned}$$

Then the following holds

$$[Q/x](\mathcal{E}[M]) = ([Q/x]\mathcal{E})[[Q/x]M] \quad (*)$$

Now, the lemma's statement is proved together with $[N^\diamond/x]\mathcal{E}_\circ^\diamond = ([N/x]\mathcal{E}_\circ)^\diamond$, with $\circ = \supset, \wedge$, by simultaneous induction on M and \mathcal{E}_\circ . All cases follow by definitions, IHs, and fact (*). ■

Theorem 2 (Translation of proof reduction).

1. Let $R \in \{\beta_\supset, \beta_\wedge, \rho_\supset, \rho_\supset^\perp, \rho_\wedge, \rho_\wedge^\perp, \rho_2\}$ and let $R^\diamond = R$. If $M \rightarrow_R N$ in λ_Δ , then $M^\diamond \rightarrow_{R^\diamond} N^\diamond$ in $\lambda_\Delta^{-\vee}$.
2. Let $R = \rho_\vee^\perp$ and let $R^\diamond = \rho_\supset^\perp$. If $M \rightarrow_R N$ in λ_Δ , then $M^\diamond \rightarrow_{R^\diamond} N^\diamond$ in $\lambda_\Delta^{-\vee}$.
3. Let $R = \beta_\vee$ (resp. $R = \pi_\supset, R = \pi_\wedge$) and let $R^\diamond = \beta\rho_2$ (resp. $R^\diamond = \beta_\supset\rho_\supset, R^\diamond = \beta_\supset\rho_\wedge$). If $M \rightarrow_R N$ in λ_Δ , then $M^\diamond \rightarrow_{R^\diamond}^+ N^\diamond$ in $\lambda_\Delta^{-\vee}$.
4. Let $R \in \{\rho_\vee, \rho_\vee^\perp\}$ and $R^\diamond = \rho_\supset^\perp$. If $M \rightarrow_R N$ in λ_Δ then $M^\diamond \rightarrow_{R^\diamond} N^\diamond$ in $\lambda_\Delta^{-\vee}$, for some N' such that $N^\diamond \rightarrow_{\rho_4} N'$.

Proof. By induction on $M \rightarrow_R N$. First we see the base cases.

Case β_\supset . Let $LHS := (\lambda x.M)N \rightarrow [N/x]M =: RHS$.

$$\begin{aligned} LHS^\diamond &= (\lambda x.M^\diamond)N^\diamond \quad (\text{by def.}) \\ &\rightarrow_{\beta_\supset} [N^\diamond/x]M^\diamond \\ &= ([N/x]M)^\diamond \quad (\text{by Lemma 1}) \\ &= RHS^\diamond \end{aligned}$$

Case ρ_\circ . Let $LHS := \mathcal{E}_\circ[\Delta k.M] \rightarrow \Delta k'[\lambda z.k'(\mathcal{E}_\circ[z])/k]M =: RHS$, with $\circ = \supset, \wedge$.

$$\begin{aligned} LHS^\diamond &= \mathcal{E}_\circ^\diamond[\Delta k.M^\diamond] \quad (\text{by def.}) \\ &\rightarrow_{\rho_\circ} \Delta k'.[\lambda z.k'(\mathcal{E}_\circ^\diamond[z])/k]M^\diamond \\ &= \Delta k'.([\lambda z.k'(\mathcal{E}_\circ[z])]^\diamond/k)M^\diamond \quad (\text{by def.}) \\ &= \Delta k'.([\lambda z.k'(\mathcal{E}_\circ[z])/k]M)^\diamond \quad (\text{by Lemma 1}) \\ &= RHS^\diamond \quad (\text{by def.}) \end{aligned}$$

The other base cases of R in statement 1 are equally straightforward.

Case $\rho_{\perp}^{\downarrow}$. Let $LHS := \text{case}(\Delta k.M, x.P, y.Q) \rightarrow [\lambda z.\text{case}(z, x.P, y.Q)/k]M =: RHS$. Let $N := \langle \lambda x.P^{\diamond}, \lambda y.Q^{\diamond} \rangle$.

Then:

$$\begin{aligned} LHS^{\diamond} &= (\Delta k.M^{\diamond})N && \text{(by def.)} \\ &\rightarrow_{\rho_{\perp}^{\downarrow}} [\lambda z.zN/k]M^{\diamond} \\ &= [(\lambda z.\text{case}(z, x.P, y.Q))^{\diamond}/k]M^{\diamond} && \text{(by def.)} \\ &= RHS^{\diamond} && \text{(by Lemma 1)} \end{aligned}$$

Case β_{\vee} . Let $LHS := \text{case}(\text{in}_i(M), x_1.P_1, x_2.P_2) \rightarrow [M/x_i]P_i =: RHS$.

$$\begin{aligned} LHS^{\diamond} &= \Delta k.(\lambda w.\pi_i(w)M^{\diamond})\langle \lambda x_1.kP_1^{\diamond}, \lambda x_2.kP_2^{\diamond} \rangle && \text{(by def.)} \\ &\rightarrow_{\beta}^3 \Delta k.k([M^{\diamond}/x_i]P_i^{\diamond}) \\ &\rightarrow_{\rho_2} [M^{\diamond}/x_i]P_i^{\diamond} \\ &= ([M/x_i]P_i)^{\diamond} && \text{(by Lemma 1)} \\ &= RHS^{\diamond} \end{aligned}$$

Case π_{\circ} . Let $LHS := \mathcal{E}_{\circ}[\text{case}(M, x.P, y.P)] \rightarrow \text{case}(M, x.\mathcal{E}_{\circ}[P], y.\mathcal{E}_{\circ}[Q]) =: RHS$, with $\circ = \supset, \wedge$.

$$\begin{aligned} LHS^{\diamond} &= \mathcal{E}_{\circ}^{\diamond}[\Delta k.M^{\diamond}\langle \lambda x.kP^{\diamond}, \lambda y.kQ^{\diamond} \rangle] && \text{(by def.)} \\ &\rightarrow_{\rho_{\circ}} \Delta k'[\lambda z.k'(\mathcal{E}_{\circ}^{\diamond}[z])/k](M^{\diamond}\langle \lambda x.kP^{\diamond}, \lambda y.kQ^{\diamond} \rangle) \\ &= \Delta k'.M^{\diamond}\langle \lambda x.(\lambda z.k'(\mathcal{E}_{\circ}^{\diamond}[z]))P^{\diamond}, \lambda y.(\lambda z.k'(\mathcal{E}_{\circ}^{\diamond}[z]))Q^{\diamond} \rangle \\ &\rightarrow_{\beta_{\circ}}^2 \Delta k'.M^{\diamond}\langle \lambda x.k'(\mathcal{E}_{\circ}^{\diamond}[P^{\diamond}]), \lambda y.k'(\mathcal{E}_{\circ}^{\diamond}[Q^{\diamond}]) \rangle \\ &= RHS^{\diamond} && \text{(by def.)} \end{aligned}$$

Case ρ_{\vee} . Let $LHS := \text{case}(\Delta k.M, x.P, y.Q) \rightarrow \Delta k'[\lambda z.k'\text{case}(z, x.P, y.Q)/k]M =: RHS$. Let $N := \langle \lambda x.k'P^{\diamond}, \lambda y.k'Q^{\diamond} \rangle$. Then:

$$\begin{aligned} LHS^{\diamond} &= \Delta k'.(\Delta k.M^{\diamond})N && \text{(by def.)} \\ &\rightarrow_{\rho_{\perp}^{\downarrow}} \Delta k'.[\lambda z.zN/k]M^{\diamond} \end{aligned}$$

On the other hand:

$$\begin{aligned} RHS^{\diamond} &= \Delta k'.([\lambda z.k'\text{case}(z, x.P, y.Q)/k]M)^{\diamond} && \text{(by def.)} \\ &= \Delta k'.([\lambda z.k'\text{case}(z, x.P, y.Q)]^{\diamond}/k)M^{\diamond} && \text{(by Lemma 1)} \\ &= \Delta k'.[\lambda z.k'(\Delta k.z\langle \lambda x.kP^{\diamond}, \lambda y.kQ^{\diamond} \rangle)/k]M^{\diamond} && \text{(by def.)} \\ &\rightarrow_{\rho_4} \Delta k'.[\lambda z.[k'/k]z\langle \lambda x.kP^{\diamond}, \lambda y.kQ^{\diamond} \rangle/k]M^{\diamond} \\ &= \Delta k'.[\lambda z.zN/k]M^{\diamond} \end{aligned}$$

Case π_{\vee} is proved by a similar argument.

As to inductive cases, it suffices to say that all the relations that hold between M^{\diamond} and N^{\diamond} , namely

- $\cdot \rightarrow_R \cdot$
- $\cdot \rightarrow_R^{\dagger} \cdot$
- $\exists N(\cdot \rightarrow_R N \wedge \cdot \rightarrow_{\rho_4} N)$

are congruences (i.e. compatible with the syntactic formation operations). So all the inductive cases follow routinely by induction hypothesis. ■

3.2 Optimization

Statement 4 of Theorem 2 is an obstacle for the ready lifting of strong normalization. We now overcome this obstacle.

Lemma 2 (Commutation of reduction steps). *In λ_Δ :*

1. *Let R be a reduction rule different from ρ_2 . If $M \rightarrow_{\rho_4} N_1$ and $M \rightarrow_R N_2$, then there is N_3 such that $N_1 \rightarrow_R N_3$ and $N_2 \rightarrow_{\rho_4}^* N_3$.*
2. *If $M \rightarrow_{\rho_4} N_1$ and $M \rightarrow_{\rho_2} N_2$, then there is N_3 such that: (i) $N_1 \rightarrow_{\rho_2} N_3$ or $N_1 = N_3$; and (ii) $N_2 \rightarrow_{\rho_4}^* N_3$.*


Proof. There is only one case where a ρ_4 -redex overlaps non-trivially with another R -redex, which is when $R = \rho_2$ and $M = k'(\Delta k.kM)$, with $k \notin M$. In this case, take $N_3 = k'M$. ■

If s is a reduction sequence, let $|s|$ denote its length, that is the number of reduction steps in s .

Theorem 3 (Translation of reduction sequences). *If s is a reduction sequence in λ_Δ from M to N , then there is N' such that:*

1. *There is a reduction sequence s' in $\lambda_\Delta^{-\vee}$ from M^\diamond to N' , and $|s'| \geq |s| - m$, where m is the number of ρ_2 -reduction steps in s .*
2. $N^\diamond \rightarrow_{\rho_4}^* N'$.

Proof. By induction on $|s|$. The base case $|s| = 0$ is trivial, just take $N' = N^\diamond$. The inductive case follows from this diagram (where double-headed arrows denote \rightarrow^*):



If any other item of Theorem 2 applies instead, then $M^\diamond \rightarrow^+ P^\diamond$, and Lemma 2 is not needed. ■

This theorem says that a reduction sequence in the full system from a proof M determines, essentially in a length-preserving way, a reduction sequence in the disjunction-free system from the De Morgan translation M^\diamond of the proof.

Corollary 1 (Lifting of strong normalization). *If the disjunction-free system enjoys strong normalization, so does the full system.*

Proof. Suppose there is an infinite reduction sequence from typable M in the full system λ_Δ . Thanks to Proposition 1, $(\cdot)^\blacklozenge$ preserves typability, so M^\blacklozenge is typable. We prove that, for any n , there is in $\lambda_\Delta^{-\vee}$ a reduction sequence s' from M^\blacklozenge of length n . The existence of an infinite reduction sequence from M^\blacklozenge then follows by König's Lemma. Let n be given. Then, there is an initial segment s of the infinite reduction sequence from M such that $|s| - m \geq n$, where m is the number of ρ_2 -reduction steps in s . That such s exists follows from termination of ρ_2 -reduction. From the previous theorem, there is in $\lambda_\Delta^{-\vee}$ a reduction sequence s' from M^\blacklozenge such that $|s'| \geq |s| - m$. ■

This is the positive answer to the main question raised in this paper.

4 Strong normalization

We confirm that the target of the De Morgan translation is a system enjoying strong normalization.

Since we want to lift strong normalization from $\lambda_\Delta^{-\vee\wedge}$ (Theorem 1), we have to translate $\lambda_\Delta^{-\vee}$ into $\lambda_\Delta^{-\vee\wedge}$, i.e., we have to get rid of conjunction. Following [3], we use the map $(\cdot)^\dagger$ of formulas induced by

$$(A \wedge B)^\dagger = \neg(A^\dagger \supset \neg B^\dagger) .$$

The map of proofs is defined by

$$\begin{aligned} \langle M, N \rangle^\dagger &= \lambda f. f M^\dagger N^\dagger \\ \pi_i(M)^\dagger &= \Delta k. M^\dagger (\lambda x_1 x_2. k x_i) \\ \pi_i(M)^\dagger &= M^\dagger (\lambda x_1 x_2. x_i) \quad (M : A_1 \wedge A_2, \quad A_i = \perp) \end{aligned}$$

with the remaining cases defined homomorphically. It turns out that the study of this map follows the same patterns as that of $(\cdot)^\blacklozenge$. We will be very brief now.

Theorem 4 (Translation of proof reduction). *Let $S := \lambda_\Delta^{-\vee\wedge} + \rho_5^\perp$.*

1. *Let R be a reduction rule of $\lambda_\Delta^{-\vee}$ different from ρ_\wedge . If $M \rightarrow_R N$ in $\lambda_\Delta^{-\vee}$, then $M^\dagger \rightarrow^+ N^\dagger$ in S .*
2. *If $M \rightarrow_{\rho_\wedge} N$ in $\lambda_\Delta^{-\vee}$ then $M^\dagger \rightarrow_{\rho_5^\perp} N^\dagger$ in S , for some N' such that $N^\dagger \rightarrow_{\rho_4} N'$.*

Proof. By induction on $M \rightarrow_R N$. ■

Why do we temporarily need reduction rule ρ_5^\perp in the target of $(\cdot)^\dagger$? Not only because the rule exists in the source calculus $\lambda_\Delta^{-\vee}$, but also because it is needed to map ρ_\wedge and ρ_\wedge^\perp , in the same way as it happened before with $(\cdot)^\blacklozenge$ and rules ρ_\vee and ρ_\vee^\perp (recall items 2 and 4 of Theorem 2).

Before, we have used Lemma 2 to obtain Theorem 3 from Theorem 2. Exactly in the same way, we use Lemma 2 to obtain from Theorem 4 the following: if s is a reduction sequence from M in $\lambda_\Delta^{-\vee}$, then there is a reduction sequence s' in $\lambda_\Delta^{-\vee\wedge} + \rho_5^\perp$ from M^\dagger with length $|s'| \geq |s| - m$, where m is the number of ρ_2 -reduction steps in s . Now, ρ_5^\perp becomes a derived rule, if we accept (temporarily) rule ρ_3 (recall (1)). Hence, we can rephrase the result just obtained:

Theorem 5 (Translation of reduction sequences). *Let $S := \lambda_\Delta^{-\vee\wedge} + \rho_3$. If s is a reduction sequence from M in $\lambda_\Delta^{-\vee}$, then there is a reduction sequence s' in S from M^\dagger with length $|s'| \geq |s| - m$, where m is the number of ρ_2 -reduction steps in s .*

Rule ρ_3 makes a timely appearance, now that the commutation argument of Lemma 2 is no longer needed. If the rule were present from the beginning, the commutation of ρ_3 and ρ_4 would fail, and the on-going strategy would be wrong.

Corollary 2 (Lifting of strong normalization). *If $\lambda_{\Delta}^{-\vee\wedge} + \rho_3$ enjoys strong normalization, so does $\lambda_{\Delta}^{-\vee}$.*

Proof. From the previous theorem, termination of ρ_2 -reduction, and preservation of typability by $(\cdot)^{\dagger}$. ■

Since strong normalization of $\lambda_{\Delta}^{-\vee\wedge}$ (Theorem 1) does not comprehend ρ_3 , the last task is to get rid of this rule. This is done by postponement.

Lemma 3 (Postponement). *In $\lambda_{\Delta}^{-\vee\wedge} + \rho_3$, let κ be the rule $kN \rightarrow N$ at type \perp .*

1. *If $M \rightarrow_{\rho_3} P \rightarrow_R Q$, then there is P' such that $M \rightarrow_{R'} P' \rightarrow_{\rho_3}^* Q$, where $R' = R$, except in the case $R = \beta_{\supset}$, of the particular form $IN \rightarrow N$ at type \perp , for which R' may be R or κ .*
2. *If $M \rightarrow_{\kappa} P \rightarrow_R Q$, then there is P' such that $M \rightarrow_R P' \rightarrow_{\kappa}^* Q$*

Proof. The exception mentioned in item 1 occurs when $M = \Delta k^{-\perp}.M'$ and $P = [I/k]M'$, with Q resulting from the reduction of some IN created by substitution $[I/k]_{\perp}$. Let M'' be the result of applying in M' , to the relevant occurrence of k , the rule $kN \rightarrow N$. Put $P' = \Delta k.M''$. ■

Corollary 3 (Lifting of strong normalization). *If $\lambda_{\Delta}^{-\vee\wedge}$ enjoys strong normalization, so does $\lambda_{\Delta}^{-\vee\wedge} + \rho_3$.*

Proof. Suppose there is an infinite reduction sequence from M in $\lambda_{\Delta}^{-\vee\wedge} + \rho_3$. We prove that, for any n , there is in $\lambda_{\Delta}^{-\vee\wedge}$ a reduction sequence s' from M of length n . Let n be given. Let ι be the rule $IN \rightarrow N$ at type \perp . There is an initial segment s of the infinite reduction sequence from M such that $|s| - m - i \geq n$, where m is the number of ρ_3 -reduction steps in s and i is the number of ι -reduction steps in s . That such s exists follows from termination of $(\rho_3 \cup \iota)$ -reduction. By applying postponement of ρ_3 (item 1 of Lemma 3) to s , one obtains a reduction sequence s' from M , without ρ_3 -steps, of length $|s| - m$, where some ι -steps are converted into κ -steps. The number of steps in s' which are not κ -steps is a number $n' \geq |s| - m - i$. By postponement of κ (item 2 of Lemma 3) applied to s , we get a reduction sequence s'' from M , without κ -steps, of length n' . This is a reduction in $\lambda_{\Delta}^{-\vee\wedge}$ with length $\geq n$. ■

Theorem 6. *The disjunction-free system $\lambda_{\Delta}^{-\vee}$ enjoys strong normalization.*

Proof. From Corollaries 2 and 3 and Theorem 1. ■

Theorem 7. *The full system λ_{Δ} enjoys strong normalization.*

Proof. From Corollary 1 and Theorem 6. ■

Acknowledgments: This research was financed by Portuguese Funds through FCT Fundação para a Ciência e a Tecnologia, within the Project UID/MAT/00013/2013.

References

- [1] P.-L. Curien & H. Herbelin (2000): *The duality of computation*. In: *Proceedings of the Fifth ACM SIGPLAN International Conference on Functional Programming (ICFP '00)*, Montreal, Canada, September 18-21, 2000, SIGPLAN Notices 35(9), ACM, pp. 233–243, doi:10.1145/351240.351262.
- [2] D. Prawitz (1965): *Natural Deduction. A Proof-Theoretical Study*. Almqvist and Wiksell, Stockholm.
- [3] N. Rehof & M. Sorensen (1994): *The λ_{Δ} -calculus*. In: *TACS'94, Lecture Notes in Computer Science 789*, Springer Verlag, pp. 516–542, doi:10.1007/3-540-57887-0_113.
- [4] G. Stalmarck (1991): *Normalization theorems for full first order classical natural deduction*. *The Journal of Symbolic Logic* 56(1), pp. 129–149, doi:10.2307/2274910.
- [5] R. Statman (1974): *Structural complexity of proofs*. Ph.D. thesis, Stanford University.