

Molecular profiling, including *TERT* promoter mutations, of acral lentiginous melanomas

Vinicius de Lima Vazquez^{a,b}, Anna L. Vicente^a, Adriana Carloni^a, Gustavo Berardinelli^a, Paula Soares^d, Cristovam Scapulatempo^{a,c}, Olga Martinho^{a,e,f} and Rui M. Reis^{a,e,f}

Acral lentiginous melanoma (ALM) is the less common subtype with singular characterization. *TERT* (human telomerase reverse transcriptase) promoter mutations have been described as recurrent in melanomas and infrequent in ALM, but their real incidence and clinical relevance is unclear. The objectives of this study were to describe the prevalence of *TERT* promoter mutations in ALM, and correlate with the molecular profile of other drive genes and clinical features. Sixty-one samples from 48 patients with ALM were analyzed. After DNA isolation, the mutation profiles of the hotspot region of *BRAF*, *NRAS*, *KIT*, *PDGFRA*, and *TERT* genes were determined by PCR amplification followed by direct Sanger sequencing. *KIT*, *PDGFRA*, and *VEGFR2* gene amplification was performed by quantitative PCR. Clinical information such as survival, clinical stage, and Breslow tumor classification were obtained from medical records. *TERT* promoter mutations were found in 9.3% of the cases, *BRAF* in 10.3%, *NRAS* in 7.5%, *KIT* in 20.7%, and *PDGFRA* in 14.8% of ALM. None of the cases showed *KIT*, *PDGFRA*, or *VEGFR2* gene amplification. We found an association between *KIT* mutations and advanced Clark level (IV and V, $P=0.043$) and *TERT* promoter mutations with low mitotic index. No other significant

associations were observed between mutation profile and patients' clinical features nor survival rates. Oncogenic *TERT* promoter mutations are present in a fraction of ALMs. No relevant associations were found between *TERT* mutation status and clinical/molecular features nor survival. Mutations of *KIT* and *PDGFRA* are the most common genetic alterations, and they can be therapeutic targets for these patients. *Melanoma Res* 26:93–99 Copyright © 2016 Wolters Kluwer Health, Inc. All rights reserved.

Melanoma Research 2016, 26:93–99

Keywords: acral lentiginous, melanoma, molecular profile, *TERT* promoter mutation

^aMolecular Oncology Research Center, ^bDepartment of Surgery, Melanoma and Sarcoma Unity, ^cDepartment of Pathology, Barretos Cancer Hospital, Barretos, São Paulo, Brazil, ^dInstitute of Pathology and Molecular Immunology of University of Porto, (IPATIMUP), Porto, ^eLife and Health Sciences Research Institute (ICVS), Health Sciences School, University of Minho and ^fICVS/3B's-PT Government Associate Laboratory, Braga, Portugal

Correspondence to Vinicius de Lima Vazquez, MD, PhD, Molecular Oncology Research Center, Barretos Cancer Hospital, Rua Antenor Duarte Villela, 1331, CEP 14784 400, Barretos, São Paulo, Brazil
Tel/fax: +55 173 321 6600; e-mails: viniciusvazquez@gmail.com

Received 21 May 2015 Accepted 6 November 2015

Introduction

Cutaneous melanoma is the most aggressive skin cancer, with a rising incidence worldwide in the past few decades [1]. Different from the main melanoma subtypes, usually diagnosed in white populations with fair skin and hair, the acral lentiginous melanoma (ALM) is an infrequent counterpart, accounting for less than 2% of all melanoma cases [2,3]. This subtype exhibits unique clinical characteristics, affecting only the palms of the hands, the soles of the feet, and the subungual area [4,5]. In regions with vast contingent of non-white population, such as Asia, Africa, and part of America, ALM is a major concern, as it occurs in populations and skin areas that are not generally at risk for skin cancer, leading frequently to late diagnosis and low survival rates [6–8]. In Brazil, a country with high miscegenated and heterogeneous population with African, European and indigenous heritage, ALM is prevalent in some geographic areas [9–11].

Over the past few years we have witnessed a fast growth in molecular characterization of melanomas, with great emphasis in alterations of cell signaling pathways [12,13]. The mitogen-activated protein kinase pathway has a crucial role in melanoma development and progression [12,14]. *BRAF*, an intracellular serine-threonine kinase, is mutated in ~60% of cutaneous melanomas [12,15,16]. Ninety-five percent of such cases result in a V600E mutation, which involves a valine to glutamic acid substitution at position 600, which leads to a constitutive activation of the mitogen-activated protein kinase pathway [12,14,17]. Mutations are also frequent in another intracellular kinase, particularly in the *NRAS* gene, with a rate of 15–18% of mutated cases [12,14,17]. It has been reported that ALM exhibited a distinct molecular profile when compared with cutaneous melanoma [18]. In the ALM subtype, *BRAF* mutations are uncommon, and mutations were identified in upstream receptor tyrosine kinase, such as *KIT* (15–30%) and *PDGFRA* (6.8%) [12,14,17]. It was also reported that *KIT* and *PDGFRA*

All supplementary digital content is available directly from the corresponding author.

could be altered in ALM by gene amplification mechanisms of their 4p12 locus [18–20]. Besides these two oncogenes, this locus also harbors another important oncogene, *KDR*, also known as *VEGFR2* (vascular endothelial growth factor receptor-2 gene) [21,22]. Importantly, many of the mutated molecules in ALM have emerged for therapeutic targets, such as *BRAF*, *KIT*/*PDGFR*, and *VEGFR2* [23–25].

Recently, another important cancer-related gene, telomerase reverse transcriptase (*TERT*), was described to be mutated in melanomas [26,27]. *TERT* encodes a subunit of telomerase that, together with other components, elongates the telomere maintaining genomic integrity. Its upregulation has been demonstrated in several human cancers, and the promoter region of the gene is considered the most regulatory element for telomerase expression. *TERT* promoter mutations, namely at positions c. –124 C>T and c. –146 C>T, have been reported in up to 50% of cutaneous melanoma [26]. In ALM, the data are scarce, with few studies addressing its biological and clinical impacts [28–30].

In the present study, we intend to determine the frequency of hotspot *TERT* promoter mutations in a series of ALM melanomas. Furthermore, we aimed to assess other major molecular features, such as *BRAF*, *NRAS*, *KIT*, *PDGFRA* mutations, as well as amplification of 4q12 locus (*KIT*/*PDGFRA*/*VEGFR2*), and correlate with *TERT* mutation status and with ALM clinicopathological features.

Materials and methods

Sixty-one formalin-fixed paraffin-embedded tissues from 48 ALM patients were retrieved from the files of the Department of Pathology at Barretos Cancer Hospital. All the patients were diagnosed between 1999 and 2013. Primary tumor samples were obtained from 48 cases. Eleven cases also presented lymph node (LN) samples and in two cases we assessed the distant metastasis. All cases were re-evaluated by a pathologist who confirmed the diagnosis and identified the tumor region for further molecular analysis. The study was conducted according to the national and institutional ethical policies, and it was previously approved by the Local Ethics Committee (CEP-548/2011).

DNA isolation from FFPE tissue

DNA was obtained from formalin-fixed paraffin-embedded tissue sections representative of the tumor lesions, as previously described with some modifications [31]. Briefly, serially 10- μ m-thick unstained sections of paraffin blocks were sectioned and one haematoxylin and eosin (H&E) section was used for identification and selection of the tumor area, which was macrodissected into a microfuge tube using a sterile needle (25 G – 0.5 mm, Neolus, Terumo Corporation, Tokyo, Japan). The macrodissected tissue was deparaffinized by a serial

wash with xylol and ethanol (100–70–50%) and allowed to air-dry. DNA was extracted using Qiagen's QIAamp DNA Micro Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions and quantified by NanoDrop 2000 (Thermo Scientific, Waltham, Massachusetts, USA). DNA samples were stored at –20°C until further genetic analysis.

Mutation analysis of *TERT*, *KIT*, *PDGFRA*, and *BRAF* and *NRAS* hotspot regions

The analysis of hotspot mutations of *TERT* promoter regions (contained the sites of c. –124:C>T and c. –146:C>T), *KIT* (exons 9, 11, 13, and 17), *PDGFRA* [12,14,18], *BRAF* (exon 15), and *NRAS* (codon12/13, 61), was performed by PCR followed by direct Sanger sequencing, as described previously [28,31–33]. Briefly, using specific pairs of primers (Supplementary Table 1), the target regions were amplified by PCR with an initial denaturation at 95°C for 15 min, followed by 40 cycles of 95°C denaturation for 45 s, specific annealing temperature for 90 s and 72°C elongation for 45 s, and 72°C final elongation for 7 min, in a Verity PCR machine (Applied Biosystems, Carlsbad, California, USA). Amplification of PCR products was confirmed by gel electrophoresis. Sequencing PCR was performed using a Big Dye terminator v3.1 cycle sequencing ready reaction kit (Applied Biosystems) and the ABI PRISM 3500 xL Genetic Analyzer (Applied Biosystems). All mutated cases were confirmed with a second PCR followed by sequencing.

Gene amplification analysis of *KIT*, *PDGFRA*, and *VEGFR2*

KIT, *PDGFRA*, and *VEGFR2* gene amplification was performed by quantitative real-time PCR with LightCycler (Roche Molecular Biochemicals, Mannheim, Germany), using fluorescent hybridization probes and fluorescence resonance energy transfer for the detection of PCR amplification product, as previously described [34,35]. In all assays, appropriate positive and negative controls of 4q12 loci amplification were included [36]. These controls had also been previously assessed by other methodologies, such as array-CGH and CISH [34,36,37].

Statistical analysis

The SPSS 19.0 software (IBM Corp., Armonk, New York, USA) was used for all statistics analyses. To identify associations among clinicopathological characteristics and the molecular alterations, and to compare the prevalence of molecular findings with the previous reports, the Fisher exact test was used. Associations of the same variables with specific cancer survival were analyzed by the Kaplan–Meier method to estimate specific cancer survival, with log-rank testing used to evaluate differences between curves. The *P*-value established for the statistics significance was up to 0.05.

Table 1 Demographic and clinicopathological features of 48 acral lentiginous melanomas

Characteristics	n (%)
Sex	
Male	23 (47.9)
Female	25 (52.1)
Age [mean (SD)]	62 (14.8)
Skin	
White	32 (66.7)
Pigmented	15 (31.3)
Not available	1 (2)
Anatomical location	
Hands	3 (6.3)
Feet	45 (93.7)
Breslow depth (mm)	
Up to 1	4 (8.3)
1.1–2	7 (14.6)
2.1–4	16 (33.3)
More than 4	17 (35.4)
Not available	4 (8.3)
Ulceration	
Yes	31 (64.6)
No	10 (20.8)
Not available	7 (14.6)
Mitotic rate (mm ²)	
0	0 (0)
1	11 (22.9)
> 1	37 (52.1)
Not available	12 (25)
TNM stage	
I	5 (10.4)
II	22 (45.8)
III	17 (35.4)
IV	4 (8.3)

Results

Clinicopathological features

The characteristics of the patients are shown in Table 1. The age ranged from 26 to 85 years old (mean 62 ± 14.8), and men and women had similar distribution. One-third of the cases presented non-white skin phenotype. Primary lesions were thick and ulcerated in most cases and arose only on the hands in three (6.3%) cases. Localized disease represented half of the cases. Breslow tumor thickness ranged from 0.62 to 20.0 mm, with a median of 4.9 (± 4.6). Five-year-specific cancer survival rate for all patients was 51.5%. Follow-up ranged from 2 to 174 months, with a mean of 35.5 (± 30.4) and a median of 29.6. At the last evaluation, 20 patients (41.7%) were alive without disease, nine were alive with disease (18.8%), 17 were dead because of melanoma (35.4%), and two were dead because of other causes (4.1%).

Table 3 Exons/codons location of the mutated genes

Mutation	n (%)
<i>TERT</i>	
c. – 146 C > T	4 (100)
<i>BRAF</i>	
Exon 15 (V600E)	5 (100)
<i>NRAS</i>	
Codon 61	3 (100)
<i>KIT</i>	
Exon 9	1 (16.6)
Exon 11	3 (50)
Exon 13	1 (16.6)
Exon 17	1 (16.6)
<i>PDGFRA</i>	
Exon 12	3 (60)
Exon 14	1 (20)
Exon 18	1 (20)

TERT, telomerase reverse transcriptase.

Molecular profile

Because of the low amount and/or poor DNA quality yield in some samples, we were not able to obtain molecular profiles from all ALM cases. Table 2 summarizes the molecular results. In primary tumors, we observed the presence of 7.0% (3/43) *TERT* mutations, harboring the c. – 146 C > T mutation in all cases (Tables 2 and 3). We found a frequency of mutations of 20.7% for *KIT*, 14.8% for *PDGFRA*, 7.5% for *NRAS*, and 10.3% for *BRAF* genes (Table 2). *KIT*, *PDGFRA*, and *VEGFR2* copy numbers were normal in all analyzed cases. One case exhibited two mutations in *PDGFRA*: one at exon 12 and another on exon 14 (Table 3). In the comparison of primary and LN or skin metastasis, we observed a similar molecular status, except in one case that showed *TERT* mutated only in the LN metastasis (Table 2). One case exhibited the same *BRAF* mutation in both primary and LN metastases (Table 2). The co-occurrence of mutations across specimens is demonstrated in Fig. 1. Only *BRAF* mutations were exclusive. *NRAS* was mutated mutually only in one case with *PDGFRA* mutation, and the same was observed with *TERT* and *KIT*; *KIT* and *PDGFRA* presented comutated with two other mutations (Fig. 1).

Clinicopathological and molecular associations


We found a significant association between *KIT* mutation status and higher Clark levels (Table 4). We also observed an association between *TERT* mutations and lower mitosis index (Table 4). There were no significant

Table 2 Molecular profile of acral lentiginous melanomas

Patient group	Mutations [n/N (%)]					Gene amplification [n/N (%)]		
	TERT	KIT	PDGFRA	NRAS	BRAF	KIT	PDGFRA	VEGFR2
Primary	3/43 (7.0)	6/29 (20.7)	4/27 (14.8)	3/40 (7.5)	4/39 (10.3)	0/35 (0)	0/36 (0)	0/8 (0)
Lymph node	1/10 (10)	0/10 (0)	0/8 (0)	0/10 (0)	1/11 (9.1)	0/10 (0)	0/10 (0)	0/2 (0)
Metastasis	0/2 (0)	0/2 (0)	0/2 (0)	0/2 (0)	0/2 (0)	0/2 (0)	0/2 (0)	0/0 (0)

TERT, telomerase reverse transcriptase; VEGFR2, vascular endothelial growth factor receptor-2.

Fig. 1



Distribution of *BRAF/NRAS/KIT/PDGFRA* and *TERT* promoter mutations of 48 acral lentiginous melanomas. *TERT*, telomerase reverse transcriptase.

Table 4 Association of molecular profile with clinicopathological features in acral lentiginous melanomas

Variables	n (%)									
	BRAF		NRAS		KIT		PDGFRA		TERT	
	WT	Mutated	WT	Mutated	WT	Mutated	WT	Mutated	WT	Mutated
Breslow (mm)										
Up to 1	2 (5.7)	0 (0)	3 (8.1)	0 (0)	1 (4.3)	0 (0)	1 (4.3)	0 (0)	2 (5.1)	1 (25)
1.1–2.0	5 (14.3)	1 (25)	6 (16.2)	0 (0)	4 (17.4)	1 (16.7)	4 (17.4)	1 (25)	7 (17.9)	0 (0)
2.1–4.0	12 (34.3)	2 (50)	12 (32.4)	3 (100)	9 (39.1)	3 (50)	10 (43.5)	2 (50)	13 (33.3)	1 (25)
More than 4	13 (37.1)	1 (25)	13 (35.1)	0 (0)	6 (36.1)	2 (33.3)	6 (26.1)	1 (25)	14 (35.9)	2 (50)
Unknown	3 (8.6)	0 (0)	3 (8.1)	0 (0)	3 (13)	0 (0)	2 (8.7)	0 (0)	3 (7.7)	0 (0)
		<i>P</i> =0.899		<i>P</i> =0.248		<i>P</i> =0.886		<i>P</i> =1.00		<i>P</i> =0.594
Clark										
II	1 (3.1)	0 (0)	1 (2.9)	0 (0)	–	–	–	–	1 (2.8)	0 (0)
III	2 (6.3)	0 (0)	3 (8.8)	0 (0)	3 (15)	0 (0)	2 (9.5)	1 (25)	4 (11.1)	1 (25)
IV	16 (50)	3 (75)	17 (50.0)	3 (100)	14 (70)	2 (33.3)	12 (57.1)	3 (75)	19 (52.8)	1 (25)
V	13 (40.6)	1 (25)	13 (38.2)	0 (0)	3 (15)	4 (66.7)	7 (33.3)	0 (0)	12 (33.3)	2 (50)
		<i>P</i> =0.736		<i>P</i> =0.428		<i>P</i> =0.043		<i>P</i> =0.400		<i>P</i> =0.491
Ulceration										
Absent	6 (19.4)	0 (0)	6 (18.8)	2 (66.6)	4 (19)	2 (33.3)	5 (22.7)	1 (25)	8 (22.9)	1 (33.3)
Present	25 (80.6)	4 (100)	26 (81.2)	1 (33.3)	17 (81)	4 (66.7)	17 (77.3)	3 (75)	27 (77.1)	2 (66.7)
		<i>P</i> =0.581		<i>P</i> =1.00		<i>P</i> =0.588		<i>P</i> =1.00		<i>P</i> =0.164
Mitosis										
1	6 (22.2)	2 (66.7)	9 (32.1)	0 (0)	5 (27.8)	2 (33.3)	7 (35)	0 (0)	6 (20)	3 (100)
> 1	21 (77.8)	1 (33.3)	19 (67.9)	3 (100)	13 (72.2)	4 (66.7)	13 (65)	3 (100)	24 (80)	0 (0)
		<i>P</i> =0.166		<i>P</i> =0.537		<i>P</i> =1.00		<i>P</i> =0.526		<i>P</i> =0.015
TNM stage										
I	3 (8.6)	0 (0)	4 (10.8)	0 (0)	2 (8.7)	0 (0)	1 (4.3)	1 (25)	4 (10.3)	1 (25)
II	17 (48.6)	2 (50)	18 (48.6)	2 (66.7)	10 (43.5)	4 (66.7)	13 (56.5)	1 (25)	18 (46.2)	1 (25)
III	11 (31.4)	2 (50)	12 (32.4)	1 (33.3)	8 (34.8)	2 (33.3)	7 (30.4)	2 (50)	13 (33.3)	2 (50)
IV	4 (11.4)	0 (0)	3 (8.1)	0 (0)	3 (13)	0 (0)	2 (8.7)	0 (0)	4 (10.3)	0 (0)
		<i>P</i> =1.00		<i>P</i> =1.00		<i>P</i> =0.776		<i>P</i> =0.440		<i>P</i> =0.751

Fisher's exact test was applied in all cases. Bold number indicates significant association. *TERT*, telomerase reverse transcriptase.

differences in the survival rates according to *TERT*, *BRAF*, *NRAS*, *KIT*, and *PDGFRA* mutation status in the univariate analysis (data not shown).

Discussion

In the present study, we described for the first time the molecular profile of Brazilian acrolentiginous melanomas. We showed that Brazilian patients exhibited a similar profile to the one described in the international literature [12,18–20,24,28–30,38–64] (Table 5).

We found that *TERT* c. –146C>T promoter mutation is present in 9.3% of patients, and this frequency was in concordance with the reported 7.6% (7/92) (*P*=0.732) (Table 5). However, our study represents the larger analysis of the gene hotspot region of *TERT* in this melanoma subtype (Table 5). Previous studies in thyroid cancer and skin melanomas suggested an association of *TERT* mutation with other molecular alterations, such as *BRAF* [28]. In superficial melanomas, thyroid cancer, and gliomas, *TERT* promoter mutations have been also reported to be

Table 5 Summary of molecular features of acral lentiginous melanomas

References	Country	Mutations [n/N (%)]					Gene amplification [n/N (%)]		
		TERT	KIT	PDGFRA	NRAS	BRAF	KIT	PDGFRA	VEGFR2
This study	Brazil	4/43 (9.3)	6/29 (20.7)	4/27 (14.8)	3/40 (7.5)	4/39 (10.3)	0/35 (0)	0/36 (0)	0/8 (0)
Heidenreich et al. [29]	Spain	4/42 (9.5)	NA	NA	2/38 (5.3)	12/30 (28.6)	NA	NA	NA
Pozzobon et al. [38]	Spain	NA	NA	NA	0/10 (0)	2/10 (20)	NA	NA	NA
Liau et al. [30]	China	2/32 (6.25)	NA	NA	NA	NA	NA	NA	NA
Dai et al. [39]	China	NA	NA	9/132 (9.8)	NA	NA	NA	NA	NA
Vinagre et al. [28]	Portugal	1/14 (7.1)	NA	NA	NA	2/14 (14.3)	NA	NA	NA
Hodi et al. [20]	Multicenter	NA	2/6 (33.3)	NA	NA	NA	4/4 (100)	NA	NA
Zebary et al. [41]	Sweden	NA	13/88 (14.8)	NA	13/88 (14.8)	15/88 (17.0)	NA	NA	NA
Greaves et al. [48]	USA	NA	NA	NA	NA	18/111 (16.2)	NA	NA	NA
Dai et al. [39]	China	NA	9/39 (23.1)	NA	NA	NA	NA	NA	NA
Lin et al. [40]	China	NA	0/20 (0)	NA	NA	NA	NA	NA	NA
Puig-Butillé et al. [42]	Spain	NA	NA	NA	3/17 (17.6)	0/17 (0)	NA	NA	NA
Minor et al. [24]	USA	NA	3/22 (13.6)	NA	6/22 (27.3)	7/22 (31.8)	6/22 (27.3)	NA	NA
Si et al. [47]	China	NA	NA	NA	13/148 (8.8)	23/148 (15.5)	NA	NA	NA
Ashida et al. [43]	Japan	NA	4/44 (9.1)	NA	NA	NA	NA	NA	NA
Jakob et al. [44]	USA	NA	NA	NA	7/44 (15.9)	6/44 (13.6)	NA	NA	NA
Yun et al. [45]	Korea	NA	4/40 (10)	NA	NA	NA	7/20 (35.0)	NA	NA
Kong et al. [46]	China	NA	23/193 (11.9)	NA	NA	NA	14/193 (7.3)	NA	NA
Handolias and colleagues [49,50]	Australia	NA	2/6 (33.3)	NA	NA	NA	NA	NA	NA
Terada [51]	Japan	NA	1/2 (50)	0/2 (0)	NA	NA	NA	NA	NA
Handolias and colleagues [49,50]	Australia	NA	1/16 (6.25)	NA	NA	NA	NA	NA	NA
Torres-Cabala et al. [52]	USA	NA	5/39 (12.8)	NA	NA	NA	NA	NA	NA
Ashida et al. [53]	Japan	NA	2/16 (12.5)	NA	0/22 (0)	3/22 (13.6)	3/16 (18.8)	NA	NA
Curtin et al. [54]	USA	NA	NA	0/7 (0)	NA	NA	NA	3/7 (42.9)	NA
Akslen et al. [55]	Africa	NA	NA	NA	3/26 (11.5)	1/25 (4.0)	NA	NA	NA
Beadling et al. [56]	USA	NA	3/13 (23.1)	NA	1/9 (11.1)	2/12 (16.7)	3/11 (27.3)	NA	NA
Takata et al. [57]	Japan	NA	NA	NA	1/6 (16.7)	2/8 (24.05)	NA	NA	NA
Liu et al. [59]	Australia	NA	NA	NA	NA	0/6 (0)	NA	NA	NA
Curtin et al. [19]	USA	NA	3/28 (10.7)	NA	0/10 (0)	1/10 (10.0)	2/28 (7.1)	NA	NA
Saldanha et al. [18]	UK	NA	NA	NA	9/19 (47.4)	2/21 (9.5)	NA	NA	NA
Davison et al. [60]	USA	NA	NA	NA	NA	0/3 (0)	NA	NA	NA
Curtin et al. [12]	Multicenter	NA	NA	NA	3/35 (8.6)	7/35 (20.0)	NA	NA	NA
Takata et al. [58]	Japan	NA	NA	NA	1/28 (3.6)	3/28 (10.8)	NA	NA	NA
Lang and MacKie [61]	UK	NA	NA	NA	NA	2/17 (11.8)	NA	NA	NA
Sasaki et al. [62]	Germany	NA	NA	NA	0/15 (0)	5/15 (33.3)	NA	NA	NA
Reifenberger et al. [63]	Germany	NA	NA	NA	1/4 (25.0)	2/4 (50.0)	NA	NA	NA
Maldonado et al. [64]	USA/Japan	NA	NA	NA	NA	4/30 (13.3)	NA	NA	NA
All the referred studies	–	7/92 (7.6)	75/572 (13.1)	9/145 (6.2)	63/544 (11.6)	119/724 (16.4)	39/294 (13.3)	3/7 (42.9)	NA
Total	–	11/131 (8.4)	81/601 (13.5)	13/168 (7.7)	66/581 (11.4)	123/759 (16.2)	39/329 (11.9)	3/43 (7.0)	0/8 (0)

NA, not available; TERT, telomerase reverse transcriptase; VEGFR2, vascular endothelial growth factor receptor-2.

associated with older patients and worse patient prognosis [65]. In our study, we did not observe such molecular and clinical associations in acrolentiginous melanomas. The association with low mitotic index could be related to better prognosis, but the small number of mutated cases needs further confirmation in a larger series.

Acrolentiginous melanomas are known to harbor a distinct molecular profile of superficial melanomas. Compelling evidence showed the paramount role of *KIT* and *PDGFRA* alterations in the melanoma subtype [66]. In the present study, we found a mutation frequency of 16.7 and 12.5% for *KIT* and *PDGFRA*, respectively. Among *KIT* mutations, 50% were located on exon 11 and the others were distributed among the other exons [9,13,17], whereas 60% of *PDGFRA* mutations were located on exon 12 and the others were distributed on exons 14 and 18. These mutation rates are in accordance with previous descriptions of 13.1% (75/572) and 6.2% (9/145) ($P=0.145$ and 0.108) (Table 5). Some authors have

identified *KIT* gene amplification as an alternative mechanism for *KIT* upregulation in these tumors [18–20]. However, we did not find any case with *KIT* gene amplification using quantitative real-time PCR, a very sensitive and specific methodology, previously optimized by us and reported by our group and other groups [34,36,67,68]. We observed an association between *KIT* mutation and higher Clark staging, indicating that *KIT* could be associated with a more aggressive disease. The presence of *KIT* and *PDGFRA* alterations, namely mutations, has an important therapeutic impact. It has been shown that some melanoma patients exhibiting *KIT/PDGFRA* mutations, rather than gene amplification, can benefit from imatinib-based targeted therapy [69]. Recently, *KIT/PDGFRA* mutations have been found as targets for other small inhibitors, such as crenolanib and nilotinib [39,70]. Therefore, we can anticipate that between 15 and 20% of Brazilian acrolentiginous melanoma patients would benefit from these above-mentioned therapeutic modalities.

We also analyzed in this study the frequency of *NRAS* (codon12/13 and 61) and *BRAF* (exon 15) hotspot mutations. We found that *NRAS* was mutated in 6.3% of cases. This frequency was somehow below what is reported worldwide, which is above 11% (63/544) ($P=0.787$) of cases (Table 5). However, *NRAS* frequency reports range from 0 to 47% (Table 5); therefore, there is not really a consensus in the literature, and this could be because of the distinct sizes of series analyzed in each study, or it could reflect the differences between populations.

With regard to *BRAF*, we observed the presence of 10.6% (all V600E) of mutated cases. This result is in line with the reported (16.4% – 119/724) ($P=0.639$) (Table 5), where the values range from 0 to 33% of *BRAF* mutated cases. *BRAF* mutations were exclusive events. Besides the low frequency of all mutations, they were mutually present at least in one-third of the cases, except for the *BRAF*, corroborating the high spectrum of melanoma mutations reported.

In conclusion, we have molecularly characterized a series of 48 ALMs. We have shown that less than 10% of patients harbor the recurrent *TERT* promoter mutation, and *KIT* represents the most mutated gene (21%), followed by *PDGFRA* (15%), in this series. These results could help in the selection of other alternative and potential therapeutic options for Brazilian acral lentiginous melanoma patients.

Acknowledgements

This project was supported by FAPESP – Brazil (2012/4194-1) to Vinicius de Lima Vazquez.

Conflicts of interest

There are no conflicts of interest.

References

- Erdmann F, Lortet-Tieulent J, Schuz J, Zeeb H, Greinert R, Breitbart EW, et al. International trends in the incidence of malignant melanoma 1953-2008 – are recent generations at higher or lower risk? *Int J Cancer* 2013; **132**:385–400.
- Katalinic A, Kunze U, Schäfer T. Epidemiology of cutaneous melanoma and non-melanoma skin cancer in Schleswig-Holstein, Germany: incidence, clinical subtypes, tumour stages and localization (epidemiology of skin cancer). *Br J Dermatol* 2003; **149**:1200–1206.
- Bradford PT, Goldstein AM, McMaster ML, Tucker MA. Acral lentiginous melanoma: incidence and survival patterns in the United States, 1986–2005. *Arch Dermatol* 2009; **145**:427–434.
- Clark WH Jr, From L, Bernardino EA, Mihm MC. The histogenesis and biologic behavior of primary human malignant melanomas of the skin. *Cancer Res* 1969; **29**:705–727.
- McGovern VJ, Mihm MC Jr, Bailly C, Booth JC, Clark WH Jr, Cochran AJ, et al. The classification of malignant melanoma and its histologic reporting. *Cancer* 1973; **32**:1446–1457.
- Hudson DA, Krige JE. Melanoma in black South Africans. *J Am Coll Surg* 1995; **180**:65–71.
- Kabigting FD, Nelson FP, Kauffman CL, Popoveniuc G, Dasanu CA, Alexandrescu DT. Malignant melanoma in African-Americans. *Dermatol Online J* 2009; **15**:3.
- Chang JW. Acral melanoma: a unique disease in Asia. *JAMA Dermatol* 2013; **149**:1272–1273.
- Brandão FV, Pereira AF, Gontijo B, Bittencourt FV. Epidemiological aspects of melanoma at a university hospital dermatology center over a period of 20 years. *An Bras Dermatol* 2013; **88**:344–353.
- Vilanova CM, Lages RB, Ribeiro SM, Almeida IP, Santos LG, Vieira SC. Epidemiological and histopathological profile of cutaneous melanoma at a center in northeastern Brazil from 2000 to 2010. *An Bras Dermatol* 2013; **88**:545–553.
- Vazquez Vde L, Silva TB, Vieira Mde A, de Oliveira AT, Lisboa MV, de Andrade DA, et al. Melanoma characteristics in Brazil: demographics, treatment, and survival analysis. *BMC Res Notes* 2015; **8**: ():4.
- Curtin JA, Fridlyand J, Kageshita T, Patel HN, Busam KJ, Kutzner H, et al. Distinct sets of genetic alterations in melanoma. *N Engl J Med* 2005; **353**:2135–2147.
- Bello DM, Ariyan CE, Carvajal RD. Melanoma mutagenesis and aberrant cell signaling. *Cancer Control* 2013; **20**:261–281.
- Gray-Schopfer V, Wellbrock C, Marais R. Melanoma biology and new targeted therapy. *Nature* 2007; **445**:851–857.
- Davies H, Bignell GR, Cox C, Stephens P, Edkins S, Clegg S, et al. Mutations of the *BRAF* gene in human cancer. *Nature* 2002; **417**:949–954.
- Pollock PM, Meltzer PS. A genome-based strategy uncovers frequent *BRAF* mutations in melanoma. *Cancer Cell* 2002; **2**:5–7.
- Davies MA, Gershenwald JE. Targeted therapy for melanoma: a primer. *Surg Oncol Clin N Am* 2011; **20**:165–180.
- Saldanha G, Potter L, Daformo P, Pringle JH. Cutaneous melanoma subtypes show different *BRAF* and *NRAS* mutation frequencies. *Clin Cancer Res* 2006; **12**:4499–4505.
- Curtin JA, Busam K, Pinkel D, Bastian BC. Somatic activation of *KIT* in distinct subtypes of melanoma. *J Clin Oncol* 2006; **24**:4340–4346.
- Hodi FS, Corless CL, Giobbie-Hurder A, Fletcher JA, Zhu M, Marino-Enriquez A, et al. Imatinib for melanomas harboring mutationally activated or amplified *KIT* arising on mucosal, acral, and chronically sun-damaged skin. *J Clin Oncol* 2013; **31**:3182–3190.
- Yin LY, Wu Y, Ballinger CA, Patterson C. Genomic structure of the human *KDR/flk-1* gene. *Mamm Genome* 1998; **9**:408–410.
- Holtkamp N, Ziegenhagen N, Malzer E, Hartmann C, Giese A, von Deimling A. Characterization of the amplicon on chromosomal segment 4q12 in glioblastoma multiforme. *Neuro Oncol* 2007; **9**:291–297.
- Flaherty KT, Infante JR, Daud A, Gonzalez R, Keefe RF, Sosman J, et al. Combined *BRAF* and *MEK* inhibition in melanoma with *BRAF* V600 mutations. *N Engl J Med* 2012; **367**:1694–1703.
- Minor DR, Kashani-Sabet M, Garrido M, O'Day SJ, Hamid O, Bastian BC. Sunitinib therapy for melanoma patients with *KIT* mutations. *Clin Cancer Res* 2012; **18**:1457–1463.
- Johansson I, Aaltonen KE, Ebbesson A, Grabau D, Wigerup C, Hedenfalk I, Rydén L. Increased gene copy number of *KIT* and *VEGFR2* at 4q12 in primary breast cancer is related to an aggressive phenotype and impaired prognosis. *Genes Chromosomes Cancer* 2012; **51**:375–383.
- Horn S, Figl A, Rachakonda PS, Fischer C, Sucker A, Gast A, et al. *TERT* promoter mutations in familial and sporadic melanoma. *Science* 2013; **339**:959–961.
- Huang FW, Hodis E, Xu MJ, Kryukov GV, Chin L, Garraway LA. Highly recurrent *TERT* promoter mutations in human melanoma. *Science* 2013; **339**:957–959.
- Vinagre J, Almeida A, Pópulo H, Batista R, Lyra J, Pinto V, et al. Frequency of *TERT* promoter mutations in human cancers. *Nat Commun* 2013; **4**:2185.
- Heidenreich B, Rachakonda PS, Hemminki K, Kumar R. *TERT* promoter mutations in cancer development. *Curr Opin Genet Dev* 2014; **24**:30–37.
- Liau JY, Tsai JH, Jeng YM, Chu CY, Kuo KT, Liang CW. *TERT* promoter mutation is uncommon in acral lentiginous melanoma. *J Cutan Pathol* 2014; **41**:504–508.
- Yamane LS, Scapulatempo-Neto C, Alvarenga L, Oliveira CZ, Berardinelli GN, Almodova E, et al. *KRAS* and *BRAF* mutations and MSI status in precursor lesions of colorectal cancer detected by colonoscopy. *Oncol Rep* 2014; **32**:1419–1426.
- Martinho O, Gouveia A, Viana-Pereira M, Silva P, Pimenta A, Reis RM, Lopes JM. Low frequency of *MAP* kinase pathway alterations in *KIT* and *PDGFRA* wild-type GISTs. *Histopathology* 2009; **55**:53–62.
- Campanella NC, Celestino R, Pestana A, Scapulatempo-Neto C, de Oliveira AT, Brito MJ, et al. Low frequency of *TERT* promoter mutations in gastrointestinal stromal tumors (GISTs). *Eur J Hum Genet* 2015; **23**:877–879.
- Gomes AL, Reis-Filho JS, Lopes JM, Martinho O, Lambros MB, Martins A, et al. Molecular alterations of *KIT* oncogene in gliomas. *Cell Oncol* 2007; **29**:399–408.
- Martinho O, Longatto-Filho A, Lambros MB, Martins A, Pinheiro C, Silva A, et al. Expression, mutation and copy number analysis of platelet-derived

- growth factor receptor A (PDGFRA) and its ligand PDGFA in gliomas. *Br J Cancer* 2009; **101**:973–982.
- 36 Martinho O, Silva-Oliveira R, Miranda-Gonçalves V, Clara C, Almeida JR, Carvalho AL, *et al.* In vitro and in vivo analysis of RTK inhibitor efficacy and identification of its novel targets in glioblastomas. *Transl Oncol* 2013; **6**:187–196.
- 37 Martinho O, Longatto-Filho A, Lambros MB, Martins A, Pinheiro C, Silva A, *et al.* Expression, mutation and copy number analysis of platelet-derived growth factor receptor A (PDGFRA) and its ligand PDGFA in gliomas. *Br J Cancer* 2009; **101**:973–982.
- 38 Pozzobon FC, Puig-Butillé JA, González-Alvarez T, Carrera C, Aguilera P, Alos L, *et al.* Dermoscopic criteria associated with BRAF and NRAS mutation status in primary cutaneous melanoma. *Br J Dermatol* 2014; **171**:754–759.
- 39 Dai J, Kong Y, Si L, Chi Z, Cui C, Sheng X, *et al.* Large-scale analysis of PDGFRA mutations in melanomas and evaluation of their sensitivity to tyrosine kinase inhibitors imatinib and crenolanib. *Clin Cancer Res* 2013; **19**:6935–6942.
- 40 Lin YC, Chang YM, Ho JY, Lin HC, Tsai YM, Chiang CP, *et al.* C-kit expression of melanocytic neoplasm and association with clinicopathological parameters and anatomic locations in Chinese people. *Am J Dermatopathol* 2013; **35**:569–575.
- 41 Zebary A, Omholt K, Vassilaki I, Höimö V, Lindén D, Viberg L, *et al.* KIT, NRAS, BRAF and PTEN mutations in a sample of Swedish patients with acral lentiginous melanoma. *J Dermatol Sci* 2013; **72**:284–289.
- 42 Puig-Butillé JA, Badenas C, Ogbah Z, Carrera C, Aguilera P, Malveyh J, Puig S. Genetic alterations in RAS-regulated pathway in acral lentiginous melanoma. *Exp Dermatol* 2013; **22**:148–150.
- 43 Ashida A, Uhara H, Kiniwa Y, Oguchi M, Murata H, Goto Y, *et al.* Assessment of BRAF and KIT mutations in Japanese melanoma patients. *J Dermatol Sci* 2012; **66**:240–242.
- 44 Jakob JA, Bassett RL Jr, Ng CS, Curry JL, Joseph RW, Alvarado GC, *et al.* NRAS mutation status is an independent prognostic factor in metastatic melanoma. *Cancer* 2012; **118**:4014–4023.
- 45 Yun J, Lee J, Jang J, Lee EJ, Jang KT, Kim JH, Kim KM. KIT amplification and gene mutations in acral/mucosal melanoma in Korea. *APMIS* 2011; **119**:330–335.
- 46 Kong Y, Si L, Zhu Y, Xu X, Corless CL, Flaherty KT, *et al.* Large-scale analysis of KIT aberrations in Chinese patients with melanoma. *Clin Cancer Res* 2011; **17**:1684–1691.
- 47 Si L, Kong Y, Xu X, Flaherty KT, Sheng X, Cui C, *et al.* Prevalence of BRAF V600E mutation in Chinese melanoma patients: large scale analysis of BRAF and NRAS mutations in a 432-case cohort. *Eur J Cancer* 2012; **48**:94–100.
- 48 Greaves WO, Verma S, Patel KP, Davies MA, Barkoh BA, Galbincea JM, *et al.* Frequency and spectrum of BRAF mutations in a retrospective, single-institution study of 1112 cases of melanoma. *J Mol Diagn* 2013; **15**:220–226.
- 49 Handolias D, Hamilton AL, Salemi R, Tan A, Moodie K, Kerr L, *et al.* Clinical responses observed with imatinib or sorafenib in melanoma patients expressing mutations in KIT. *Br J Cancer* 2010; **102**:1219–1223.
- 50 Handolias D, Salemi R, Murray W, Tan A, Liu W, Viros A, *et al.* Mutations in KIT occur at low frequency in melanomas arising from anatomical sites associated with chronic and intermittent sun exposure. *Pigment Cell Melanoma Res* 2010; **23**:210–215.
- 51 Terada T. Low incidence of KIT gene mutations and no PDGFRA gene mutations in primary cutaneous melanoma: an immunohistochemical and molecular genetic study of Japanese cases. *Int J Clin Oncol* 2010; **15**:453–456.
- 52 Torres-Cabala CA, Wang WL, Trent J, Yang D, Chen S, Galbincea J, *et al.* Correlation between KIT expression and KIT mutation in melanoma: a study of 173 cases with emphasis on the acral-lentiginous/mucosal type. *Mod Pathol* 2009; **22**:1446–1456.
- 53 Ashida A, Takata M, Murata H, Kido K, Saida T. Pathological activation of KIT in metastatic tumors of acral and mucosal melanomas. *Int J Cancer* 2009; **124**:862–868.
- 54 Curtin JA, Pincel D, Bastian BC. Absence of PDGFRA mutations in primary melanoma. *J Invest Dermatol* 2008; **128**:488–489.
- 55 Akslen LA, Puntervoll H, Bachmann IM, Straume O, Vuhahula E, Kumar R, Molven A. Mutation analysis of the EGFR-NRAS-BRAF pathway in melanomas from black Africans and other subgroups of cutaneous melanoma. *Melanoma Res* 2008; **18**:29–35.
- 56 Beadling C, Jacobson-Dunlop E, Hodi FS, Le C, Warrick A, Patterson J, *et al.* KIT gene mutations and copy number in melanoma subtypes. *Clin Cancer Res* 2008; **14**:6821–6828.
- 57 Takata M, Lin J, Takayanagi S, Suzuki T, Ansai S, Kimura T, *et al.* Genetic and epigenetic alterations in the differential diagnosis of malignant melanoma and spitzoid lesion. *Br J Dermatol* 2007; **156**:1287–1294.
- 58 Takata M, Goto Y, Ichii N, Yamaura M, Murata H, Koga H, *et al.* Constitutive activation of the mitogen-activated protein kinase signaling pathway in acral melanomas. *J Invest Dermatol* 2005; **125**:318–322.
- 59 Liu W, Kelly JW, Trivett M, Murray WK, Dowling JP, Wolfe R, *et al.* Distinct clinical and pathological features are associated with the BRAF(T1799A (V600E)) mutation in primary melanoma. *J Invest Dermatol* 2007; **127**:900–905.
- 60 Davison JM, Rosenbaum E, Barrett TL, Goldenberg D, Hoque MO, Sidransky D, Westra WH. Absence of V599E BRAF mutations in desmoplastic melanomas. *Cancer* 2005; **103**:788–792.
- 61 Lang J, MacKie RM. Prevalence of exon 15 BRAF mutations in primary melanoma of the superficial spreading, nodular, acral, and lentigo maligna subtypes. *J Invest Dermatol* 2005; **125**:575–579.
- 62 Sasaki Y, Niu C, Makino R, Kudo C, Sun C, Watanabe H, *et al.* BRAF point mutations in primary melanoma show different prevalences by subtype. *J Invest Dermatol* 2004; **123**:177–183.
- 63 Reizenberger J, Knobbe CB, Sterzinger AA, Blaschke B, Schulte KW, Ruzicka T, Reizenberger G. Frequent alterations of Ras signaling pathway genes in sporadic malignant melanomas. *Int J Cancer* 2004; **109**:377–384.
- 64 Maldonado JL, Fridlyand J, Patel H, Jain AN, Busam K, Kageshita T, *et al.* Determinants of BRAF mutations in primary melanomas. *J Natl Cancer Inst* 2003; **95**:1878–1890.
- 65 Griewank KG, Murali R, Puig-Butillé JA, Schilling B, Livingstone E, Potrony M, *et al.* TERT promoter mutation status as an independent prognostic factor in cutaneous melanoma. *J Natl Cancer Inst* 2014; **106**:dju246.
- 66 Woodman SE, Davies MA. Targeting KIT in melanoma: a paradigm of molecular medicine and targeted therapeutics. *Biochem Pharmacol* 2010; **80**:568–574.
- 67 Baruchel S, Sharp JR, Bartels U, Hukin J, Odame I, Portwine C, *et al.* A Canadian paediatric brain tumour consortium (CPBTC) phase II molecularly targeted study of imatinib in recurrent and refractory paediatric central nervous system tumours. *Eur J Cancer* 2009; **45**:2352–2359.
- 68 Ozawa T, Brennan CW, Wang L, Squatrito M, Sasayama T, Nakada M, *et al.* PDGFRA gene rearrangements are frequent genetic events in PDGFRA-amplified glioblastomas. *Genes Dev* 2010; **24**:2205–2218.
- 69 Bello DM, Dematteo RP, Ariyan CE. The GIST of targeted therapy for malignant melanoma. *Ann Surg Oncol* 2014; **21**:2059–2067.
- 70 Carvajal RD, Lawrence DP, Weber JS, Gajewski TF, Gonzalez R, Lutzky J, *et al.* Phase II study of nilotinib in melanoma harboring KIT alterations following progression to prior KIT inhibition. *Clin Cancer Res* 2015; **21**:2289–2296.