



Molecular and physiological basis of *Saccharomyces cerevisiae* tolerance to adverse lignocellulose-based process conditions

Joana T. Cunha¹ · Aloia Romani¹ · Carlos E. Costa¹ · Isabel Sá-Correia² · Lucília Domingues¹ 

Received: 20 July 2018 / Revised: 19 October 2018 / Accepted: 22 October 2018 / Published online: 5 November 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Lignocellulose-based biorefineries have been gaining increasing attention to substitute current petroleum-based refineries. Biomass processing requires a pretreatment step to break lignocellulosic biomass recalcitrant structure, which results in the release of a broad range of microbial inhibitors, mainly weak acids, furans, and phenolic compounds. *Saccharomyces cerevisiae* is the most commonly used organism for ethanol production; however, it can be severely distressed by these lignocellulose-derived inhibitors, in addition to other challenging conditions, such as pentose sugar utilization and the high temperatures required for an efficient simultaneous saccharification and fermentation step. Therefore, a better understanding of the yeast response and adaptation towards the presence of these multiple stresses is of crucial importance to design strategies to improve yeast robustness and bioconversion capacity from lignocellulosic biomass. This review includes an overview of the main inhibitors derived from diverse raw material resultants from different biomass pretreatments, and describes the main mechanisms of yeast response to their presence, as well as to the presence of stresses imposed by xylose utilization and high-temperature conditions, with a special emphasis on the synergistic effect of multiple inhibitors/stressors. Furthermore, successful cases of tolerance improvement of *S. cerevisiae* are highlighted, in particular those associated with other process-related physiologically relevant conditions. Decoding the overall yeast response mechanisms will pave the way for the integrated development of sustainable yeast cell-based biorefineries.

Keywords Lignocellulosic biomass · Inhibitory compounds · Stress response mechanisms · *S. cerevisiae* · Metabolic engineering

Introduction

Depletion of fossil resources and environmental concerns related to their exploitation promote the transition of petroleum-based refinery towards a bio-based economy. The bioeconomy aims at the manufacturing of products and fuels from renewable materials using efficient biotechnologies, contributing to the creation of new jobs and industries.

Lignocellulosic biomass is the most available renewable resource on earth and may be used for the production of liquid biofuels, such as bioethanol. Nevertheless, the large-scale

production of lignocellulosic bioethanol is not extensively implemented due to the elevated initial investment and operational costs related to the process. For instance, a physicochemical pretreatment is required to break down the recalcitrant and complex structure of lignocellulosic biomass to obtain fermentable sugars, significantly increasing the complexity and length of the process. Several strategies have been considered to reduce capital costs such as the use of whole slurry (liquid and solid phases) or slurries after pretreatment, to eliminate unnecessary washing steps, and operating at high solid loading, to reduce distillation costs (Romaní et al. 2014). Moreover, the use of all sugars present in the hemicellulosic fraction is also required for cost-effective lignocellulosic ethanol production. Xylose is the most abundant sugar in the hemicellulosic fraction; however, *Saccharomyces cerevisiae*, the preferred microorganism for bioethanol production, is not naturally capable of metabolizing this sugar. Considering this, several efforts have been applied in the last years for the development of *S. cerevisiae* strains capable of xylose consumption through the expression of heterologous pathways, such as

✉ Lucília Domingues
luciliad@deb.uminho.pt

¹ Centre of Biological Engineering (CEB), University of Minho, 4710-057 Braga, Portugal

² Institute for Bioengineering and Biosciences, Department of Bioengineering, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisbon, Portugal

xylose reductase and xylitol dehydrogenase (XR/XDH) from *Pichia stipitis*, or of xylose isomerases (XIs) from different bacterial and fungal species (Moysés et al. 2016). Both pathways convert xylose into xylulose, which is subsequently phosphorylated into xylulose-5P and then further metabolized in the pentose phosphate pathway (PPP).

Nevertheless, the hemicellulosic fraction (liquid phase) also contains toxic compounds, in concentrations dependent of pretreatment severity, which are inhibitors of the subsequent saccharification and fermentation steps (Modenbach and Nokes 2012). These inhibitory compounds are weak acids, furans, and phenolic compounds. Acetic acid is the most abundant weak acid in lignocellulosic hydrolysates and is present due to the deacetylation of acetyl groups linked to the main chain of hemicelluloses. Other weak acids, such as formic and levulinic acids, can also be present in hydrolysates resulting from furan compound degradation. Furfural and hydroxymethylfurfural (HMF) are produced by dehydration of pentoses and hexoses, respectively. On the other hand, phenolic compounds (such as syringic acid, vanillin, ferulic acid, vanillic acid, and coumaric acid) are produced by depolymerization of lignin. The amount of inhibitory compounds in the lignocellulosic hydrolysates is dependent on lignocellulosic source (e.g., agricultural residues, hardwoods, or softwoods), the selected pretreatment (hydrothermal treatment, diluted acid treatment, alkali treatment), and operational conditions (solid loading of lignocellulosic biomass, temperature, time, percentage of catalyst) (Modenbach and Nokes 2012; Ko et al. 2015; Dominguez et al. 2017). Additionally, the concentration of hemicellulose-derived sugars (xylose and xylooligosaccharides) can also vary depending on raw material and pretreatment selected for the processing of lignocellulosic biomass (Table 1). As seen in Table 1, the increase of temperature (from 210 to 220 °C) in steam explosion treatment increased the acetic acid concentration, furfural, and HMF in wheat straw hydrolysate (Alvira et al. 2011). Moreover, the acid-diluted treatment yields a higher concentration of hemicellulose-derived compounds as monomers (xylose, furfural, HMF, or acetic acid) than autohydrolysis treatment (using only water as reaction medium), since part of hemicellulose-derived compounds is solubilized as oligomers (xylooligosaccharides or acetyl groups) in autohydrolysis treatments (Yáñez et al. 2009; Jesus et al. 2017). On the other hand, the total inhibitory load in hardwood (e.g., eucalyptus) hydrolysates is superior to the inhibitory load in hydrolysates from agricultural residues (corn cob and wheat straw) (Costa et al. 2017). HMF concentration is higher than furfural in softwood hydrolysates as hemicellulose is composed mainly by hexoses (Table 1), while furfural is the predominant furan compound in hardwood and agricultural residue hydrolysates (Westman et al. 2012; Dominguez et al. 2017). Phenolic compounds are generally present in hydrolysates at lower concentration (Table 1), and their inhibitory effect is more described for enzymes in cellulose conversion (Ko

et al. 2015). Under acid conditions, the formation of phenolic compounds can differ, depending of lignocellulosic source and treatment conditions (Ko et al. 2015). These compounds are generally considered inhibitory for *S. cerevisiae* growth, affecting its fermentative performance by increasing the fermentation lag phase and decreasing ethanol yield and productivity (Guo and Olsson 2014; Larsson et al. 2000; Liu et al. 2004). Furthermore, besides the presence of more than one inhibitory compound in the hydrolysate, for an efficient conversion of cellulose to glucose, higher fermentation temperatures are desirable in order to facilitate simultaneous saccharification and fermentation (SSF) processes, which can increase the yield of lignocellulosic ethanol (Kelbert et al. 2016), representing an additional stress factor. To partially overcome these physiological hurdles, proper nutrient supplementation together with an adequate yeast genetic background has shown to increase process efficiency (Kelbert et al. 2015).

The adaptive response mechanism of yeast cells towards the presence of a single inhibitor such as acetic acid (Dong et al. 2017; Giannattasio et al. 2013; Guerreiro et al. 2016; Lindberg et al. 2013; Mira et al. 2010), formic acid (Henriques et al. 2017), furfural (Allen et al. 2010; Gorsich et al. 2006), HMF (Ma and Liu 2010), and vanillin (Nguyen et al. 2014a, b; Wang et al. 2016) has been extensively described and studied. Nevertheless, yeast response towards the synergistic effect of multiple inhibitors is generally less approached (Pereira et al. 2011a, 2014b). Furthermore, aforementioned conditions required for cost-efficient production of ethanol (high temperature and xylose co-consumption) together with the presence of inhibitory compounds will further increase the negative effects on lignocellulosic fermentation performance. Recently, the focus on more robust or tolerant yeast is emerging as a desirable strategy for the fermentation of lignocellulosic hydrolysates, with the previous phenotypic selection of stress-tolerant strains being an essential step (Jin et al. 2013; Wimalasena et al. 2014; Romani et al. 2015). In this sense, this review aims to describe the negative effects caused by the presence of multiple lignocellulose-derived inhibitors linked to required process conditions (high temperature and xylose co-consumption), as well as the mechanisms of yeast response and tolerance towards the simultaneous presence of all these fermentation constrains (Fig. 1). Different from other reviews covering yeast lignocellulosic tolerance, this work will focus on the overall effect caused by the mixture of the main lignocellulose-derived inhibitors and not in the detached individual effects. In addition, special attention is given to the heterogeneous composition of different hydrolysates, which is considered of major importance to the yeast tolerance response. Furthermore, rational metabolic engineering strategies successfully applied to yeast under industrial-like conditions are discussed.

Table 1 Main hemicellulose-derived compounds (monosaccharide and oligosaccharide) and degradation compounds (weak acids, furans, and phenolic compounds) derived from the pretreatment of lignocellulosic biomass

Raw material	Pretreatment (operational conditions)	Sugar ^a	Main inhibitor compounds derived from lignocellulosic pretreatment			Reference
			Weak acid ^b	Furan compounds ^c	Phenolic compounds	
Agricultural residues Wheat straw	Steam explosion (220 °C for 2.5 min)	4.4 g/L of X; 19.2 g/L of XOS	6.4 g/L of AA; 2.6 g/L of FA	1.8 g/L of F; 0.4 g/L of HMF	NA	Alvira et al. (2011)
	Steam explosion (210 °C for 2.5 min)	2.7 g/L of X; 20.7 g/L of XOS	4.0 g/L of AA; 1.7 g/L of FA	0.7 g/L of F; 0.2 g/L of HMF	NA	Costa et al. (2017)
Wheat straw	Autohydrolysis treatment ($S_0 = 3.92$); acid post-hydrolysis (165 °C, 0.5% H_2SO_4 , 168 min)	16 g/L of X	2.6 g/L of AA	0.13 g/L of HMF; 0.35 g/L of F	NA	Costa et al. (2017)
Sweet sorghum hybrid CSSH 45	Steam process (200 °C for 1.5 min)	6.7% of X	0.1% of GA; 0.2% of FA; 1.1% of AA; 0.1% of LeA	0.4% of F; 0.1% of HMF	3.4 g GAE/L	Damay et al. (2018)
	Steam process (215 °C for 1.5 min)	12.8% of X	0.2% of GA; 0.6% of FA; 2.6% of AA; 0.1% of LeA	1.2% of F; 0.1% of HMF	NA	
Vine pruning residue	Steam process (215 °C for 6 min)	5.2% of X	0.4% of GA; 0.9% of FA; 2.9% of AA; 0.4% of LeA	2.0% of F; 0.1% of HMF	5.3 g GAE/L	Jesus et al. (2017)
	Autohydrolysis treatment (180 °C for 60 min)	1.99 g/L of X; 17.22 g/L of XOS	2.66 g/L of AA; 6.10 g/L of AG	0.36 g/L of F; 0.66 g/L of HMF	2.35 g GAE/L; 47.54 mg/L of catechin; 34.70 mg/L of chlorogenic acid	Costa et al. (2017)
Corn cob	Autohydrolysis treatment (S_0); acid post-hydrolysis (165 °C, 0.5% H_2SO_4 , 168 min)	26 g/L of X	4 g/L of AA	0.2 g/L of HMF; 0.4 g/L of F	NA	Costa et al. (2017)
Corn cob	Wet oxidation treatment (195 °C for 15 min) with Na_2CO_3	1.3% of X	2.1 g/L of GA; 3.3 g/L of FA; 4.2 g/L of AA	NA	54 mg/L of 4-hydroxybenzaldehyde; 57 mg/L of vanillin	Varga et al. (2004)
Sugarcane bagasse	Wet oxidation treatment (195 °C for 15 min) with H_2SO_4	9% of X	1.8 g/L of GA; 2.9 g/L of FA; 2.8 g/L of AA	NA	102 mg/L of 4-hydroxybenzaldehyde; 74 mg/L of vanillin	Yu et al. (2018)
	Microwave-assisted pretreatment (0.5% H_2SO_4 , 140 °C for 5 min)	20.1%	0.05 g/L of FA; 1.69 g/L of AA	0.028 g/L of HMF; 0.64 g/L of F	NA	
Hardwood <i>Eucalyptus globulus</i> wood	Autohydrolysis treatment ($S_0 = 4.08$); acid post-hydrolysis (165 °C, 0.5% H_2SO_4 , 168 min)	16 g/L of X	6.2 g/L of AA	0.33 g/L of HMF; 1.66 g/L of F	2.01 g GAE/L	Costa et al. (2017)
<i>Paulownia tomentosa</i>	Autohydrolysis treatment ($S_0 = 4.19$); acid post-hydrolysis (165 °C, 0.5% H_2SO_4 , 168 min)	14 g/L of X	5.84 g/L of AA	0.72 g/L of HMF; 1.96 g/L of F	NA	Cunha et al. (2018)
<i>Acacia dealbata</i>	Non-isothermal autohydrolysis treatment at 217 °C	1.29% of X; 10% of XOS	0.89% of AA; 4.7% of AG	0.40% of F; 0.17% of HMF	NA	Yáñez et al. (2009)
Softwood Spruce	SO_2 -impregnated steam explosion	10.09 g/L of X	4.79 g/L of AA	1.57 g/L of F; 1.09 g/L of HMF	NA	Demeke et al. (2013a)
Lodgepole pine wood	SPORL pretreatment (165 °C for 75 min)	13.9% of GA; 14.5% of M; 9.7% of X	0.76% of AA	1.7 g/L of HMF; 1.1 g/L of F	NA	Zhou et al. (2013)
Spruce chips	SO_2 steam explosion (pH 2; 18 bar for 5–7 min)		3.52 g/L of AA	0.30 g/L of F; 1.26 g/L of HMF	0.05 mg/L of catechol; 0.128 mg/L of vanillin	Westman et al. (2012)

Table 1 (continued)

Raw material	Pretreatment (operational conditions)	Sugar ^a	Main inhibitor compounds derived from lignocellulosic pretreatment		Reference
			Weak acid ^b	Phenolic compounds	
Pine wood biomass	SO ₂ steam explosion (215 °C for 5 min)	9.2 g/L of G; 12.5 g/L of M; 2.5 g/L of Ga; 5.2 g/L of X NA	0.43 g/L of FA; 0.10 g/L of LA; 2.15 g/L of HMF; 1.18 g/L of F; 0.050 mg/L of vanillic acid; 0.022 mg/L of vanillin; 0.015 mg/L of benzoic acid	0.003 mg/L of 3,4-DHBA; 0.005 mg/L of 3-HBA; 0.050 mg/L of vanillic acid; 0.022 mg/L of vanillin; 0.015 mg/L of benzoic acid	Hawkins and Doran-Peterson (2011)

The hardness of hydrothermal treatments can be expressed in terms of “severity” (S_0), defined as the logarithm of the severity factor R_0 (Lavoie et al. 2010). Total phenolic compounds were measured as gallic acid equivalent (GAE) using the Folin-Ciocalteu method (Jesus et al. 2017); % g of component per 100 g of oven-dried raw material

^a Sugar: xylose (X), xylooligomers (XOS), galactose (Ga), glucose (G), and mannose (M)

^b Weak acids: acetic acid (AA), formic acid (FA), levulinic acid (LeA), lactic acid (LA), acetyl groups (AG), and glycolic acid (GA)

^c Furan compounds: furfural (F), hydroxymethylfurfural (HMF), and furfuroic acid (FuA)

Inhibitory effects on yeast during lignocellulosic fermentation

The overall metabolic and structural effects behind the negative effects of inhibitory compounds/process conditions on yeast growth and fermentation are listed on Table 2 and are further discussed below. Nevertheless, it should be taken into consideration that the specific effects of some of these inhibitors remain unknown or not well understood.

Intracellular acidification and ATP depletion

The effects of weak acids, mainly of acetic acid, on *S. cerevisiae* physiology and performance have been studied and recently reviewed (Palma et al. 2018). In the acidic pH conditions required for ethanol production from lignocellulosic biomass, weak acids enter the yeast cell in their protonated form (–COOH) and dissociate in the nearly pH-neutral cytoplasm, releasing hydrogen ions (H⁺) and leading to intracellular acidification (Ullah et al. 2012; Fig. 1 \triangleleft). To maintain intracellular pH homeostasis, this acidification is counteracted by the activity of the H⁺-ATPase, which exports H⁺ at the expense of ATP consumption (Fig. 1 \triangleleft). Furthermore, the anionic form of the acid is presumably exported by several multidrug resistance (MDR) transporters also contributing to ATP depletion in the yeast cell (Palma et al. 2018). In turn, ATP depletion will further limit the activity of ATPases, causing the dissipation of the transmembrane electrochemical gradient of protons, compromising secondary solute transport systems and the maintenance of ion homeostasis in the yeast cell (Serrano 1984). In addition, weak acids are also known to inhibit glycolytic enzymes, preventing ATP regeneration (Pampulha and Loureiro-Dias 1990) and leading to an energy drain.

Reactive oxygen species accumulation/oxidative stress

Weak acids are also known to cause oxidative stress, being the accumulation of reactive oxygen species (ROS; Fig. 1 \triangleleft) caused both by the increase of H⁺ in the cytosol and by the decrease of the ROS scavenger reduced glutathione (GSH) (Guo and Olsson 2014). The rate of ROS production is also known to be significantly increased at high temperatures as a consequence of heat stress (Morano et al. 2012). In yeast, ROS are neutralized by non-enzymatic and enzymatic processes, with these last requiring NADPH as a source of reduction equivalents (Herrero et al. 2008). To compensate NADPH oxidation, yeast gradually increases the influx through pentose phosphate and acetic acid pathways (Celton et al. 2012). Thereby, an increase in acetic acid production is stimulated at high temperatures, representing a synergistic effect that leads to the decrease of growth and ethanol production rate




Fig. 1 Main mechanisms of the *S. cerevisiae* response towards the presence of lignocellulose-derived inhibitors. Main negative stressor effects are identified by numbered triangles: (1) intracellular acidification, (2) ATP depletion, (3) ROS oxidative stress, (4) redox imbalance, and (5) cell wall and plasma membrane perturbations. Superscript numbered red circles nearby stressors indicate its main

negative effects. Superscript numbered green circles correspond to the counteract effects on the corresponding main negative effect. Black arrows represent transport of compounds and metabolic reactions; red arrows indicate negative effects; full green arrows represent positive activation/induction; dashed green arrows indicate counteract effects on the correspondent main negative stressor effect

(Woo et al. 2014). In a similar manner, furan aldehydes such as furfural and HMF also potentiate ROS generation by acting as thiol-reactive electrophiles and depleting GSH levels (Allen et al. 2010; Kim and Hahn 2013). Some phenolic compounds have also been reported to cause oxidative stress (Nguyen et al. 2014b); however, the mechanism behind their involvement in ROS accumulation is not yet understood. ROS accumulation in the yeast will ultimately result in damage at the mitochondrial and vacuolar membranes, the nuclear chromatin, and the actin cytoskeleton (Allen et al. 2010).

Redox imbalance

In the yeast cell, furans are converted into their corresponding less toxic alcohols, through reactions mediated by NAD(P)H-dependent oxidoreductases, which will ultimately lead to redox imbalance (Ask et al. 2013; Fig. 1). This decrease/drop in the reduction potential of the yeast intracellular compartment also contributes to oxidative stress, as NADPH is

required for the reduction of oxidized glutathione, thus also decreasing GSH levels. The yeast *S. cerevisiae* also has the capacity to detoxify phenolic compounds, converting them into less toxic derivatives, either through a series of decarboxylations and oxidations (Adeboye et al. 2015; Adeboye et al. 2017) or by NADPH-dependent reductions (Nguyen et al. 2014a; Wang et al. 2016), which may also contribute to a redox imbalance depending on the prevalence of phenolic compounds derived from the lignocellulose pretreatment.

Structural defects

In addition to these metabolic effects, the lignocellulose-derived inhibitors also cause structural changes in the yeast cell (Fig. 1), mainly in its cellular envelope. The presence of these compounds is known to lead to the reduction of plasma membrane stability and its selective permeability, by reducing its ergosterol content (Godinho et al. 2018) or by changing its protein-to-lipid ratio (Campos et al. 2009).

Table 2 Main metabolic and structural effects of the presence of lignocellulose-derived inhibitors in the yeast cell

Effects	Stress (reference)
Metabolic	
Redox imbalance	Furans (Ask et al. 2013) Phenolics (Adeboye et al. 2014, 2017; Nguyen et al. 2014a; Wang et al. 2016)
ROS accumulation	Furans (Kim and Hahn 2013) Weak acids (Guo and Olsson 2014; Woo et al. 2014) Phenolics (Nguyen et al. 2014b) High temperature (Woo et al. 2014)
Intracellular acidification (ATP depletion)	Weak acids (Ullah et al. 2012)
Inhibition of glycolytic enzymes (ATP depletion)	Weak acids (Pampulha and Loureiro-Dias 1990; Pearce et al. 2001) Sugar co-fermentation (Subtil and Boles 2012)
Structural	
Membrane and cell wall integrity	Weak acids (Godinho et al. 2018) Phenolics (Campos et al. 2009) High temperature (Verghese et al. 2012)
Organelle integrity	Weak acids (Pereira et al. 2010a; Verghese et al. 2012)
Macromolecule production and/or stability	Furans (Iwaki et al. 2013a) Phenolics (Iwaki et al. 2013b) High temperature (Foretek et al. 2016; Verghese et al. 2012)

Furthermore, the integrity and organization of the cell wall is also compromised by these inhibitors; e.g., weak acids are capable of increasing cell wall porosity and decreasing its robustness (Simões et al. 2006). The decrease on the integrity of the yeast cellular envelope significantly facilitates and increases the entry of inhibitory compounds into the yeast cell, synergistically contributing to their toxic effect. In fact, Ding and collaborators (2011) have observed that the severe effects of acetic acid on the yeast cell were potentiated by the presence of phenol and furfural, due to the loss of membrane integrity and metabolism inhibition. In fact, the synergistic effect between weak acids, furans, and phenolic compounds has been for long recognized as the main cause of the high toxicity of lignocellulosic hydrolysates, as the cumulative effect of the inhibitors present in a hydrolysate is far beyond that of the sum of their individual toxic effects (Ding et al. 2011; Keating et al. 2006; Klinke et al. 2003; Palmqvist et al. 1999).

Effects of high temperature

High temperature is one of the conditions required for simultaneous saccharification and fermentation from lignocellulosic biomass, and it can significantly affect yeast. Heat stress is known to disturb protein stability, cell membrane, and cytoskeleton structures, which leads to protein dysfunction, metabolic imbalances (Verghese et al. 2012), loss of metabolic activity (Woo et al. 2014), and defects in transfer RNA (tRNA) maturation by the accumulation of aberrant tRNA

processing intermediates upon shift of cells to high-temperature conditions (Foretek et al. 2016). Heat shock response is a fundamental cytoprotective pathway that enables yeast to cope with high-temperature stress, by activation of heat shock protein (HSP) synthesis (Verghese et al. 2012).

Contribution of modifications for xylose consumption to the inhibitory effects

Xylose consumption, by expression of heterologous pathways, on *S. cerevisiae* presents another hurdle on the production of second-generation bioethanol, as it has been described to increase yeast susceptibility to the inhibitory effects of the compounds present in lignocellulosic hydrolysates (Bellissimi et al. 2009). In fact, the genetic modifications used for xylose consumption can disturb the metabolic homeostasis of the yeast cell, decreasing its tolerance. For instance, it is known that expression of the *P. stipitis* XR/XDH pathway results in a redox imbalance caused by the co-factor difference between XR and XDH (while XR mainly uses NADPH, and XDH co-factor is NAD⁺) (Zhang et al. 2012), which may interfere with the metabolic effects of the inhibitory compounds, in addition to the undesirable accumulation of the by-product xylitol. Xylose uptake is mediated by hexose transporters in yeast, which are unspecific for pentose sugars (Subtil and Boles 2012). In low concentrations, glucose improves xylose uptake by activating these transporters; however, in higher concentrations, it outcompetes xylose, with high-glucose phosphorylation rates

repressing sugar co-consumption (Lane et al. 2018). Hexose and pentose catabolism converges at the level of phosphofruktokinase, and glucose limits glycolytic enzyme activity at this level (Subtil and Boles 2012). Also, after glucose depletion in a medium containing glucose and xylose (which occurs in lignocellulosic fermentations), cell growth and xylose consumption rate decrease sharply to values even lower than those in media containing xylose as the sole carbon source, and cells cease to respond to residual xylose, entering a new lag phase, named post-glucose effect lag phase (Wei et al. 2018). Intracellular xylose can, in fact, trigger a signal similar to carbon limitation in yeast cells actively metabolizing xylose, which causes low assimilation rates (Osiro et al. 2018). Xylose metabolism can also lead to downregulation of genes encoding gluconeogenic enzymes (Salusjärvi et al. 2006) and cause upregulation of genes involved in response to stress, starvation, DNA damage, and lipid metabolism by being forced to metabolize unconventional substrates (Gopinarayanan and Nair 2018).

End-product inhibition

On top of these stress factors inherent to efficient processing of lignocellulosic materials, the target product itself will, in most of the cases, affect negatively yeast cell metabolism. The most well-described end-product inhibition is ethanol, having pleiotropic effects on yeast cell (Deparis et al. 2017) affecting cell growth and viability (Pereira et al. 2011b) mainly by distressing cell wall and membrane integrity. An adequate medium supplementation partly counteracts the ethanol negative effects (Pereira et al. 2010b).

Mechanisms of yeast response to the presence of multiple inhibitors/hydrolysates

S. cerevisiae has developed several mechanisms to cope with the presence of lignocellulose-derived inhibitors and their effects (Fig. 1). Additionally, the yeast also exhibits responses towards the hurdles typical of lignocellulosic processes, such as high temperatures and xylose co-consumption (Fig. 1). As lignocellulosic materials are a platform to obtain several different compounds, the end-product inhibition response will not be addressed in here.

Oxidative stress response

One of the most toxic effects of the presence of lignocellulose-derived inhibitors on the yeast cell is oxidative stress, an imbalance between ROS generation and antioxidant response. The *YAP1* gene encodes a transcription factor, activated by the presence of both furans (Kim and Hahn 2013) and some phenolic compounds (Nguyen et al. 2014b), and is the major

regulon in oxidative stress response (Herrero et al. 2008). It induces expression of genes involved in the detoxification of superoxide anions (*SOD1*), reduction of hydrogen peroxide (*GPX2*, *CTT1*, *TSA1*), and thiol reduction (*TRX2*, *TRR1*), as well as expression of genes involved in the glutathione system (*GSH1*, *GSH2*, *GLR1*, *GRX1*, *YCF1*) (Hélène et al. 2000). *YAP1* also regulates the expression of other genes involved in response to several stressful conditions, such as MDR proteins (*FLR1*, *ATRI*) (Sundström et al. 2010) and HSPs (*SSA1*) (Maeta et al. 2004).

Furthermore, *YAP1* is known to induce *STB5* (Ouyang et al. 2011), a transcription factor that regulates most genes of the PPP, being a key player for NADPH regeneration required for oxidative stress response (Larochelle et al. 2006), but also for the detoxification of inhibitory compounds (Gorsich et al. 2006; Nguyen et al. 2014b). As already mentioned, in the presence of furfural and HMF, the yeast cell responds with the activity of NAD(P)H-dependent oxidoreductases to convert them into the less toxic furfuryl alcohol and furan dimethanol, respectively (Heer et al. 2009; Liu et al. 2008; Xianxian et al. 2015). In addition, being PPP the primary pathway for xylose metabolism, *STB5* regulation is important not only for tolerance towards inhibitory compounds but also for the consumption of alternative carbon sources present in lignocellulosic biomass (Kim et al. 2015). The detoxification of some phenolic compounds, involving genes such as *ALD5*, *PAD1*, *ATF1*, and *ATF2* (Adeboye et al. 2017), and several decarboxylation and oxidation reactions (Adeboye et al. 2015) could hypothetically counteract the redox imbalance created by the reduction of furan compounds. Nevertheless, the detoxification of other phenolic compounds, such as vanillin, involves NADPH-dependent reductases, and in this sense, the effects of the phenolic compounds in the redox homeostasis of the yeast will strongly depend on their chemical nature (Adeboye et al. 2014). Additionally, *S. cerevisiae* induces the synthesis of diverse molecules with antioxidant activity against heat-induced oxidative stress (Morano et al. 2012), with several molecules being identified as important for yeast response to heat stress, such as HSPs, H⁺-ATPases, ubiquitin, and antioxidant enzymes (Gao et al. 2016).

Structural response: cell membrane

Accumulation of trehalose is another defense mechanism activated by oxidative stress, where it plays an important protective role in the maintenance of the integrity of the cell membrane (Alvarez-Peral et al. 2002), probably by stabilization of membrane proteins (Jain and Roy 2009). In this sense, trehalose accumulation has been described to be activated in response to membrane-disrupting stresses, such as high temperatures (Mensonides et al. 2014) and exposure to weak acids (Guo and Olsson 2014). Accordingly, the genes involved in trehalose synthesis have been found to be regulated by the *MSN2/4*

transcription factors, which are activated upon oxidative stress (Gasch et al. 2000; Hasan et al. 2002), but also by stressors such as high temperature and low pH (Causton et al. 2001).

Another factor that has been identified as a determinant for yeast tolerance is its capacity to largely rearrange the lipid composition of the plasma membrane (e.g., sphingolipids and sterols) (Lindberg et al. 2013). Sphingolipid content was found to be increased in response to acetic acid stress (Lindberg et al. 2013), and the upregulation of sphingolipid biosynthesis was described to be mediated by the TORC2-Ypk1 signaling complex (Roelants et al. 2011), which is activated not only by acetic acid (Guerreiro et al. 2016) but also by heat stress (Sun et al. 2012). Ergosterol, a major constituent of the yeast plasma membrane, is another molecule required to maintain membrane integrity. In fact, a possible interaction has been suggested between ergosterol biosynthesis and the oxidative stress response (Higgins et al. 2003). Furthermore, several genes from the ergosterol biosynthetic pathway were upregulated in response to acetic acid stress, as well as *PDR18*, which was found to have a physiological role in ergosterol transport and proper incorporation into the plasma membrane, increasing its lipid order and decreasing the non-specific membrane permeability (Godinho et al. 2018). *PDR16* (positively regulated by YAP1) and *PDR17* have also been described to be important for lipid biosynthesis (ergosterol and phospholipids, respectively), not only playing an important role on plasma membrane integrity but also controlling lipid content in various compartments of the cell, providing mechanisms for multidrug resistance (van den Hazel et al. 1999). In fact, the expression of genes of the pleiotropic drug resistance (PDR) family was found to be enhanced in response to the presence of furfural and HMF (Liu et al. 2008; Ma and Liu 2010). The PDR family mainly consists of membrane- and transport-related proteins, such as the ATP-binding cassette (ABC) transporters, including the weak acid-inducible *PDR12* which contributes for the efflux of anions (Ullah et al. 2012). In fact, *PDR12* is regulated by *WAR1*, a transcription factor that is activated by phosphorylation in the presence of weak acids (Frohner et al. 2010; Gregori et al. 2008; Kren et al. 2003). Nevertheless, *Pdr12* role in response to weak acid stress is not common to weak acids in general, as its absence leads to high susceptibility to the more lipophilic weak acids but seems to be advantageous for tolerance to shorter acids, such as acetic and formic (Nygård et al. 2014). In fact, *TPO2* and *TPO3*, encoding MDR transporters of the major facilitator superfamily, have been found to confer resistance to acetic, propionic, benzoic, and octanoic acids (with a slightly more evident effect for the more hydrophilic acids), presumably through the active export of the counter ions (Fernandes et al. 2005). More recently, *TRK1*, encoding for a high-affinity potassium transporter, has been found to have a detrimental effect in the yeast response to formic acid, presumably by contributing to the influx of this acid into the cell

(Henriques et al. 2017). The fact that *TRK1* is a determinant of yeast tolerance towards acetic acid is another example of how diverse weak acids may activate different response mechanisms. Accordingly, it has been proposed that dissimilar weak acids may activate unique tolerance mechanisms: while less lipophilic acids (acetate and propionate) were found to mainly regulate membrane-associated transport processes, the transcriptional response to more strongly lipophilic acids (benzoate and sorbate) mainly regulates genes related to the cell wall (Abbott et al. 2007).

Structural response: cell wall

The cell wall integrity (CWI) signaling pathway in *S. cerevisiae* is activated in response to several forms of cell wall stress and acts on cell wall remodeling (through control of wall biosynthetic enzymes), transcriptional regulation of cell wall-related genes, and organization of actin cytoskeleton (Levin 2005, 2011). CWI pathway has been found to play an important role in yeast tolerance towards major components of lignocellulosic hydrolysates, such as acetic acid (Nishida et al. 2014), furfural (Liu et al. 2018), and HMF (Liu et al. 2018; Zhou et al. 2014). Weak acid stress is also known to cause the activation of HAA1, a transcription factor responsible for yeast adaptation and tolerance to short-chain weak acids, such as acetic and formic acids (Fernandes et al. 2005; Henriques et al. 2017). HAA1 has been found to transcriptionally regulate cell wall proteins, such as SPI1 and YGP1; proteins from the plasma membrane, such as the MDR transporters TPO2 and TPO3; and proteins involved in the biosynthesis of lipids (contributing to the integrity of the plasma membrane) (Fernandes et al. 2005; Mira et al. 2010; Simões et al. 2006). More recently, HAA1 has been hypothesized to play a role in acetic acid tolerance through the activation of the CWI pathway (Cunha et al. 2018).

ATP and NADH regeneration

Another inhibitory effect occurring during lignocellulosic ethanol fermentation is ATP depletion, mainly caused by the activity of ATP-dependent pumps required to cope with the intracellular acidification caused by weak acids, in particular the plasma and vacuolar H⁺-ATPases and multidrug efflux pumps. In this situation, the yeast cell adjusts its carbon flux distribution between respiratory and fermentative growth to achieve energy homeostasis through optimal ATP regeneration (Guo and Olsson 2014). Furthermore, the trehalose synthase (TPS1) has been found to be essential to maintain ATP levels during heat shock (Petitjean et al. 2015). MSN2/4 transcription factors, known to regulate trehalose biosynthesis genes, were also reported to induce glycolysis, increasing the levels of acetyl-CoA, an essential metabolite to generate ATP in the tricarboxylic acid (TCA) cycle and to promote yeast cell growth and proliferation (Kuang et al. 2017).

Additionally, the presence of furan compounds was found to result in the activation of glycolysis and TCA cycle, contributing to both ATP and NADH regenerations (Lin et al. 2009).

Successful cases of yeast robustness improvement for industrial-like conditions

Industrial-derived tolerant strains

The use of *S. cerevisiae* strains isolated from industrial harsh conditions (such as high sugar and ethanol concentrations, elevated temperatures, pH variations, and presence of toxic compounds) for the production of second-generation bioethanol has been receiving increased attention in the last years (Della-Bianca and Gombert 2013). These isolated strains have shown superior abilities than laboratory strains, with the differences in fermentation performance being related to metabolic activity, not only with sugar consumption and ethanol production (Pereira et al. 2010c) but also with furan conversion (Brandberg et al. 2004; Pereira et al. 2014a). Interestingly, the better fermentation performance of industrial isolates compared to laboratory strains in very high-gravity conditions was related with an increased accumulated content of sterols, glycogen, and trehalose in the industrial isolates (Pereira et al. 2011b). On the other hand, under second-generation inhibitory conditions, the *S. cerevisiae* ATCC96581 strain (isolated from spent sulfite liquor at Swedish pulp plant) converted almost completely the furfural of spruce hydrolysate, whereas the laboratory strain CBS 8066 only detoxified 25% (Brandberg et al. 2004). This fact could be explained by a higher activity of alcohol dehydrogenase responsible for the conversion of furfural into less toxic alcohols. Pereira and co-workers (2014a) also reported a faster bioconversion/detoxification of furfural and HMF in eucalyptus hydrolysate by two industrial strains, PE-2 and flocculating CCUG53310 isolated from first- and second-generation bioethanol industries, respectively. The authors concluded that the ability for detoxification of furan compounds is dependent on strain background, which is determinant for an efficient ethanol production (Pereira et al. 2014a). Moreover, the flocculant character of strains, which has well-known process-related advantages (Gomes et al. 2012), has been also related to inhibitor tolerance (Purwadi et al. 2007; Westman et al. 2014). The mechanism and robustness of the flocculating CCUG53310 strain have been investigated and compared with the laboratorial *S. cerevisiae* CBS 8066 (Westman et al. 2012). The flocculant strain showed higher tolerance to the inhibitors present in a spruce hydrolysate, even though it presented lower expression levels of *YAPI*, *ATRI*, and *FLRI* genes (known to confer resistance to lignocellulose-derived inhibitors) than the laboratorial strain, highlighting flocculation as a physiological trait determinant of yeast tolerance. The authors also hypothesized that the lower expression

of *YAPI* (normally activated in response to oxidative stress) in the CCUG53310 strain indicated that flocculation may prevent ROS accumulation, through mechanisms that are still not elucidated but are likely related with a reduction of toxic concentrations around the cell and in the cell interior.

Therefore, the selection of robust yeast chassis for metabolic engineering purposes (such as xylose consumption) shows a further edge for the lignocellulose-to-ethanol fermentations (Costa et al. 2017). In fact, Romani et al. (2015) expressed a xylose consumption pathway in three different *S. cerevisiae* strains: the laboratorial CEN.PK113-5D and two industrial isolates from first-generation bioethanol plants (PE-2 and CAT-1), and observed that the two industrial strains presented higher xylose consumption and ethanol production than the strain with laboratorial background, both in synthetic media and in a corn cob hydrolysate. Kim et al. (2017b) also evaluated the host strain background of a haploid derivative of the industrial strain *S. cerevisiae* ATCC 4124 and of the laboratory D452-2 strain by genetically engineering them for xylose consumption. They observed that the industrial-derived strain had a superior fermentative performance in a *Miscanthus* hydrolysate (superior efficiency of xylose fermentation and ethanol production) than the laboratorial strain containing the same genetic modification, highlighting the importance of selecting a naturally robust host strain. In addition, Costa et al. (2017) showed differences among metabolically engineered industrial strains for xylose consumption depending of the hemicellulosic hydrolysate used.

Moreover, these desirable traits for inhibitor tolerance of the industrial isolates can still be improved through metabolic engineering, mutagenesis, genome shuffling, or evolutionary engineering. The work developed by Liu et al. (2005, 2008, 2018) and Liu and Moon (2009) is a clear example of the development of new improved strains. The industrial *S. cerevisiae* NRRL Y-12632, isolated from the brewer's top yeast in Netherlands in 1925, was subjected to evolutionary engineering in HMF- and furfural-containing media, resulting in the reduction of lag phase, improvement of glucose consumption, and ethanol production in media containing these inhibitors. It was further described that these improved traits resulted from determinant yeast response mechanisms, such as enhanced expression of PDR gene family, increased NAD(P)H-dependent aldehyde reduction activities, increased expression of genes from glycolysis, and PPP for NAD(P)H regeneration and robust cell wall integrity pathway.

Rational metabolic engineering strategies to improve tolerance to lignocellulosic hydrolysates

The use of industrial strains as hosts for metabolic engineering is a promising approach for the feasibility of second-generation bioethanol industry. An extensive knowledge of the mechanisms required for the yeast response towards lignocellulose-

derived inhibitors has been guiding the use of several strategies to develop *S. cerevisiae* strains capable of withstanding acute stresses with improved growth/fermentative performances (Table 3).

Several of these strategies have focused on the detoxification of inhibitory compounds. Jayakody and collaborators (2018) improved the fermentation of a *Miscanthus* hydrolysate by overexpressing of *GRE2* (encoding a NADPH-dependent aldehyde reductase), increasing the yeast capacity to detoxify aldehyde inhibitors, such as vanillin and glycolaldehyde. The overexpression of *PRS3*, responsible for the synthesis of PRPP (a precursor of nucleotide and histidine biosynthesis), was found to improve yeast fermentation rates and productivities in different lignocellulosic hydrolysates, through a hypothesized increase in NADH regeneration which facilitates detoxification furans (Cunha et al. 2015). Nevertheless, it should be noted that this positive effect was dependent of the strain background and composition of the fermentation media, highlighting the importance of the selection of yeast chassis and fermentation conditions for effective metabolic engineering (Cunha et al. 2015). Detoxification of phenolic compounds has also been addressed to improve yeast tolerance: the expression of a laccase from the white rot fungus *Trametes versicolor* in a laboratorial *S. cerevisiae* strain has increased the yeast ability to convert coniferyl aldehyde into less toxic compounds, increasing yeast growth and ethanol production in a dilute acid spruce hydrolysate (Larsson et al. 2001). Wallace-Salinas and collaborators (2014) decreased the lag phase and improved the growth rate of an Ethanol Red strain (previously modified for xylose consumption) (Demeke et al. 2013a, b), in a spruce hydrolysate, by overexpressing *YAP1*, a transcription factor involved in oxidative stress response and tolerance. Furthermore, these authors also overexpressed *MCR1*, coding for the mitochondrial NADH-cytochrome b5 reductase, resulting in a faster furaldehyde reduction capacity with positive effects on yeast growth (similar to the ones resultant from *YAP1* overexpression). Nevertheless, no cumulative effect of the simultaneous overexpression of these two genes on yeast tolerance was observed (Wallace-Salinas et al. 2014).

Other studies have also attempted to improve yeast tolerance together with xylose consumption capacity. In fact, a haploid derivative of an industrial strain, isolated from a molasses distillery, was modified for xylose consumption with the XR/XDH pathway and for acetate consumption by expression of a NADH-dependent acetate reduction pathway (*adhE* gene from *Escherichia coli* coding for an acetylating acetaldehyde dehydrogenase) (Kim et al. 2017b). This later modification not only allowed the in situ detoxification of acetic acid but also increased intracellular NAD⁺ levels, potentiating XDH activity and reducing xylitol accumulation, leading to a higher ethanol yield. Hasunuma et al. (2014) also improved ethanol production from wheat straw-derived xylose (in an industrial strain also expressing hemicellulolytic enzymes) through the

overexpression of *TAL1* and *FDH1* and expression of a mutant NADH-dependent *ADH1*, which resulted in formate detoxification and faster detoxification of furfural, leading to a higher regeneration of NAD⁺ co-factor, improving the XR/XDH consumption pathway. More recently, *HAA1* (encoding a transcription factor involved in adaptation and tolerance to weak acid stress) and *PRS3* (encoding a 5-phospho-ribosyl-1(alpha)-pyrophosphate synthetase that synthesizes PRPP, which is required for nucleotide, histidine, and tryptophan biosynthesis) have been expressed in a first-generation bioethanol strain (PE-2), previously modified for xylose consumption, improving its adaptation to a non-detoxified Paulownia hydrolysate (Cunha et al. 2018). Furthermore, the simultaneous overexpression of both genes had a cumulative positive effect on yeast growth, and expression of both *HAA1* and *PRS3* was found to play a role in yeast cell wall integrity.

These successful strategies show that a thorough knowledge of the mechanisms involved in yeast response towards the presence of inhibitory compounds is a determinant for the development of tolerant strains to attain an efficient and economical production of lignocellulosic bioethanol.

Final remarks and future perspectives

The lignocellulosic process-derived stress factors lead to negative effects in the yeast cell at molecular, metabolic, and structural levels, being the most noteworthy, intracellular acidification, ATP depletion, ROS-induced oxidative stress, redox imbalance, and cell wall and plasma membrane perturbations. In order to cope with these conditions, the cell falls back on several global mechanisms that counteract their synergistic negative effects. In spite of their complexity, some of these mechanisms are nowadays relatively well described and linked with successful cases of yeast engineering. Nonetheless, the majority of studies regarding this subject use laboratorial yeast strains and focus on the effect and response to a single inhibitor. As depicted in this review, in process-like conditions, the synergetic effect of the presence of several inhibitors is of major influence in the process and should always be considered and evaluated in order to efficiently develop lignocellulosic hydrolysate-tolerant strains. Furthermore, the selection of chassis' strains for metabolic engineering strategies should be regarded as a crucial step to attain more robust and efficient strains. In fact, industrial isolates has been receiving growing attention in this field, as they naturally present advantageous traits (such as higher capacity for inhibitor tolerance/detoxification, thermotolerance, faster sugar consumption) that could represent a leverage for the attainment of efficient second-generation bioethanol processes. However, metabolic engineering of these industrial strains still poses some constrains that are being overcome by the presently available molecular toolbox for *S. cerevisiae* (in constant evolution), which facilitates the development of highly engineered

Table 3 Successful cases of rational metabolic engineering of *S. cerevisiae* for improved tolerance towards lignocellulosic hydrolysates

Modification	<i>S. cerevisiae</i> strain	Lignocellulosic hydrolysate	Effect	Mechanism	Reference
Laboratory strains					
Expression of laccase from <i>Trametes versicolor</i> and overexpression of <i>SSO2</i>	INVSC1 (MATa his3-Δ1 leu2 trp1-289 ura3-52)	Dilute acid spruce hydrolysate in g/L: 2.8 acetic acid, 0.7 formic acid, 1.1 levulinic acid, 1.4 furfural, 2.3 HMF, and 2.9 phenolic compounds	Faster growth (μ_{max} increased from 0.000 to 0.012 h ⁻¹) and ethanol formation (ethanol yield increased from 0.02 to 0.44 g/g of total sugars)	Improved detoxification of phenolic compounds	Larsson et al. (2001)
Overexpression of <i>YAP1</i>	INVSC1 (MATa his3-Δ1 leu2 trp1-289 ura3-52)	Diluted spruce hydrolysate	Improved ethanol productivity (0.17 g/L/h vs. 0.05 g/L/h of the control)	<i>YAP1</i> plays an important role in the oxidative stress response	Ariksson et al. (2010)
Expression of the <i>adhE</i> gene from <i>E. coli</i> (coding for acetylating acetaldehyde dehydrogenase) and of the mutant <i>Salmonella</i> ACS gene (coding for a acetyl-CoA synthetase)	D452-2 (MATα, leu2, his3, ura3, and can1)	<i>Miscanthus</i> hydrolysate in g/L: 20 glucose, 50 xylose, 10 acetate, 1 HMF, and 2 furfural	Acetate consumption and faster xylose consumption	In situ detoxification of acetic acid	Zhang et al. (2016)
Xylose consumption: <i>XYL1</i> , <i>XYL2</i> , and <i>XYL3</i> , evolutionary engineered in xylose-containing media and deletion of <i>PHO13</i> and <i>ALD6</i>			Increased ethanol yield (18.4%) and decreased glycerol and xylitol yields (40.3%)	Decrease of redox imbalance of xylose consumption pathway	
Overexpression of <i>SPE3</i> and deletion of <i>TPO1</i> and <i>OAZ1</i>	D452-2 (MATα, leu2, his3, ura3, and can1)	Corn stover hydrolysate in g/L: 3.3 acetic acid, 0.8 HMF, and 0.4 furfural	Improved ethanol productivity (14% higher than the control strain)	Spermidine extends the lifespan of <i>S. cerevisiae</i> (Eisenberg et al. 2009)	Kim et al. (2017a)
Overexpression of <i>YAP1</i> , <i>STB5</i> , <i>WARI</i> , <i>PDR8</i> , <i>CAT8</i> , <i>PUT3</i> , and <i>GZF3</i> , separately	BY4741 (MATa, his3Δ1, leu2Δ0, met15Δ0, ura3Δ0)	Sugarcane bagasse hydrolysate in g/L: 0.89 furfural, 0.11 HMF, 1.4 acetic acid, 0.03 formic acid, and 0.05 levulinic acid	Enhanced relative growth rates: 2.85- and 2.75-fold higher than those of the control strain in bagasse and spruce hydrolysate	Transcription factors involved in oxidative stress (<i>YAP1</i> and <i>STB5</i>), acid stress adaptation (<i>WARI</i>), pleiotropic drug resistance (<i>PDR8</i>), carbon source responsiveness (<i>CAT1</i>), amino acid biosynthesis (<i>PUT3</i>), and nitrogen catabolism (<i>GZF3</i>)	Wu et al. (2017)
Overexpression of <i>GRE2</i>	D452-2 (MATα, leu2, his3, ura3, and can1)	Spruce hydrolysate in g/L: 0.36 furfural, 0.03 HMF, 0.72 acetic acid, 0.27 formic acid, and 0.12 levulinic acid	Growth in the toxic hydrolysate at low inoculum (vs. no growth of the control strain)	Possible detoxification of glycolaldehyde and other major inhibitory compounds	Jayakody et al. (2018)
Xylose consumption: <i>XYL1</i> , <i>XYL2</i> , and <i>XYL3</i> , evolutionary engineered in xylose-containing media, and knockout <i>ALD6</i>			Improved xylose consumption and ethanol yields at higher inoculum	<i>GRE2</i> also converts vanillin into the less toxic vanillin alcohol through an NADPH-dependent reaction	
Overexpression of <i>TRX1</i> (coding for thioredoxin)	BY4742 (MATa, his3Δ1, leu2Δ0, lys2Δ0, ura3Δ0)	Diluted bagasse hydrolysate in g/L: 44 glucose, 5.8 xylose, 4.1 acetic acid, 0.6 furfural, and 0.2 HMF	Higher ethanol titer (10.50 g/L vs. 8.61 g/L), yield (0.30 g/L vs. 0.23 g/L), and productivity (0.46 g/L/h vs. 0.30 g/L/h) than the control	Maintenance of energy and redox homeostasis and minimization of stress-induced cell damages	Unrean et al. (2018)
Industrial strains				Increased levels of trehalose, fatty acids, GABA, and putrescine provided additional defense against oxidative and redox stresses	
				Detoxification of formate (<i>FDH1</i>)	

Table 3 (continued)

Modification	<i>S. cerevisiae</i> strain	Lignocellulosic hydrolysate	Effect	Mechanism	Reference
Overexpression of <i>FDHI</i> and <i>TAL1</i> and expression of a mutant NADH-dependent <i>ADHI</i> Xylose consumption: <i>Scheffersomyces stipitis</i> <i>Xyl1</i> and <i>Xyl2</i> and <i>S. cerevisiae</i> <i>XKS1</i> Hemicellulolytic enzymes: <i>Trichoderma reesei</i> <i>XYNII</i> , <i>Aspergillus oryzae</i> <i>XylA</i> , and <i>Aspergillus aculeatus</i> <i>BGLI</i>	Sun049, obtained from Suntory Limited (brewing and distilling company)	Rice straw hydrolysate in mM: 17.6 formate, 28.6 acetate, 0.3 vanillin, 12.6 furfural, and 0.9 5-HMF	2.7-fold higher ethanol titers and improved xylose consumption	Faster furfural detoxification, regenerating NAD ⁺ which improved ethanol yield from xylose (<i>ADHI</i>) <i>TAL1</i> overexpression improves PPP	Hasunuma et al. (2014)
Overexpression of <i>YAP1</i> and <i>MCRI</i> , separately Previous modifications: GSE16 (<i>XylA</i> ; <i>XKS1</i> ; <i>TAL1</i> ; <i>TKL1</i> ; <i>RPE1</i> ; <i>RKI1</i> ; <i>HXT7</i> ; <i>AraF</i> ; <i>AraA</i> ; <i>AraB</i> ; <i>AraD</i> ; <i>TAL2</i> ; <i>TKL2</i> + mutagenesis (<i>EMS</i>) + genome shuffling and evolutionary adaptation + backcrossing with a haploid segregant of Ethanol Red that is tolerant towards acetic acid)	Ethanol Red, a strain currently commonly used in industrial bioethanol fermentations	Spruce hydrolysate in g/L: 3.7 acetate, 0.96 HMF, and 0.78 furfural	Decreased lag phase (9 h to 5.3 h) and improved growth rate (around 60% higher)	Faster furfuraldehyde reduction capacity	Wallace-Salinas et al. (2014)
Overexpression of <i>PRS3</i>	PE2 and CCUG53310, isolated from the first- and second-generation bioethanol plants, respectively	<i>Eucalyptus globulus</i> wood hydrolysate in g/L: 2.1 acetic acid, 1.4 furfural, and 0.25 HMF Corn cob hydrolysate in g/L: 1.6 acetic acid, 1.6 furfural, and 0.12 HMF <i>Miscanthus</i> hydrolysate	Improved fermentation rate (up to 32%) and productivity (up to 48%) in the different hydrolysates	Possible increase in redox balance through NADH regeneration for furfural and HMF detoxification	Cunha et al. (2015)
Expression of the <i>adhE</i> gene from <i>E. coli</i> (coding for acetylating acetaldehyde dehydrogenase)	Haploid derivative of ATCC 4124 strain, isolated from a molasses distillery		Higher ethanol yield and lower by-product yield	In situ detoxification of acetic acid Decrease of redox imbalance of xylose consumption pathway	Kim et al. (2017b)
Xylose consumption: expression of <i>XYL1</i> , <i>XYL2</i> , and <i>XYL3</i> from <i>S. stipitis</i> and deletion of <i>PHO13</i> and <i>ALD6</i>	PE-2, isolated from first-generation bioethanol plant	Paulownia hydrolysate in g/L: 5.84 acetic acid, 1.96 furfural, and 0.719 HMF	Improved yeast adaptation to non-detoxified hydrolysate with high acetic acid content, resulting in higher ethanol titers (≥ 12%)	Increased robustness of yeast cell wall when challenged with acetic acid stress, caused by a possible involvement of <i>HAA1</i> and/or <i>PRS3</i> in the modulation of the cell wall integrity pathway	Cunha et al. (2018)
Overexpression of <i>HAA1</i> and/or <i>PRS3</i>					
Xylose consumption: expression of <i>XYL1</i> and <i>XYL2</i> from <i>S. stipitis</i> , overexpression of <i>TAL1</i> and <i>XKS</i> , and deletion of <i>GRE3</i>					

yeast strains. Accordingly, more recent studies have been using industrial yeast and lignocellulosic hydrolysates to develop more tolerant strains. Nevertheless, there is a lack of fundamental understanding regarding the response mechanisms that confer higher tolerance and robustness to these industrial isolates, being a subject requiring further investigation. As the complexity of yeast cell response is unraveled, an increasing number of metabolic engineering strategies will become successful, feeding back the accumulated knowledge. Nowadays, works to improve yeast tolerance still mainly focus on only one part of the inhibitory effects (such as oxidative stress or specific inhibitor detoxification). Due to the complexity of the multifactorial yeast tolerance to stress and in order to be effective, metabolic engineering strategies should be rationally designed to simultaneously overcome all the stresses imposed by the lignocellulosic hydrolysates. Additionally, the heterogeneity of lignocellulosic hydrolysates (dependent on the raw material and pretreatments used) should also be taken into consideration, as possible synergetic and antagonistic effects may arise from different inhibitory compositions and trigger different yeast responses. Taken together, this knowledge can unlock a wide range of strategies to develop tailor-made *S. cerevisiae* strains through rational metabolic engineering approaches for industrial processes, ultimately resulting in improved robustness when challenged in lignocellulosic hydrolysates, greatly contributing to the development of sustainable growth based on a bioeconomy.

Funding This study was supported by the Portuguese Foundation for Science and Technology (FCT) by the strategic funding of UID/BIO/04469/2013 unit, MIT Portugal Program (Ph.D. grant PD/BD/128247/2016 to Joana T. Cunha), Ph.D. grant SFRH/BD/130739/2017 to Carlos E. Costa, COMPETE 2020 (POCI-01-0145-FEDER-006684), BioTecNorte operation (NORTE-01-0145-FEDER-000004), YeasTempTation (ERA-IB-2-6/0001/2014), and MultiBiorefinery project (POCI-01-0145-FEDER-016403). Funding by the Institute for Bioengineering and Biosciences (IBB) from FCT (UID/BIO/04565/2013) and from Programa Operacional Regional de Lisboa 2020 (Project N. 007317) was also received.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

References

- Abbott DA, Knijnenburg TA, de Poorter LMI, Reinders MJT, Pronk JT, van Maris AJA (2007) Generic and specific transcriptional responses to different weak organic acids in anaerobic chemostat cultures of *Saccharomyces cerevisiae*. *FEMS Yeast Res* 7(6):819–833. <https://doi.org/10.1111/j.1567-1364.2007.00242.x>
- Adeboye PT, Bettiga M, Olsson L (2014) The chemical nature of phenolic compounds determines their toxicity and induces distinct physiological responses in *Saccharomyces cerevisiae* in lignocellulose hydrolysates. *AMB Express* 4:46–46. <https://doi.org/10.1186/s13568-014-0046-7>
- Adeboye PT, Bettiga M, Aldaeus F, Larsson PT, Olsson L (2015) Catabolism of coniferyl aldehyde, ferulic acid and p-coumaric acid by *Saccharomyces cerevisiae* yields less toxic products. *Microb Cell Factories* 14:149. <https://doi.org/10.1186/s12934-015-0338-x>
- Adeboye PT, Bettiga M, Olsson L (2017) *ALD5*, *PAD1*, *ATF1* and *ATF2* facilitate the catabolism of coniferyl aldehyde, ferulic acid and p-coumaric acid in *Saccharomyces cerevisiae*. *Sci Rep* 7:42635. <https://doi.org/10.1038/srep42635>
- Allen SA, Clark W, McCaffery JM, Cai Z, Lancot A, Slininger PJ, Liu ZL, Gorsich SW (2010) Furfural induces reactive oxygen species accumulation and cellular damage in *Saccharomyces cerevisiae*. *Biotechnol Biofuels* 3:2. <https://doi.org/10.1186/1754-6834-3-2>
- Alriksson B, Horváth IS, Jönsson L (2010) Overexpression of *Saccharomyces cerevisiae* transcription factor and multidrug resistance genes conveys enhanced resistance to lignocellulose-derived fermentation inhibitors. *Process Biochem* 45(2):264–271. <https://doi.org/10.1016/j.procbio.2009.09.016>
- Alvarez-Peral FJ, Zaragoza O, Pedreno Y, Argüelles J-C (2002) Protective role of trehalose during severe oxidative stress caused by hydrogen peroxide and the adaptive oxidative stress response in *Candida albicans*. *Microbiol* 148(8):2599–2606. <https://doi.org/10.1099/00221287-148-8-2599>
- Alvira P, Negro MJ, Ballesteros M (2011) Effect of endoxylanase and α -L-arabinofuranosidase supplementation on the enzymatic hydrolysis of steam exploded wheat straw. *Bioresour Technol* 102(6):4552–4558. <https://doi.org/10.1016/j.biortech.2010.12.112>
- Ask M, Bettiga M, Mapelli V, Olsson L (2013) The influence of HMF and furfural on redox-balance and energy-state of xylose-utilizing *Saccharomyces cerevisiae*. *Biotechnol Biofuels* 6(1):22. <https://doi.org/10.1186/1754-6834-6-22>
- Bellissimi E, van Dijken JP, Pronk JT, van Maris AJ (2009) Effects of acetic acid on the kinetics of xylose fermentation by an engineered, xylose-isomerase-based *Saccharomyces cerevisiae* strain. *FEMS Yeast Res* 9(3):358–364. <https://doi.org/10.1111/j.1567-1364.2009.00487.x>
- Brandberg T, Franzén CJ, Gustafsson L (2004) The fermentation performance of nine strains of *Saccharomyces cerevisiae* in batch and fed-batch cultures in dilute-acid wood hydrolysate. *J Biosci Bioeng* 98:122–125. [https://doi.org/10.1016/S1389-1723\(04\)70252-2](https://doi.org/10.1016/S1389-1723(04)70252-2)
- Campos FM, Couto JA, Figueiredo AR, Toth IV, Rangel AO, Hogg TA (2009) Cell membrane damage induced by phenolic acids on wine lactic acid bacteria. *Int J Food Microbiol* 135(2):144–151. <https://doi.org/10.1016/j.ijfoodmicro.2009.07.031>
- Causton HC, Ren B, Koh SS, Harbison CT, Kanin E, Jennings EG, Lee TI, True HL, Lander ES, Young RA (2001) Remodeling of yeast genome expression in response to environmental changes. *Mol Biol Cell* 12(2):323–337. <https://doi.org/10.1091/mbc.12.2.323>
- Celton M, Goelzer A, Camarasa C, Fromion V, Dequin S (2012) A constraint-based model analysis of the metabolic consequences of increased NADPH oxidation in *Saccharomyces cerevisiae*. *Metab Eng* 14:366–379. <https://doi.org/10.1016/j.ymben.2012.03.008>
- Costa CE, Romani A, Cunha JT, Johansson B, Domingues L (2017) Integrated approach for selecting efficient *Saccharomyces cerevisiae* for industrial lignocellulosic fermentations: importance of yeast chassis linked to process conditions. *Bioresour Technol* 227:24–34. <https://doi.org/10.1016/j.biortech.2016.12.016>
- Cunha JT, Aguiar TQ, Romani A, Oliveira C, Domingues L (2015) Contribution of *PRS3*, *RPB4* and *ZWF1* to the resistance of industrial *Saccharomyces cerevisiae* CCUG53310 and PE-2 strains to lignocellulosic hydrolysate-derived inhibitors. *Bioresour Technol* 191:7–16. <https://doi.org/10.1016/j.biortech.2015.05.006>
- Cunha JT, Costa CE, Ferraz L, Romani A, Johansson B, Sa-Correia I, Domingues L (2018) *HAA1* and *PRS3* overexpression boosts yeast

- tolerance towards acetic acid improving xylose or glucose consumption: unravelling the underlying mechanisms. *Appl Microbiol Biotechnol* 102(10):4589–4600. <https://doi.org/10.1007/s00253-018-8955-z>
- Damay J, Boboescu I-Z, Duret X, Lalonde O, Lavoie J-M (2018) A novel hybrid first and second generation hemicellulosic bioethanol production process through steam treatment of dried sorghum biomass. *Bioresour Technol* 263:103–111. <https://doi.org/10.1016/j.biortech.2018.04.045>
- Della-Bianca BE, Gombert AK (2013) Stress tolerance and growth physiology of yeast strains from the Brazilian fuel ethanol industry. *Antonie Van Leeuwenhoek* 104:1083–1095. <https://doi.org/10.1007/s10482-013-0030-2>
- Demeke MM, Dietz H, Li Y, Foulquié-Moreno MR, Mutturi S, Deprez S, Den Abt T, Bonini BM, Liden G, Dumortier F, Verplaetse A, Boles E, Thevelein JM (2013a) Development of a D-xylose fermenting and inhibitor tolerant industrial *Saccharomyces cerevisiae* strain with high performance in lignocellulose hydrolysates using metabolic and evolutionary engineering. *Biotechnol Biofuels* 6(1):89. <https://doi.org/10.1186/1754-6834-6-89>
- Demeke MM, Dumortier F, Li Y, Broeckx T, Foulquié-Moreno MR, Thevelein JM (2013b) Combining inhibitor tolerance and D-xylose fermentation in industrial *Saccharomyces cerevisiae* for efficient lignocellulose-based bioethanol production. *Biotechnol Biofuels* 6:120–120. <https://doi.org/10.1186/1754-6834-6-120>
- Deparis Q, Claes A, Foulquié-Moreno MR, Thevelein JM (2017) Engineering tolerance to industrially relevant stress factors in yeast cell factories. *FEMS Yeast Research* 17(4). <https://doi.org/10.1093/femsyr/fox036>
- Ding MZ, Wang X, Yang Y, Yuan YJ (2011) Metabolomic study of interactive effects of phenol, furfural, and acetic acid on *Saccharomyces cerevisiae*. *OMICS* 15(10):647–653. <https://doi.org/10.1089/omi.2011.0003>
- Dominguez E, Romani A, Domingues L, Garrote G (2017) Evaluation of strategies for second generation bioethanol production from fast growing biomass *Paulownia* within a biorefinery scheme. *Appl Energy* 187:777–789. <https://doi.org/10.1016/j.apenergy.2016.11.114>
- Dong Y, Hu J, Fan L, Chen Q (2017) RNA-Seq-based transcriptomic and metabolomic analysis reveal stress responses and programmed cell death induced by acetic acid in *Saccharomyces cerevisiae*. *Sci Rep* 7:42659. <https://doi.org/10.1038/srep42659>
- Eisenberg T, Knauer H, Schauer A, Büttner S, Ruckenstuhl C, Carmona-Gutierrez D, Ring J, Schroeder S, Magnes C, Antonacci L, Fussi H, Deszcz L, Hartl R, Schraml E, Criollo A, Megalou E, Weiskopf D, Laun P, Heeren G, Breitenbach M, Grubeck-Loebenstein B, Herker E, Fahrenkrog B, Fröhlich K-U, Sinner F, Tavernarakis N, Minois N, Kroemer G, Madeo F (2009) Induction of autophagy by spermidine promotes longevity. *Nat Cell Biol* 11:1305–1314. <https://doi.org/10.1038/ncb1975>
- Fernandes AR, Mira NP, Vargas RC, Canelhas I, Sá-Correia I (2005) *Saccharomyces cerevisiae* adaptation to weak acids involves the transcription factor Haa1p and Haa1p-regulated genes. *Biochem Biophys Res Commun* 337(1):95–103. <https://doi.org/10.1016/j.bbrc.2005.09.010>
- Foretek D, Wu J, Hopper AK, Boguta M (2016) Control of *Saccharomyces cerevisiae* pre-tRNA processing by environmental conditions. *RNA* 22:339–349. <https://doi.org/10.1261/ma.054973.115>
- Frohner IE, Gregori C, Anrather D, Roitinger E, Schüller C, Ammerer G, Kuchler K (2010) Weak organic acid stress triggers hyperphosphorylation of the yeast zinc-finger transcription factor War1 and dampens stress adaptation. *OMICS* 14(5):575–586. <https://doi.org/10.1089/omi.2010.0032>
- Gao L, Liu Y, Sun H, Li C, Zhao Z, Liu G (2016) Advances in mechanisms and modifications for rendering yeast thermotolerance. *J Biosci Bioeng* 121(6):599–606. <https://doi.org/10.1016/j.jbiosc.2015.11.002>
- Gasch AP, Spellman PT, Kao CM, Carmel-Harel O, Eisen MB, Storz G, Botstein D, Brown PO (2000) Genomic expression programs in the response of yeast cells to environmental changes. *Mol Biol Cell* 11(12):4241–4257. <https://doi.org/10.1091/mbc.11.12.4241>
- Giannattasio S, Guaragnella N, Ždravlević M, Marra E (2013) Molecular mechanisms of *Saccharomyces cerevisiae* stress adaptation and programmed cell death in response to acetic acid. *Front Microbiol* 4:33. <https://doi.org/10.3389/fmicb.2013.00033>
- Godinho CP, Prata CS, Pinto SN, Cardoso C, Bandarra NM, Fernandes F, Sá-Correia I (2018) *Pdr18* is involved in yeast response to acetic acid stress counteracting the decrease of plasma membrane ergosterol content and order. *Sci Rep* 8(1):7860. <https://doi.org/10.1038/s41598-018-26128-7>
- Gomes DG, Guimarães PMR, Pereira FB, Teixeira JA, Domingues L (2012) Plasmid-mediate transfer of FLO-1 into industrial *Saccharomyces cerevisiae* PE-2 strain creates a strain useful for repeat-batch fermentations involving flocculation-sedimentation. *Bioresour Technol* 108:162–168. <https://doi.org/10.1016/j.biortech.2011.12.089>
- Gopinarayanan VE, Nair NU (2018) A semi-synthetic regulon enables rapid growth of yeast on xylose. *Nat Commun* 9:1233. <https://doi.org/10.1038/s41467-018-03645-7>
- Gorsich SW, Dien BS, Nichols NN, Slininger PJ, Liu ZL, Skory CD (2006) Tolerance to furfural-induced stress is associated with pentose phosphate pathway genes *ZWF1*, *GND1*, *RPE1*, and *TKL1* in *Saccharomyces cerevisiae*. *Appl Microbiol Biotechnol* 71(3):339–349. <https://doi.org/10.1007/s00253-005-0142-3>
- Gregori C, Schuller C, Frohner IE, Ammerer G, Kuchler K (2008) Weak organic acids trigger conformational changes of the yeast transcription factor War1 in vivo to elicit stress adaptation. *J Biol Chem* 283(37):25752–25764. <https://doi.org/10.1074/jbc.M803095200>
- Guerreiro JF, Muir A, Ramachandran S, Thorne J, Sa-Correia I (2016) Sphingolipid biosynthesis upregulation by TOR complex 2-Ypk1 signaling during yeast adaptive response to acetic acid stress. *Biochem J* 473(23):4311–4325. <https://doi.org/10.1042/BCJ20160565>
- Guo Z, Olsson L (2014) Physiological response of *Saccharomyces cerevisiae* to weak acids present in lignocellulosic hydrolysate. *FEMS Yeast Res* 14(8):1234–1248. <https://doi.org/10.1111/1567-1364.12221>
- Hasan R, Leroy C, Isnard A-D, Labarre J, Boy-Marcotte E, Toledano MB (2002) The control of the yeast H₂O₂ response by the Msn2/4 transcription factors. *Mol Microbiol* 45(1):233–241. <https://doi.org/10.1046/j.1365-2958.2002.03011.x>
- Hasunuma T, Hori Y, Sakamoto T, Ochiai M, Hatanaka H, Kondo A (2014) Development of a GIN11/FRT-based multiple-gene integration technique affording inhibitor-tolerant, hemicellulolytic, xylose-utilizing abilities to industrial *Saccharomyces cerevisiae* strains for ethanol production from undetoxified lignocellulosic hemicelluloses. *Microb Cell Factories* 13:145. <https://doi.org/10.1186/s12934-014-0145-9>
- Hawkins GM, Doran-Peterson J (2011) A strain of *Saccharomyces cerevisiae* evolved for fermentation of lignocellulosic biomass displays improved growth and fermentative ability in high solids concentrations and in the presence of inhibitory compounds. *Biotechnol Biofuels* 4:49. <https://doi.org/10.1186/1754-6834-4-49>
- Heer D, Heine D, Sauer U (2009) Resistance of *Saccharomyces cerevisiae* to high concentrations of furfural is based on NADPH-dependent reduction by at least two oxidoreductases. *Appl Environ Microbiol* 75(24):7631–7638. <https://doi.org/10.1128/aem.01649-09>
- Hélène D, Nolwenn D, Moïse P, Monique BF (2000) A large-scale study of Yap1p-dependent genes in normal aerobic and H₂O₂-stress conditions: the role of Yap1p in cell proliferation control in yeast. *Mol Microbiol* 36(4):830–845. <https://doi.org/10.1046/j.1365-2958.2000.01845.x>

- Henriques SF, Mira NP, Sá-Correia I (2017) Genome-wide search for candidate genes for yeast robustness improvement against formic acid reveals novel susceptibility (Trk1 and positive regulators) and resistance (Haa1-regulon) determinants. *Biotechnol Biofuels* 10:96. <https://doi.org/10.1186/s13068-017-0781-5>
- Herrero E, Ros J, Bellí G, Cabisco E (2008) Redox control and oxidative stress in yeast cells. *Biochim Biophys Acta* 1780(11):1217–1235. <https://doi.org/10.1016/j.bbagen.2007.12.004>
- Higgins VJ, Beckhouse AG, Oliver AD, Rogers PJ, Dawes IW (2003) Yeast genome-wide expression analysis identifies a strong ergosterol and oxidative stress response during the initial stages of an industrial lager fermentation. *Appl Environ Microbiol* 69(8):4777–4787. <https://doi.org/10.1128/aem.69.8.4777-4787.2003>
- Iwaki A, Kawai T, Yamamoto Y, Izawa S (2013a) Biomass conversion inhibitors furfural and 5-hydroxymethylfurfural induce formation of messenger RNP granules and attenuate translation activity in *Saccharomyces cerevisiae*. *Appl Environ Microbiol* 79(5):1661–1667. <https://doi.org/10.1128/AEM.02797-12>
- Iwaki A, Ohnuki S, Suga Y, Izawa S, Ohya Y (2013b) Vanillin inhibits translation and induces messenger ribonucleoprotein (mRNP) granule formation in *Saccharomyces cerevisiae*: application and validation of high-content, image-based profiling. *PLoS One* 8(4):e61748. <https://doi.org/10.1371/journal.pone.0061748>
- Jain NK, Roy I (2009) Effect of trehalose on protein structure. *Protein Sci* 18(1):24–36. <https://doi.org/10.1002/pro.3>
- Jayakody LN, Turner TL, Yun EJ, Kong II, Liu JJ, Jin YS (2018) Expression of Gre2p improves tolerance of engineered xylose-fermenting *Saccharomyces cerevisiae* to glycolaldehyde under xylose metabolism. *Appl Microbiol Biotechnol* 102(18):8121–8133. <https://doi.org/10.1007/s00253-018-9216-x>
- Jesus MS, Romani A, Genisheva Z, Teixeira JA, Domingues L (2017) Integral valorization of vine pruning residue by sequential autohydrolysis stages. *J Clea Prod* 168:74–86. <https://doi.org/10.1016/j.jclepro.2017.08.230>
- Jin M, Sarks C, Gunawan C, Bice BD, Simonett SP, Narasimhan RA, Willis LB, Dale BE, Venkatesh B, Sato TK (2013) Phenotypic selection of a wild *Saccharomyces cerevisiae* strain for simultaneous saccharification and co-fermentation of AFEX™ pretreated corn stover. *Biotechnol Biofuels* 6:108. <https://doi.org/10.1186/1754-6834-6-108>
- Keating JD, Panganiban C, Mansfield SD (2006) Tolerance and adaptation of ethanologenic yeasts to lignocellulosic inhibitory compounds. *Biotechnol Bioeng* 93(6):1196–1206. <https://doi.org/10.1002/bit.20838>
- Kelbert M, Romani A, Coelho E, Pereira FB, Teixeira JA, Domingues L (2015) Lignocellulosic bioethanol production with revalorization of low-cost agroindustrial by-products as nutritional supplements. *Ind Crop Prod* 64:16–24. <https://doi.org/10.1016/j.indcrop.2014.10.056>
- Kelbert M, Romani A, Coelho E, Pereira FB, Teixeira JA, Domingues L (2016) Simultaneous saccharification and fermentation of hydrothermal pretreated lignocellulosic biomass: evaluation of process performance under multiple stress conditions. *Bioen Res* 9:750–762. <https://doi.org/10.1007/s12155-016-9722-6>
- Kim D, Hahn J-S (2013) Roles of the Yap1 transcription factor and antioxidants in *Saccharomyces cerevisiae*'s tolerance to furfural and 5-hydroxymethylfurfural, which function as thiol-reactive electrophiles generating oxidative stress. *Appl Environ Microbiol* 79(16):5069–5077. <https://doi.org/10.1128/aem.00643-13>
- Kim SR, Xu H, Lesmana A, Kuzmanovic U, Au M, Florencia C, Oh EJ, Zhang G, Kim KH, Jin Y-S (2015) Deletion of *PHO13*, encoding haloacid dehalogenase type iia phosphatase, results in upregulation of the pentose phosphate pathway in *Saccharomyces cerevisiae*. *Appl Environ Microbiol* 81(5):1601–1609. <https://doi.org/10.1128/aem.03474-14>
- Kim S-K, Jo J-H, Jin Y-S, Seo J-H (2017a) Enhanced ethanol fermentation by engineered *Saccharomyces cerevisiae* strains with high spermidine contents. *Bioprocess Biosyst Eng* 40(5):683–691. <https://doi.org/10.1007/s00449-016-1733-3>
- Kim SR, Skerker JM, Kong II, Kim H, Maurer MJ, Zhang GC, Peng D, Wei N, Arkin AP, Jin YS (2017b) Metabolic engineering of a haploid strain derived from a triploid industrial yeast for producing cellulosic ethanol. *Metab Eng* 40:176–185. <https://doi.org/10.1016/j.ymben.2017.02.006>
- Klinke HB, Olsson L, Thomsen AB, Ahring BK (2003) Potential inhibitors from wet oxidation of wheat straw and their effect on ethanol production of *Saccharomyces cerevisiae*: wet oxidation and fermentation by yeast. *Biotechnol Bioeng* 81(6):738–747. <https://doi.org/10.1002/bit.10523>
- Ko JK, Um Y, Park YC, Seo JH, Kim KH (2015) Compounds inhibiting the bioconversion of hydrothermally pretreated lignocellulose. *Appl Microbiol Biotechnol* 99(10):4201–4212. <https://doi.org/10.1007/s00253-015-6595-0>
- Kren A, Mamnun YM, Bauer BE, Schüller C, Wolfger H, Hatzixanthos K, Mollapour M, Gregori C, Piper P, Kuchler K (2003) War1p, a novel transcription factor controlling weak acid stress response in yeast. *Mol Cell Biol* 23(5):1775–1785. <https://doi.org/10.1128/mcb.23.5.1775-1785.2003>
- Kuang Z, Pingley S, Ji H, Boeke JD (2017) Msn2/4 regulate expression of glycolytic enzymes and control transition from quiescence to growth. *eLife* 6:e29938. <https://doi.org/10.7554/eLife.29938>
- Lane S, Xu H, Oh EJ, Kim H, Lesmana A, Jeong D, Guochang Z, Tsai C-S, Jin Y-S, Kim SR (2018) Glucose repression can be alleviated by reducing glucose phosphorylation rate in *Saccharomyces cerevisiae*. *Sci Rep* 8:2613. <https://doi.org/10.1038/s41598-018-20804-4>
- Larochelle M, Drouin S, Robert F, Turcotte B (2006) Oxidative stress-activated zinc cluster protein Stb5 has dual activator/repressor functions required for pentose phosphate pathway regulation and NADPH production. *Mol Cell Biol* 26(17):6690–6701. <https://doi.org/10.1128/mcb.02450-05>
- Larsson S, Quintana-Sainz A, Reimann A, Nilvebrant NO, Jonsson LJ (2000) Influence of lignocellulose-derived aromatic compounds on oxygen-limited growth and ethanolic fermentation by *Saccharomyces cerevisiae*. *Appl Biochem Biotechnol* 84-86:617–632
- Larsson S, Cassland P, Jönsson LJ (2001) Development of a *Saccharomyces cerevisiae* strain with enhanced resistance to phenolic fermentation inhibitors in lignocellulose hydrolysates by heterologous expression of laccase. *Appl Environ Microbiol* 67(3):1163–1170. <https://doi.org/10.1128/aem.67.3.1163-1170.2001>
- Lavoie JM, Capek-Menard E, Gauvin H, Chomet E (2010) Production of pulp from *Salix viminalis* energy crops using the FIRSST process. *Bioresour Technol* 101:4940–4946. <https://doi.org/10.1016/j.biortech.2009.09.021>
- Levin DE (2005) Cell wall integrity signaling in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev* 69(2):262–291. <https://doi.org/10.1128/MMBR.69.2.262-291.2005>
- Levin DE (2011) Regulation of cell wall biogenesis in *Saccharomyces cerevisiae*: the cell wall integrity signaling pathway. *Genetics* 189(4):1145–1175. <https://doi.org/10.1534/genetics.111.128264>
- Lin FM, Qiao B, Yuan YJ (2009) Comparative proteomic analysis of tolerance and adaptation of ethanologenic *Saccharomyces cerevisiae* to furfural, a lignocellulosic inhibitory compound. *Appl Environ Microbiol* 75(11):3765–3776. <https://doi.org/10.1128/AEM.02594-08>
- Lindberg L, Santos AXS, Riezman H, Olsson L, Bettiga M (2013) Lipidomic profiling of *Saccharomyces cerevisiae* and *Zygosaccharomyces bailii* reveals critical changes in lipid composition in response to acetic acid stress. *PLoS One* 8(9):e73936. <https://doi.org/10.1371/journal.pone.0073936>
- Liu ZL, Moon J (2009) A novel NADPH-dependent aldehyde reductase gene from *Saccharomyces cerevisiae* NRRL Y-12632 involved in the detoxification of aldehyde inhibitors derived from

- lignocellulosic biomass conversion. *Gene* 446(1):1–10. <https://doi.org/10.1016/j.gene.2009.06.018>
- Liu ZL, Slininger PJ, Dien BS, Berhow MA, Kurtzman CP, Gorsich SW (2004) Adaptive response of yeasts to furfural and 5-hydroxymethylfurfural and new chemical evidence for HMF conversion to 2,5-bis-hydroxymethylfuran. *J Ind Microbiol Biotechnol* 31(8):345–352. <https://doi.org/10.1007/s10295-004-0148-3>
- Liu ZL, Slininger PJ, Gorsich SW (2005) Enhanced biotransformation of furfural and hydroxymethylfurfural by newly developed ethanologenic yeast strains. *Appl Biochem Biotechnol* 121:451–460. <https://doi.org/10.1385/ABAB:121:1-3:0451>
- Liu ZL, Moon J, Andersh BJ, Slininger PJ, Weber S (2008) Multiple gene-mediated NAD(P)H-dependent aldehyde reduction is a mechanism of in situ detoxification of furfural and 5-hydroxymethylfurfural by *Saccharomyces cerevisiae*. *Appl Microbiol Biotechnol* 81(4):743–753. <https://doi.org/10.1007/s00253-008-1702-0>
- Liu ZL, Wang X, Weber SA (2018) Tolerant industrial yeast *Saccharomyces cerevisiae* posses a more robust cell wall integrity signaling pathway against 2-furaldehyde and 5-(hydroxymethyl)-2-furaldehyde. *J Biotechnol* 276-277:15–24. <https://doi.org/10.1016/j.jbiotec.2018.04.002>
- Ma M, Liu ZL (2010) Comparative transcriptome profiling analyses during the lag phase uncover *YAP1*, *PDR1*, *PDR3*, *RPN4*, and *HSF1* as key regulatory genes in genomic adaptation to the lignocellulose derived inhibitor HMF for *Saccharomyces cerevisiae*. *BMC Genomics* 11:660–660. <https://doi.org/10.1186/1471-2164-11-660>
- Maeta K, Izawa S, Okazaki S, Kuge S, Inoue Y (2004) Activity of the Yap1 transcription factor in *Saccharomyces cerevisiae* is modulated by methylglyoxal, a metabolite derived from glycolysis. *Mol Cell Biol* 24(19):8753–8764. <https://doi.org/10.1128/mcb.24.19.8753-8764.2004>
- Menonides FIC, Brul S, Hellingwerf KJ, Bakker BM, Teixeira de Mattos MJ (2014) A kinetic model of catabolic adaptation and protein reprofiling in *Saccharomyces cerevisiae* during temperature shifts. *FEBS J* 281(3):825–841. <https://doi.org/10.1111/febs.12649>
- Mira NP, Becker JD, Sa-Correia I (2010) Genomic expression program involving the Haa1p-regulon in *Saccharomyces cerevisiae* response to acetic acid. *OMICS* 14(5):587–601. <https://doi.org/10.1089/omi.2010.0048>
- Modenbach AA, Nokes SE (2012) The use of high-solids loadings in biomass pretreatment—a review. *Biotechnol Bioeng* 109(6):1430–1442. <https://doi.org/10.1002/bit.24464>
- Morano KA, Grant CM, Moye-Rowley WS (2012) The response to heat shock and oxidative stress in *Saccharomyces cerevisiae*. *Genetics* 190:1157–1195. <https://doi.org/10.1534/genetics.111.128033>
- Moyés DN, Reis VC, de Almeida JR, de Moraes LM, Torres FA (2016) Xylose fermentation by *Saccharomyces cerevisiae*: challenges and prospects. *Int J Mol Sci* 17:207. <https://doi.org/10.3390/ijms17030207>
- Nguyen TT, Kitajima S, Izawa S (2014a) Importance of glucose-6-phosphate dehydrogenase (G6PDH) for vanillin tolerance in *Saccharomyces cerevisiae*. *J Biosci Bioeng* 118(3):263–269. <https://doi.org/10.1016/j.jbiosc.2014.02.025>
- Nguyen TTM, Iwaki A, Ohya Y, Izawa S (2014b) Vanillin causes the activation of Yap1 and mitochondrial fragmentation in *Saccharomyces cerevisiae*. *J Biosci Bioeng* 117(1):33–38. <https://doi.org/10.1016/j.jbiosc.2013.06.008>
- Nishida N, Jing D, Kuroda K, Ueda M (2014) Activation of signaling pathways related to cell wall integrity and multidrug resistance by organic solvent in *Saccharomyces cerevisiae*. *Curr Genet* 60(3):149–162. <https://doi.org/10.1007/s00294-013-0419-5>
- Nygård Y, Mojzita D, Toivari M, Penttilä M, Wiebe MG, Ruohonen L (2014) The diverse role of Pdr12 in resistance to weak organic acids. *Yeast* 31(6):219–232. <https://doi.org/10.1002/yea.3011>
- Osiro KO, Brink DP, Borgström C, Wasserstrom L, Carlquist M, Gorwa-Grauslund MF (2018) Assessing the effect of d-xylose on the sugar signaling pathways of *Saccharomyces cerevisiae* in strains engineered for xylose transport and assimilation. *FEMS Yeast Res* 18(1). <https://doi.org/10.1093/femsyr/fox096>
- Ouyang X, Tran QT, Goodwin S, Wible RS, Sutter CH, Sutter TR (2011) Yap1 activation by H₂O₂ or thiol-reactive chemicals elicits distinct adaptive gene responses. *Free Radic Biol Med* 50(1):1–13. <https://doi.org/10.1016/j.freeradbiomed.2010.10.697>
- Palma M, Guerreiro JF, Sa-Correia I (2018) Adaptive response and tolerance to acetic acid in *Saccharomyces cerevisiae* and *Zygosaccharomyces bailii*: a physiological genomics perspective. *Front Microbiol* 9:274. <https://doi.org/10.3389/fmicb.2018.00274>
- Palmqvist E, Grage H, Meinander NQ, Hahn-Hägerdal B (1999) Main and interaction effects of acetic acid, furfural, and p-hydroxybenzoic acid on growth and ethanol productivity of yeasts. *Biotechnol Bioeng* 63(1):46–55. [https://doi.org/10.1002/\(SICI\)10970290\(19990405\)63:1<46::AID-BIT5>3.0.CO;2-J](https://doi.org/10.1002/(SICI)10970290(19990405)63:1<46::AID-BIT5>3.0.CO;2-J)
- Pampulha ME, Loureiro-Dias MC (1990) Activity of glycolytic enzymes of *Saccharomyces cerevisiae* in the presence of acetic acid. *Appl Microbiol Biotechnol* 34(3):375–380. <https://doi.org/10.1007/bf00170063>
- Pearce AK, Booth IR, Brown AJ (2001) Genetic manipulation of 6-phosphofructo-1-kinase and fructose 2,6-bisphosphate levels affects the extent to which benzoic acid inhibits the growth of *Saccharomyces cerevisiae*. *Microbiol* 147:403–410. <https://doi.org/10.1099/00221287-147-2-403>
- Pereira C, Chaves S, Alves S, Salin B, Camougrand N, Manon S, Sousa MJ, Corte-Real M (2010a) Mitochondrial degradation in acetic acid-induced yeast apoptosis: the role of Pep4 and the ADP/ATP carrier. *Mol Microbiol* 76(6):1398–1410. <https://doi.org/10.1111/j.1365-2958.2010.07122.x>
- Pereira FB, Guimarães PMR, Teixeira JA, Domingues L (2010b) Optimization of low-cost medium for very high gravity ethanol fermentations by *Saccharomyces cerevisiae* using statistical experimental designs. *Bioresour Technol* 101:7856–7863. <https://doi.org/10.1016/j.biortech.2010.04.082>
- Pereira FB, Guimarães PMR, Teixeira JA, Domingues L (2010c) Selection of *Saccharomyces cerevisiae* strains for efficient very high gravity bio-ethanol fermentation processes. *Biotechnol Lett* 32:1655–1661. <https://doi.org/10.1007/s10259-010-0330-9>
- Pereira FB, Guimarães PMR, Gomes DG, Mira NP, Teixeira MC, Sá-Correia I, Domingues L (2011a) Identification of candidate genes for yeast engineering to improve bioethanol production in very-high-gravity and lignocellulosic biomass industrial fermentations. *Biotechnol Biofuels* 4:57. <https://doi.org/10.1186/1754-6834-4-57>
- Pereira FB, Guimarães PMR, Teixeira JA, Domingues L (2011b) Robust industrial *Saccharomyces cerevisiae* strains for very high gravity bio-ethanol fermentations. *J Biosci Bioeng* 112:130–136. <https://doi.org/10.1016/j.jbiosc.2011.03.022>
- Pereira FB, Romani A, Ruiz HA, Teixeira JA, Domingues L (2014a) Industrial robust yeast isolates with great potential for fermentation of lignocellulosic biomass. *Bioresour Technol* 161:192–199. <https://doi.org/10.1016/j.biortech.2014.03.043>
- Pereira FB, Teixeira MC, Mira NP, Sá-Correia I, Domingues L (2014b) Genome-wide screening of *Saccharomyces cerevisiae* genes required to foster tolerance towards industrial wheat straw hydrolysates. *J Ind Microbiol Biotechnol* 41:1753–1761. <https://doi.org/10.1007/s10295-014-1519-z>
- Petitjean M, Teste M-A, François JM, Parrou J-L (2015) Yeast tolerance to various stresses relies on the trehalose-6p synthase (Tps1) protein, not on trehalose. *J Biol Chem* 290(26):16177–16190. <https://doi.org/10.1074/jbc.M115.653899>
- Purwadi R, Brandberg T, Taherzadeh MJ (2007) A possible industrial solution to ferment lignocellulosic hydrolyzate to ethanol: continuous cultivation with flocculating yeast. *Int J Mol Sci* 8:920–932. <https://doi.org/10.3390/i8090920>

- Roelants FM, Breslow DK, Muir A, Weissman JS, Thomer J (2011) Protein kinase Ypk1 phosphorylates regulatory proteins Orm1 and Orm2 to control sphingolipid homeostasis in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A* 108(48):19222–19227. <https://doi.org/10.1073/pnas.1116948108>
- Romani A, Ruiz HA, Pereira FB, Teixeira JA, Domingues L (2014) Integrated approach for effective bioethanol production using whole slurry from autohydrolyzed *Eucalyptus globulus* wood at high-solid loadings. *Fuel* 135:482–491. <https://doi.org/10.1016/j.fuel.2014.06.061>
- Romani A, Pereira F, Johansson B, Domingues L (2015) Metabolic engineering of *Saccharomyces cerevisiae* ethanol strains PE-2 and CAT-1 for efficient lignocellulosic fermentation. *Bioresour Technol* 179: 150–158. <https://doi.org/10.1016/j.biortech.2014.12.020>
- Salusjärvi L, Pitkänen JP, Aristidou A, Ruohonen L, Penttilä M (2006) Transcription analysis of recombinant *Saccharomyces cerevisiae* reveals novel responses to xylose. *Appl Biochem Biotechnol* 128: 237–261. <https://doi.org/10.1385/ABAB:128:3:237>
- Serrano R (1984) Plasma membrane ATPase of fungi and plants as a novel type of proton pump. *Curr Top Cell Reg* 23:87–126. <https://doi.org/10.1016/B978-0-12-152823-2.50007-6>
- Simões T, Mira NP, Fernandes AR, Sa-Correia I (2006) The *SPH1* gene, encoding a glycosylphosphatidylinositol-anchored cell wall protein, plays a prominent role in the development of yeast resistance to lipophilic weak-acid food preservatives. *Appl Environ Microbiol* 72(11):7168–7175. <https://doi.org/10.1128/AEM.01476-06>
- Subtil T, Boles E (2012) Competition between pentoses and glucose during uptake and catabolism in recombinant *Saccharomyces cerevisiae*. *Biotechnol Biofuels* 5:14. <https://doi.org/10.1186/1754-6834-5-14>
- Sun Y, Miao Y, Yamane Y, Zhang C, Shokat KM, Takematsu H, Kozutsumi Y, Drubin DG (2012) Orm protein phosphoregulation mediates transient sphingolipid biosynthesis response to heat stress via the Pkh-Ypk and Cdc55-PP2A pathways. *Mol Biol Cell* 23(12): 2388–2398. <https://doi.org/10.1091/mbc.E12-03-0209>
- Sundström L, Larsson S, Jönsson LJ (2010) Identification of *Saccharomyces cerevisiae* genes involved in the resistance to phenolic fermentation inhibitors. *Appl Biochem Biotechnol* 161(1): 106–115. <https://doi.org/10.1007/s12010-009-8811-9>
- Ullah A, Orij R, Brul S, Smits GJ (2012) Quantitative analysis of the modes of growth inhibition by weak organic acids in *Saccharomyces cerevisiae*. *Appl Environ Microbiol* 78(23):8377–8387. <https://doi.org/10.1128/AEM.02126-12>
- Unrean P, Gätgens J, Klein B, Noack S, Champreda V (2018) Elucidating cellular mechanisms of *Saccharomyces cerevisiae* tolerant to combined lignocellulosic-derived inhibitors using high-throughput phenotyping and multiomics analyses. *FEMS Yeast Res* 18(8). <https://doi.org/10.1093/femsyr/foy106>
- van den Hazel HB, Pichler H, do Valle Matta MA, Leitner E, Goffeau A, Daum G (1999) *PDR16* and *PDR17*, two homologous genes of *Saccharomyces cerevisiae*, affect lipid biosynthesis and resistance to multiple drugs. *J Biol Chem* 274(4):1934–1941
- Varga E, Klinke HB, Réczey K, Thomsen AB (2004) High solid simultaneous saccharification and fermentation of wet oxidized corn stover to ethanol. *Biotechnol Bioeng* 88:567–574. <https://doi.org/10.1002/bit.20222>
- Verghese J, Abrams J, Wang Y, Morano KA (2012) Biology of the heat shock response and protein chaperones: budding yeast (*Saccharomyces cerevisiae*) as a model system. *Microbiol Mol Biol Rev* 76:115–158. <https://doi.org/10.1128/MMBR.05018-11>
- Wallace-Salinas V, Signori L, Li Y-Y, Ask M, Bettiga M, Porro D, Thevelein JM, Branduardi P, Foulquié-Moreno MR, Gorwa-Grauslund M (2014) Re-assessment of *YAP1* and *MCR1* contributions to inhibitor tolerance in robust engineered *Saccharomyces cerevisiae* fermenting undetoxified lignocellulosic hydrolysate. *AMB Express* 4:56–56. <https://doi.org/10.1186/s13568-014-0056-5>
- Wang X, Liang Z, Hou J, Bao X, Shen Y (2016) Identification and functional evaluation of the reductases and dehydrogenases from *Saccharomyces cerevisiae* involved in vanillin resistance. *BMC Biotechnol* 16:31. <https://doi.org/10.1186/s12896-016-0264-y>
- Wei S, Liu Y, Wu M, Ma T, Bai X, Hou J, Shen Y, Bao X (2018) Disruption of the transcription factors Thi2p and Nrm1p alleviates the post-glucose effect on xylose utilization in *Saccharomyces cerevisiae*. *Biotechnol Biofuels* 11:112. <https://doi.org/10.1186/s13068-018-1112-1>
- Westman JO, Taherzadeh MJ, Franzén CJ (2012) Inhibitor tolerance and flocculation of a yeast strain suitable for second generation bioethanol production. *Electron J Biotechnol* 15(3). <https://doi.org/10.2225/vol15-issue3-fulltext-8>
- Westman JO, Mapelli V, Taherzadeh MJ, Franzén CJ (2014) Flocculation causes inhibitor tolerance in *Saccharomyces cerevisiae* for second-generation bioethanol production. *Appl Environ Microbiol* 80: 6908–6918. <https://doi.org/10.1128/AEM.01906-14>
- Wimalasena TT, Greetham D, Marvin ME, Liti G, Chandelia Y, Hart A, Louis EJ, Phister TG, Tucker GA, Smart KA (2014) Phenotypic characterisation of *Saccharomyces* spp. yeast for tolerance to stresses encountered during fermentation of lignocellulosic residues to produce bioethanol. *Microb Cell Factories* 13:47. <https://doi.org/10.1186/1475-2859-13-47>
- Woo JM, Yang KM, Kim SU, Blank LM, Park JB (2014) High temperature stimulates acetic acid accumulation and enhances the growth inhibition and ethanol production by *Saccharomyces cerevisiae* under fermenting conditions. *Appl Microbiol Biotechnol* 98:6085–6094. <https://doi.org/10.1007/s00253-014-5691-x>
- Wu G, Xu Z, Jonsson LJ (2017) Profiling of *Saccharomyces cerevisiae* transcription factors for engineering the resistance of yeast to lignocellulose-derived inhibitors in biomass conversion. *Microb Cell Factories* 16(1):199. <https://doi.org/10.1186/s12934-017-0811-9>
- Xianxian Z, Juan T, Xu W, Ruoheng Y, Xiaoping Z, Yunfu G, Xi L, Menggen M (2015) YNL134C from *Saccharomyces cerevisiae* encodes a novel protein with aldehyde reductase activity for detoxification of furfural derived from lignocellulosic biomass. *Yeast* 32(5): 409–422. <https://doi.org/10.1002/yea.3068>
- Yáñez R, Romani A, Garrote G, Alonso JL, Parajó JC (2009) Experimental evaluation of alkaline treatment as a method for enhancing the enzymatic digestibility of autohydrolysed Acacia dealbata. *J Chem Technol Biotechnol* 84(7):1070–1077. <https://doi.org/10.1002/jctb.2136>
- Yu N, Tan L, Sun Z-Y, Tang Y-Q, Kida K (2018) Production of bio-ethanol by integrating microwave-assisted dilute sulfuric acid pretreated sugarcane bagasse slurry with molasses. *Appl Biochem Biotechnol* 185: 191–206. <https://doi.org/10.1007/s12010-017-2651-9>
- Zhang G-C, Liu J-J, Ding W-T (2012) Decreased xylitol formation during xylose fermentation in *Saccharomyces cerevisiae* due to overexpression of water-forming NADH oxidase. *Appl Environ Microbiol* 78(4):1081–1086. <https://doi.org/10.1128/aem.06635-11>
- Zhang G-C, Kong II, Wei N, Peng D, Turner TL, Sung BH, Sohn J, Jin Y (2016) Optimization of an acetate reduction pathway for producing cellulosic ethanol by engineered yeast. *Biotechnol Bioeng* 113: 2587–2596. <https://doi.org/10.1002/bit.26021>
- Zhou H, Zhu JY, Luo X, Leu S-Y, Wu X, Gleisner R, Dien BS, Hector RE, Yang D, Qiu X, Horn E, Negron J (2013) Bioconversion of beetle-killed lodgepole pine using spori: process scale-up design, lignin coproduct, and high solids fermentation without detoxification. *Ind Eng Chem Res* 52:16057–16065. <https://doi.org/10.1021/ie402873y>
- Zhou Q, Liu ZL, Ning K, Wang A, Zeng X, Xu J (2014) Genomic and transcriptome analyses reveal that MAPK- and phosphatidylinositol-signaling pathways mediate tolerance to 5-hydroxymethyl-2-furaldehyde for industrial yeast *Saccharomyces cerevisiae*. *Sci Rep* 4:6556. <https://doi.org/10.1038/srep06556>