

promising method for producing large scale amounts of recombinant BMPs, in pure and bioactive form, for novel biomedical applications.

Acknowledgments: Fundação para a Ciência e Tecnologia for PhD grant SFRH/BD/17049/2004 and project ElastM POCI/CTM/57177/2004 supported by FEDER and the Fundação para a Ciência e Tecnologia; European STREP Project HIPPOCRATES (NMP3-CT-2003-505758). The work was carried out under the scope of the European NoE EXPERTISSUES (NMP3-CT-2004-500283).

(P 168) Expression, Purification and Bioactivity of Recombinant Human Bone Morphogenetic Protein-4, -9, -10, -11 and -14 Produced in Escherichia Coli for Tissue Engineering Applications

P.C. Bessa^{1,2,3}, M.T. Cerqueira^{1,2}, T. Rada^{1,2}, M.E. Gomes^{1,2}, N.M. Neves^{1,2}, M. Casal³, R.L. Reis^{1,2}

¹3B's Research Group—Biomaterials, Biodegradables and Biomimetics, University of Minho, Braga, Portugal.

²IBB—Institute for Biotechnology and Bioengineering, PT Government Associated Laboratory, Braga, Portugal.

³CBMA—Molecular and Environmental Biology Centre/ Biology Department, University of Minho, Braga, Portugal.

Bone morphogenetic proteins (BMPs) are cytokines from the TGF- β superfamily, with important roles during embryonic development and in inducing bone and cartilage in the adult body. In this contribution, we report the expression of recombinant human BMP-4, BMP-9, BMP-10, BMP-11 (or growth differentiation factor-11, GDF-11) and BMP-14 (GDF-5), using *Escherichia coli* pET-25b expression system. The BMPs were purified by affinity chromatography and its bioactivity accessed in C2C12 cell line, by screening the expression of osteogenic markers with RT-PCR. The expression of Smad-1, Smad-3, Smad-5, Runx2/cbfa1, Osterix, Bone sialoprotein, Osteopontin and ALP was increased with some or all of these BMPs, while Smad-7 was down-regulated, coinciding with the different changes in cell morphology. No cytotoxicity was observed and around 200mg of BMPs were purified per Liter of batch. Ongoing work includes bioactivity assays in human adipose stem cells and with the effect of heparin on the BMP activity. The approach described here is a