

# Monotone and Nonmonotone Implementations of an Interior-Point SQP Filter Line Search Algorithm

M. FERNANDA P. COSTA

Departamento de Matemática para a Ciência e Tecnologia

EDITE M. G. P. FERNANDES

Departamento de Produção e Sistemas

Universidade do Minho

Campus de Azurém 4800-058 Guimarães

PORTUGAL

[mfc@mct.uminho.pt](mailto:mfc@mct.uminho.pt) <http://www.mct.uminho.pt/membros/mfc.html>

[emgpf@dps.uminho.pt](mailto:emgpf@dps.uminho.pt) <http://www.norg.uminho.pt/emgpf/>

*Abstract:* - We present a sequential quadratic programming (SQP) filter line search method which relies on an interior-point paradigm for solving the QP subproblems. Besides the usual step size acceptability criteria, a nonmonotone line search approach is also implemented. A comprehensive description of the algorithm, which also includes a feasibility restoration phase, is provided. Both monotone and nonmonotone algorithms have been implemented in Fortran 90, and the numerical results confirm that relaxing the acceptability criteria is effective.

*Key-Words:* - Nonlinear programming, Interior-point SQP method, Filter method, Line search

## 1 Introduction

The proposed algorithm is a filter line search algorithm for solving nonlinear optimization problems of the form

$$\begin{aligned} \min_{x \in R^n} F(x) \\ \text{s.t. } b \leq h(x) \leq b+r, l \leq x \leq u, \end{aligned} \quad (1)$$

where  $h_k : R^n \rightarrow R$  for  $k=1, \dots, m$  and  $F : R^n \rightarrow R$  are nonlinear and twice continuously differentiable functions.  $r$  is the vector of ranges on the constraints  $h(x)$ ,  $u$  and  $l$  are the vectors of upper and lower bounds on the variables  $x$  and  $b$  is assumed to be a finite real vector. Elements of the vector  $r$ ,  $l$  and  $u$  are real numbers subject to the following limitations:

$$0 \leq r_k \leq \infty, \quad -\infty \leq l_i, u_i \leq \infty \quad \text{for } k=1, \dots, m, \quad i=1, \dots, n.$$

Constraints of the form  $b \leq h(x) \leq b+r$  are denoted by range constraints. Note that equality constraints can still be treated as range constraints with  $r=0$ .

Let  $\nabla F(x)$  denote the gradient of  $F(x)$  and  $\nabla \bar{h}(x)$  denote the Jacobian matrix of the constraint vector

$$\bar{h}(x)^T = [h(x) - b, b+r - h(x), x - l, u - x].$$

A solution of (1) will be denoted by  $x^*$ , and we assume that there is a finite number of solutions. We also assume that the first order Kuhn-Tucker (KT) conditions hold (with strict complementarity) at  $x^*$ . Thus, the constraints are verified and there exists a Lagrange multiplier vector  $\lambda^* \geq 0$  such that

$$\nabla F(x^*) = \nabla \bar{h}(x^*)^T \lambda^*, \quad \bar{h}(x^*)^T \lambda^* = 0. \quad (2)$$

Given a starting point  $x_0$ , the proposed line search algorithm generates a sequence of improved estimates  $x_k$  of the solution for the problem (1) using a sequential quadratic programming (SQP) method. At each iteration  $k$  the search direction  $\Delta_k$  is the solution of a quadratic programming subproblem whose objective function approximates the Lagrangian function  $L(x, \lambda) = F(x) - \lambda^T \bar{h}(x)$  and whose constraints are linear approximations to the constraints in (1). The usual definition of the QP subproblem is the following:

$$\min_{\Delta \in R^n} \Delta^T H_k \Delta + \nabla F_k^T \Delta \quad (3)$$

$$\text{s.t. } b \leq \nabla h_k \Delta + h_k \leq b+r, \quad l \leq \Delta + x_k \leq u$$

where  $\nabla h(x)$  denotes the Jacobian matrix of the constraint vector  $h(x)$ , and  $\nabla F_k$ ,  $h_k$  and  $\nabla h_k$  denote the relevant quantities evaluated at  $x_k$ . The matrix  $H_k$  is a symmetric positive definite approximation to the Hessian of the Lagrangian function. This problem has a solution  $\Delta_k$  and a Lagrange multiplier  $\pi_k \geq 0$  that satisfy

$$H_k \Delta_k + \nabla F_k = \nabla \bar{h}_k^T \pi_k, \quad \pi_k^T (\nabla \bar{h}_k \Delta_k + \bar{h}_k) = 0.$$

Clearly the most common approach for solving (3) considers active set methods (see, for example, [9]). Solving QP subproblems with equality constraints is straightforward. However, problems that have inequality constraints are significantly more difficult to solve than problems in which all constraints are

equations since it is not known in advance which inequality constraints are active at the solution.

In this paper, we describe a new SQP method that is based on the interior-point paradigm for solving the QP subproblems (3). To promote the global convergence, the filter technique of Fletcher and Leyffer [3] is used to globalize the SQP algorithm, avoiding the use of a merit function and the updating of the penalty parameter. The underlying concept is that trial iterates are accepted if they improve the objective function or improve the constraints violation, instead of a combination of those two measures defined by a merit function. In this context, we also test a specific implementation of a nonmonotone line search approach.

The paper is organized as follows. Section 2 describes the interior-point method used to solve the QP subproblems. The filter mechanism is presented in Section 3, Section 4 describes the nonmonotone strategy and Section 5 contains the numerical results, some conclusions and future work.

## 2 The Interior-Point Framework

This section describes an infeasible primal-dual interior-point method for solving the quadratic subproblem (3). We refer to [11] for details. Adding nonnegative slack variables  $w, p, g, t$ , (3) becomes

$$\min \Delta^T H_k \Delta + \nabla F_k^T \Delta \quad (4)$$

$$\text{s.t. } \nabla h_k \Delta - w = b - h_k, \quad \nabla h_k \Delta + p = b + r - h_k,$$

$$\Delta - g = l - x_k, \quad \Delta + t = u - x_k, \quad w, p, g, t \geq 0.$$

The nonnegativity constraints are then eliminated by incorporating them in logarithmic barrier terms in the objective function transforming (4) into

$$\begin{aligned} \min \Delta^T H_k \Delta + \nabla F_k^T \Delta - \mu \sum_{j=1}^m \ln(w_j) - \mu \sum_{j=1}^m \ln(p_j) - \\ - \mu \sum_{i=1}^n \ln(g_i) - \mu \sum_{i=1}^n \ln(t_i) \end{aligned}$$

subject to the same set of equality constraints, where  $\mu$  is a positive barrier parameter. Optimality conditions for this subproblem produce the standard primal-dual system

$$\begin{aligned} H_k \Delta + \nabla F_k^T \Delta - \nabla h_k^T y - z + s = 0, \quad y + q - v = 0, \\ WVe_1 = \mu e_1, PQe_1 = \mu e_1, GZe_2 = \mu e_2, TSe_2 = \mu e_2, \\ \nabla h_k \Delta + h_k - b - w = 0, r - w - p = 0, \\ \Delta + x_k - l - g = 0, u - \Delta - x_k - t = 0, \end{aligned} \quad (5)$$

where  $V = \text{diag}(v_j)$ ,  $Q = \text{diag}(q_j)$ ,  $Z = \text{diag}(z_i)$ ,

$$S = \text{diag}(s_i), W = \text{diag}(w_j), P = \text{diag}(p_j),$$

$G = \text{diag}(g_i)$  and  $T = \text{diag}(t_i)$  are diagonal matrices,

$y = v - q$ ,  $e_1 = (1, \dots, 1)^T$  and  $e_2 = (1, \dots, 1)^T$  are  $m$  and  $n$  vectors respectively. This is a nonlinear system of  $5n+5m$  equations in  $5n+5m$  unknowns. It has a unique solution in the strict interior of an appropriate orthant in primal-dual space  $\{(\Delta, w, g, t, p, y, z, v, s, q) : w, g, t, p, z, v, s, q \geq 0\}$ .

The central path is an arc of strictly feasible points. It is parameterized by the scalar  $\mu$ , and each point on the central path solves the primal-dual system (5). As  $\mu$  tends to zero, the central path converges to an optimal solution to both primal and dual problems. For a value of  $\mu$  let  $(\Delta, w, g, \dots, q)$  denote the current point in the orthant. Our aim is to find the direction vectors  $(\Delta\Delta, \Delta w, \dots, \Delta q)$  such that the new point  $(\Delta + \Delta\Delta, w + \Delta w, \dots, q + \Delta q)$  lies approximately on the primal-dual central path at the point  $(\Delta_\mu, w_\mu, \dots, q_\mu)$ . We see that the new point  $(\Delta + \Delta\Delta, w + \Delta w, \dots, q + \Delta q)$ , if it were to lie exactly on the central path at  $\mu$ , would be defined by

$$\begin{aligned} -H_k \Delta\Delta + \nabla h_k^T \Delta y + \Delta z - \Delta s = \sigma \\ -\Delta y - \Delta q + \Delta v = y + q - v \equiv \beta \\ V^{-1} W \Delta v + \Delta w = \mu V^{-1} e_1 - w - V^{-1} \Delta V \Delta w \equiv \gamma_w \\ P^{-1} Q \Delta p + \Delta q = \mu P^{-1} e_1 - q - P^{-1} \Delta P \Delta q \equiv \gamma_q \\ G^{-1} Z \Delta g + \Delta z = \mu G^{-1} e_2 - z - G^{-1} \Delta G \Delta z \equiv \gamma_z \\ T^{-1} S \Delta t + \Delta s = \mu T^{-1} e_2 - s - T^{-1} \Delta T \Delta s \equiv \gamma_s \end{aligned} \quad (6)$$

$$\nabla h_k \Delta\Delta - \Delta w = w + b - \nabla h_k \Delta - h_k \equiv \rho$$

$$\Delta w + \Delta p = r - w - p \equiv \alpha$$

$$\Delta\Delta - \Delta g = l - \Delta - x_k + g \equiv \nu$$

$$\Delta\Delta + \Delta t = u - \Delta - x_k - t \equiv \tau$$

where we have introduced notations

$$\sigma \equiv H_k \Delta\Delta + \nabla F_k^T \Delta\Delta - \nabla h_k^T y - z + s \quad \text{and} \quad \beta, \rho, \alpha, \tau, \nu,$$

$\gamma_w, \gamma_q, \gamma_z, \gamma_s$  as short-hands for the right-hand side

expressions. This is almost a linear system for the direction vectors  $(\Delta\Delta, \Delta w, \dots, \Delta q)$ . The only nonlinearities appear on the right-side hand of the complementarity equations (*i.e.*, in  $\gamma_w, \gamma_q, \gamma_z, \gamma_s$ , the  $\gamma$ -vectors).

The algorithm implements a *predictor-corrector* [8] approach to find a good approximation solution to the equations (6). First, a predictor direction  $(\Delta\Delta^p, \Delta w^p, \dots, \Delta q^p)$  is computed from (6) ignoring the  $\mu$  and  $\Delta$ -terms of the  $\gamma$ -vectors. Then an estimate of an appropriate target value for  $\mu$  is made using

$$\mu = \bar{\delta} \left( \bar{z}^T \bar{g} + \bar{s}^T \bar{t} + \bar{v}^T \bar{w} + \bar{p}^T \bar{q} \right) / (2m + 2n) \quad \text{with}$$

$$\bar{z} = z + \bar{\alpha}^p \Delta z^p, \bar{g} = g + \bar{\alpha}^p \Delta g^p, \dots, \bar{q} = q + \bar{\alpha}^p \Delta q^p \text{ and}$$

$\bar{\delta} = \left( \frac{(\bar{\alpha}^p - 1)}{(\bar{\alpha}^p + 10)} \right)^2$  where  $\bar{\alpha}^p$  is the longest step length that can be taken along this direction before violating the nonnegative conditions  $w, g, t, p, z, v, s, q \geq 0$  with an upper bound of 1. The corrector step  $(\Delta\Delta, \Delta w, \dots, \Delta q)$  is then obtained by reinstalling the  $\mu$  and the  $\Delta$ -terms on the  $\gamma$ -vectors in (6). This step is used to move to a new point in primal-dual space. Again we calculate the maximum step  $\bar{\alpha}$  that can be taken along this direction before violating the nonnegativity conditions, yielding the new point  $\Delta = \Delta + \bar{\alpha}\Delta\Delta, w = w + \bar{\alpha}\Delta w, \dots, q = q + \bar{\alpha}\Delta q$ .

Implementation details to provide initial values for all the variables in this interior-point paradigm as well as to solve system (6) are described in [2]. A solution of the quadratic subproblem is declared primal/dual feasible if the relative measures of primal and dual infeasibilities are less than  $10^{-4}$ . Thus, the QP subproblem has a solution  $(\Delta_k, \pi_k)$  with  $\Delta_k = \Delta$  and  $\pi_k^T = (v, q, z, s)$ .

### 3 A Line Search Filter Method in SQP

After a search direction  $\Delta_k$  has been computed, we consider a backtracking line search procedure, where a decreasing sequence of step sizes  $\alpha_{k,l} \in (0, 1]$  ( $l = 0, 1, \dots$ ), with  $\lim_l \alpha_{k,l} = 0$ , is tried until an acceptance criterion is satisfied. The procedure that decides which trial step is accepted is a “filter method”. Traditionally, a trial step size  $\alpha_{k,l}$  is accepted if the corresponding trial point  $x_k(\alpha_{k,l}) = x_k + \alpha_{k,l}\Delta_k$ ,  $\lambda_k(\alpha_{k,l}) = \lambda_k + \alpha_{k,l}\xi_k$ ,  $ss_k(\alpha_{k,l}) = ss_k + \alpha_{k,l}\zeta_k$ , provides sufficient reduction of a merit function, such as the augmented Lagrangian function [4], which has the form

$$L(x, \lambda, ss; \eta) = F(x) - \lambda^T (\bar{h}(x) - ss) + \frac{\eta}{2} \bar{\theta}(x, ss)^2 \quad (7)$$

where the infeasibility measure  $\bar{\theta}(x, ss)$  is given by

$$\bar{\theta}(x, ss) = \left\| (\bar{h}(x) - ss) \right\|_2$$

and  $\eta$  is a positive penalty parameter. Here,  $ss$  is a vector of nonnegative slack variables that are used only in the line search procedure and at the beginning of iteration  $k$  is taken as  $ss_k = \max(0, \bar{h}(x_k))$ . We treat the elements of  $\lambda$  as additional variables so that  $\pi$  is used to define a “search direction”,  $\xi$ , for the multiplier estimate  $\lambda$ , and the line search is performed with respect to  $x, \lambda$  and  $ss$ . At iteration  $k$ ,

a vector triple  $d_k^T = (\Delta_k, \xi_k, \zeta_k)$  is computed to serve as direction of search for the variables  $(x_k, \lambda_k, ss_k)$ . The vectors  $\Delta_k$  and  $\pi_k$  are found from the QP subproblem (3). The  $\xi_k$  is defined as  $\xi_k = \lambda_k - \pi_k$  and the vector  $\zeta_k$  satisfies  $\nabla \bar{h}_k \Delta_k + \bar{h}_k = \zeta_k + ss_k$ , from which we can see that  $\zeta_k + ss_k$  is simply the residual of the inequality constraints from problem (3). In order to avoid the determination of an appropriate value of the penalty parameter  $\eta$ , Fletcher and Leyffer [3] proposed the concept of a filter method in the context of a trust region SQP algorithm. The basic idea behind this approach is to interpret the optimization problem (1) as a biobjective optimization problem with two goals: minimizing the constraints violation  $\bar{\theta}(x) = \left\| \min(0, \bar{h}(x)) \right\|_2$  and minimizing the objective function  $F(x)$ . A certain emphasis is placed on the first measure, since a point has to be feasible in order to be an optimal solution of (1).

Following this paradigm, we propose an approach based on the two components of the augmented Lagrangian function (7):

$$\bar{L}(x, \lambda, ss) = F(x) - \lambda^T (\bar{h}(x) - ss) \quad (8)$$

and  $\bar{\theta}(x, ss)$  (or, equivalently,  $\bar{\theta}(x, ss)^2$ ) rather than on  $\bar{\theta}(x)$  and  $F(x)$ . (Recently, in [10] a related approach using the Lagrangian function in a filter trust region based method is proposed.) The trial point  $(x_k(\alpha_{k,l}), \lambda_k(\alpha_{k,l}), ss_k(\alpha_{k,l}))$  is accepted by the filter if it improves feasibility, i.e., if  $\bar{\theta}(x_k(\alpha_{k,l}), ss_k(\alpha_{k,l})) < \bar{\theta}(x_k, ss_k)$ , or if it improves the Lagrangian function (8), i.e., if  $\bar{L}(x_k(\alpha_{k,l}), \lambda_k(\alpha_{k,l}), ss_k(\alpha_{k,l})) < \bar{L}(x_k, \lambda_k, ss_k)$ . Note that this criterion is less demanding than the enforcement of decrease in the penalty function (7) and might in general allow larger steps.

#### 3.1 Sufficient reduction

Line search methods that use a merit function ensure sufficient progress toward the solution by enforcing an Armijo condition for the augmented Lagrangian function (7). Following this idea, we might consider the trial point  $(x_k(\alpha_{k,l}), \lambda_k(\alpha_{k,l}), ss_k(\alpha_{k,l}))$  during the backtracking line search to be acceptable, if the next iterate provides at least as much progress in one of the measures  $\bar{\theta}$  or  $\bar{L}$  that corresponds to a small fraction of the current constraints violation,  $\bar{\theta}(x_k, ss_k)$ , i.e, if

$$\bar{\theta}(x_k(\alpha_{k,l}), ss_k(\alpha_{k,l})) \leq (1 - \gamma_{\bar{\theta}}) \bar{\theta}(x_k, ss_k) \quad (9a)$$

or

$$\bar{L}(x_k(\alpha_{k,l}), \lambda_k(\alpha_{k,l}), ss_k(\alpha_{k,l})) \leq \bar{L}(x_k, \lambda_k, ss_k) - \gamma_{\bar{L}} \bar{\theta}(x_k, ss_k) \quad (9b)$$

holds for fixed constants  $\gamma_{\bar{\theta}}, \gamma_{\bar{L}} \in (0, 1)$ . However, we change to a different sufficient reduction criterion whenever, for the current iterate, we have  $\bar{\theta}(x_k, ss_k) \leq \bar{\theta}^{\min}$ , for some  $\bar{\theta}^{\min} \in (0, \infty]$ , and the following “switching conditions”

$$\begin{aligned} \nabla \bar{L}(x_k, \lambda_k, ss_k)^T d_k < 0 \quad \text{and} \\ \alpha_{k,l} \left[ -\nabla \bar{L}(x_k, \lambda_k, ss_k)^T d_k \right]^{s_{\bar{L}}} > \delta \left[ \bar{\theta}(x_k, ss_k) \right]^{s_{\bar{\theta}}}, \end{aligned} \quad (10)$$

hold with fixed constants  $\delta > 0$ ,  $s_{\bar{\theta}} > 1$ ,  $s_{\bar{L}} > 2s_{\bar{\theta}}$ . If  $\bar{\theta}(x_k, ss_k) \leq \bar{\theta}^{\min}$  and (10) is true for the current iterate, the trial point  $(x_k(\alpha_{k,l}), \lambda_k(\alpha_{k,l}), ss_k(\alpha_{k,l}))$ , has to satisfy the Armijo condition

$$\begin{aligned} \bar{L}(x_k(\alpha_{k,l}), \lambda_k(\alpha_{k,l}), ss_k(\alpha_{k,l})) \leq \bar{L}(x_k, \lambda_k, ss_k) + \\ + \eta_{\bar{L}} \alpha_{k,l} \nabla \bar{L}(x_k, \lambda_k, ss_k)^T d_k, \end{aligned} \quad (11)$$

instead of (9), in order to be acceptable. Here,  $\eta_{\bar{L}} \in (0, 0.5)$  is a constant. According to previous publications on filter methods we call a trial step size  $\alpha_{k,l}$  for which (10) holds, a “ $\bar{L}$ -step size”. Similarly, if an  $\bar{L}$ -step size is accepted as the final step size  $\alpha_k$  in iteration  $k$ , we refer to  $k$  as an “ $\bar{L}$ -type iteration”. At each iteration  $k$ , the algorithm also maintains a “filter”, here denoted by  $F_k \subseteq \{(\bar{\theta}, \bar{L}) \in R^2 : \bar{\theta} \geq 0\}$ . Following the ideas in [12, 13, 14], the filter here is not defined by a list but as a set  $F_k$  that contains those combinations of constraints violation values  $\bar{\theta}$  and Lagrangian function values  $\bar{L}$ , that are prohibited for a successful trial point in iteration  $k$ . So, during the line search, a trial point  $(x_k(\alpha_{k,l}), \lambda_k(\alpha_{k,l}), ss_k(\alpha_{k,l}))$  is rejected, if

$(\bar{\theta}(x_k(\alpha_{k,l}), ss_k(\alpha_{k,l})), \bar{L}(x_k(\alpha_{k,l}), \lambda_k(\alpha_{k,l}), ss_k(\alpha_{k,l}))) \in F_k$ . The authors in [12, 13, 14] apply this simplified notation to active set SQP and barrier interior-point line search based algorithms. At the beginning of the optimization, the filter is initialized to

$$F_0 = \left\{ (\bar{\theta}, \bar{L}) \in R^2 : \bar{\theta} \geq \bar{\theta}^{\max} \right\} \quad (12)$$

for some  $\bar{\theta}^{\max}$ , so that the algorithm will never allow trial points to be accepted that have a constraint violation larger than  $\bar{\theta}^{\max}$ . Later, the filter is augmented, using the update formula

$$F_{k+1} = F_k \cup \left\{ (\bar{\theta}, \bar{L}) \in R^2 : \bar{\theta} \geq (1 - \gamma_{\bar{\theta}}) \bar{\theta}_k \text{ and } \bar{L} \geq \bar{L}_k - \gamma_{\bar{L}} \bar{\theta}_k \right\} \quad (13)$$

after every iteration in which the accepted trial step size does not satisfy the switching conditions (10). This ensures that the iterates cannot return to the neighborhood of  $x_k$ . On the other hand, if both (10) and (11) hold for the accepted step size, the filter remains unchanged.

Overall, this procedure ensures that the algorithm cannot cycle, for example between two points that alternatively decrease the constraints violation and the Lagrangian function  $\bar{L}$ . Finally, in some cases it is not possible to find a trial step size  $\alpha_{k,l}$  that satisfies the above criteria. We define a minimum desired step size using linear models of the involved functions

$$\alpha_k^{\min} := \gamma_{\alpha} \begin{cases} \min \left\{ \gamma_{\bar{\theta}}, \frac{\gamma_{\bar{L}} \bar{\theta}_k}{-\nabla \bar{L}_k^T d_k}, \frac{\delta [\bar{\theta}_k]^{s_{\bar{\theta}}}}{[-\nabla \bar{L}_k^T d_k]^{s_{\bar{L}}}} \right\} \\ \quad \text{if } \nabla \bar{L}_k^T d_k < 0 \text{ and } \bar{\theta}_k \leq \bar{\theta}^{\min} \\ \min \left\{ \gamma_{\bar{\theta}}, \frac{\gamma_{\bar{L}} \bar{\theta}_k}{-\nabla \bar{L}_k^T d_k} \right\} \\ \quad \text{if } \nabla \bar{L}_k^T d_k < 0 \text{ and } \bar{\theta}_k > \bar{\theta}^{\min} \\ \gamma_{\bar{\theta}}, \text{ otherwise} \end{cases} \quad (14)$$

with a safety factor  $\gamma_{\alpha} \in (0, 1]$ . If the backtracking line search finds a trial step size  $\alpha_{k,l} < \alpha_k^{\min}$ , the algorithm reverts to a feasibility restoration phase. Here, the algorithm tries to find a new iterate  $(x_{k+1}, \lambda_{k+1}, ss_{k+1})$  which is acceptable to the current filter and for which (9) holds, by reducing the constraints violation within an iterative process.

Our interior-point SQP filter line search algorithm for solving inequality constrained optimization problems is as follows:

### Algorithm 1

*Given:* Starting point  $(x_0, \lambda_0, ss_0)$  with  $ss_0 = \max(0, \bar{h}(x_0))$ ; constants  $\bar{\theta}^{\max} \in (\bar{\theta}(x_0, ss_0), \infty]$ ,

$\bar{\theta}^{\min} > 0$ ;  $\gamma_{\bar{\theta}}, \gamma_{\bar{L}} \in (0, 1)$ ;  $\delta > 0$ ;  $\gamma_{\alpha} \in (0, 1]$ ;  $s_{\bar{\theta}} > 1$ ;  
 $s_{\bar{L}} > 2s_{\bar{\theta}}$ ;  $\eta_{\bar{L}}, \eta_{\bar{\theta}_2} \in (0, 1)$

1. *Initialize*. Initialize the filter (using (12)) and the iteration counter  $k \leftarrow 0$ .

2. *Check convergence*. Stop if  $x_k$  is a stationary point of the problem (1), i.e., if it satisfies the KT conditions (2) for some  $\lambda \in R^m$ .

3. *Compute search direction*. Compute the search direction  $\Delta_k$  and the Lagrange multiplier  $\pi_k$  from the linear system (6) (using the interior-point strategy presented in Section 2).

4. *Backtracking line search*.

4.1 *Initialize line search*. Set  $ss_k = \max(0, \bar{h}(x_k))$ ,

$\xi_k = \lambda_k - \pi_k$ ,  $\zeta_k = \nabla \bar{h}_k \Delta_k + \bar{h}_k - ss_k$ ,  $\alpha_{k,l} = 1$ ,  $l \leftarrow 0$ .

4.2 *Compute new trial point*. If the trial step size becomes too small, i.e.,  $\alpha_{k,l} < \alpha_k^{\min}$  with  $\alpha_k^{\min}$  defined by (14), go to feasibility restoration phase in step 8. Otherwise, compute the trial points  $x_k(\alpha_{k,l}) = x_k + \alpha_{k,l} \Delta_k$ ,  $\lambda_k(\alpha_{k,l}) = \lambda_k + \alpha_{k,l} \xi_k$ ,

$ss_k(\alpha_{k,l}) = ss_k + \alpha_{k,l} \zeta_k$ .

4.3 *Check acceptability to the filter*. If  $(\bar{\theta}(x_k(\alpha_{k,l}), ss_k(\alpha_{k,l})), \bar{L}(x_k(\alpha_{k,l}), \lambda_k(\alpha_{k,l}), ss_k(\alpha_{k,l}))) \in F_k$ , reject the trial step size and go to step 4.5.

4.4 *Check sufficient decrease with respect to current iterate*.

*Case I:  $\alpha_{k,l}$  is an  $\bar{L}$ -step size (i.e., (10) holds):* If the Armijo condition (11) for the  $\bar{L}$  function holds, accept the trial step and go to step 5. Otherwise, go to step 4.5.

*Case II:  $\alpha_{k,l}$  is not an  $\bar{L}$ -step size:* If (9) holds, accept the trial step and go to step 5. Otherwise, go to step 4.5.

4.5. *Choose new trial step size*. Set  $\alpha_{k,l+1} \leftarrow \alpha_{k,l}/2$ ,  $l \leftarrow l+1$ , and go back to step 4.2.

5. *Accept trial point*. Set  $\alpha_k \leftarrow \alpha_{k,l}$ ,  $x_{k+1} \leftarrow x_k(\alpha_k)$ , and  $\lambda_{k+1} \leftarrow \lambda_k(\alpha_k)$ .

6. *Augment filter if necessary*. If  $k$  is not an  $\bar{L}$ -type iteration, augment the filter using (13). Otherwise, leave the filter unchanged.

7. *Continue with next iteration*. Increase the iteration counter  $k \leftarrow k+1$  and go back to step 2.

8. *Feasibility restoration phase*. Use a restoration algorithm to produce a point  $(x_{k+1}, \lambda_{k+1}, ss_{k+1})$  that is acceptable to the filter, i.e.,

$(\bar{\theta}(x_{k+1}, ss_{k+1}), \bar{L}(x_{k+1}, \lambda_{k+1}, ss_{k+1})) \in F_k$ . Augment the filter using (13) and continue with the regular iteration in step 7.

### 3.2 Feasibility restoration phase

In this section we present a restoration algorithm. The task of the restoration phase is to compute a new iterate acceptable to the filter by decreasing the infeasibility, whenever the regular backtracking line search procedure cannot make sufficient progress and the step size becomes too small. To compute a trial point that sufficiently decreases infeasibility, we introduce the function  $\bar{\theta}_2(x, ss) = \frac{1}{2} \|(\bar{h}(x) - ss)\|_2^2$ . The restoration algorithm herein presented works with the step framework  $\bar{d}^T = (\Delta, \zeta)$  that should be a descent direction for  $\bar{\theta}_2(x, ss)$ . In fact,

$$\begin{aligned} \bar{d}^T \nabla \bar{\theta}_2 &= (\bar{h} - ss)^T \nabla \bar{h} \Delta - (\bar{h} - ss)^T \zeta \\ &= (\bar{h} - ss)^T (\nabla \bar{h} \Delta - \zeta) \\ &= -(\bar{h} - ss)^T (\bar{h} - ss) = -2\bar{\theta}_2 < 0. \end{aligned}$$

Additionally, we also ensure that the new iterate  $x_{k+1}$  does not deviate too much from the current iterate  $x_k$  (see step 5 in Algorithm 2).

Several other restoration algorithms are plausible but we chose the following one because it is consistent with the step calculation of our interior-point SQP filter line search method:

#### Algorithm 2 (restoration algorithm)

0. Set  $x_{k,0} = x_k$ ,  $\lambda_{k,0} = \lambda_k$ ,  $ss_{k,0} = ss_k$ ,  $l = 0$  and start with step 4.

1. If  $(x_{k,l}, \lambda_{k,l}, ss_{k,l})$  is acceptable to the filter (conditions (9)) then set  $x_{k+1} = x_{k,l}$ ,  $\lambda_{k+1} = \lambda_{k,l}$ , and stop restoration.

2. Compute  $\Delta_{k,l}$  and  $\pi_{k,l}$  by solving the QP subproblem (3), with  $(x_k, \lambda_k) = (x_{k,l}, \lambda_{k,l})$ .

3. Compute  $ss_{k,l}$ ,  $\xi_{k,l}$ ,  $\zeta_{k,l}$  and define the vector  $(\bar{d}_{k,l})^T = (\Delta_{k,l}, \zeta_{k,l})$  which is used as direction of search for the variables  $(x_{k,l}, ss_{k,l})$ .

4. Set  $\alpha_k = 1$ .

5. If

$$\bar{\theta}_2(x_{k,l}(\alpha_k), ss_{k,l}(\alpha_k)) \leq \bar{\theta}_2(x_{k,l}, ss_{k,l}) + \alpha_k \eta_{\bar{\theta}_2} \nabla \bar{\theta}_2^T \bar{d}_{k,l},$$

and  $\|x_{k,l}(\alpha_k) - x_k\| \leq \varepsilon_{\bar{\theta}_2} (1 + \|x_{k,l}(\alpha_k)\|)$

then set  $x_{k,l+1} = x_{k,l}(\alpha_k)$ ,  $ss_{k,l+1} = ss_{k,l}(\alpha_k)$ ,  $\lambda_{k,l+1} = \lambda_{k,l}(\alpha_k)$ ,  $l = l+1$ , and return to step 1. Otherwise  $\alpha_k \leftarrow \alpha_k/2$ , and repeat step 5.

#### 4 Nonmonotone Line Search

Nonmonotone line search methods that use a merit function ensure sufficient progress toward the solution by imposing that the merit function value of each new iterate satisfies an Armijo condition with respect to the maximum merit function value of a prefixed number of previous iterates, see for example [6]. Thus, the condition which implies a monotonic decrease of the merit function is relaxed without affecting the global convergence. Transposing this idea to this interior-point SQP filter line search framework, we now should consider the trial point  $(x_k(\alpha_{k,l}), \lambda_k(\alpha_{k,l}), ss_k(\alpha_{k,l}))$ , during the backtracking line search technique, to be acceptable if it leads to sufficient progress in one of the two measures ( $\bar{\theta}$  or  $\bar{L}$ ) compared to a previous iterate that yields the maximum value of the corresponding measure within the last  $m(k)$  iterates, where  $m(0) = 0$  and  $0 \leq m(k) \leq \min\{m(k-1)+1, M\}$ ,  $k \geq 1$  ( $M$  is a nonnegative integer). So, the conditions (9a) and (9b) are now rewritten as follows:

$$\bar{\theta}(x_k(\alpha_{k,l}), ss_k(\alpha_{k,l})) \leq (1 - \gamma_{\bar{\theta}}) \max_{0 \leq j \leq m(k)} \bar{\theta}(x_{k-j}, ss_{k-j}) \quad (15a)$$

or

$$\bar{L}(x_k(\alpha_{k,l}), \lambda_k(\alpha_{k,l}), ss_k(\alpha_{k,l})) \leq \max_{0 \leq j \leq m(k)} \bar{L}(x_{k-j}, \lambda_{k-j}, ss_{k-j}) - \gamma_{\bar{L}} \bar{\theta}(x_k, ss_k). \quad (15b)$$

Similarly in the switching conditions (10), only the second condition is modified as follows:

$$\alpha_{k,l} \left[ -\nabla \bar{L}(x_k, \lambda_k, ss_k)^T d_k \right]^{s_{\bar{L}}} > \delta \left[ \max_{0 \leq j \leq m(k)} \bar{\theta}(x_{k-j}, ss_{k-j}) \right]^{s_{\bar{\theta}}}. \quad (16)$$

The acceptability Armijo condition (11) for the step size is now relaxed and given by

$$\bar{L}(x_k(\alpha_{k,l}), \lambda_k(\alpha_{k,l}), ss_k(\alpha_{k,l})) \leq \max_{0 \leq j \leq m(k)} \bar{L}(x_{k-j}, \lambda_{k-j}, ss_{k-j}) + \eta_{\bar{L}} \alpha_{k,l} \nabla \bar{L}(x_k, \lambda_k, ss_k)^T d_k. \quad (17)$$

The inclusion of the nonmonotone strategy also affects the update formula to augment the filter which has now the following form

$$F_{k+1} = F_k \cup \left\{ (\bar{\theta}, \bar{L}) \in R^2 : \bar{\theta} \geq (1 - \gamma_{\bar{\theta}}) \max_{0 \leq j \leq m(k)} \bar{\theta}(x_{k-j}, ss_{k-j}) \text{ and } \bar{L} \geq \max_{0 \leq j \leq m(k)} \bar{L}(x_{k-j}, \lambda_{k-j}, ss_{k-j}) - \gamma_{\bar{L}} \bar{\theta}(x_k, ss_k) \right\}. \quad (18)$$

Finally the computation of the  $\alpha_k^{\min}$  (last (14)) has to take into account the previous relaxed conditions and is given by

$$\alpha_k^{\min} := \gamma_{\alpha} \begin{cases} \min \left\{ \gamma_{\bar{\theta}}, (\bar{L}_k - \max_{0 \leq j \leq m(k)} \bar{L}(x_{k-j}, \lambda_{k-j}, ss_{k-j})) + \gamma_{\bar{L}} \bar{\theta}_k \right\} / (-\nabla \bar{L}_k^T d_k), \frac{\delta \left[ \max_{0 \leq j \leq m(k)} \bar{\theta}(x_{k-j}, ss_{k-j}) \right]^{s_{\bar{\theta}}}}{\left[ -\nabla \bar{L}_k^T d_k \right]^{s_{\bar{L}}}} \right\} \\ \text{if } \nabla \bar{L}_k^T d_k < 0 \text{ and } \bar{\theta}_k \leq \bar{\theta}^{\min} \\ \min \left\{ \gamma_{\bar{\theta}}, (\bar{L}_k - \max_{0 \leq j \leq m(k)} \bar{L}(x_{k-j}, \lambda_{k-j}, ss_{k-j})) + \gamma_{\bar{L}} \bar{\theta}_k \right\} / (-\nabla \bar{L}_k^T d_k) \\ \text{if } \nabla \bar{L}_k^T d_k < 0 \text{ and } \bar{\theta}_k > \bar{\theta}^{\min} \\ \gamma_{\bar{\theta}}, \text{ otherwise.} \end{cases} \quad (19)$$

The algorithm for the interior-point SQP filter nonmonotone line search approach is similar to Algorithm 1. The references to equations (9), (10), (11), (13) and (14) should now be replaced by (15), (16), (17), (18) and (19) respectively. Further, in step 1,  $m(0)$  should be initialized to 0 and in step 7, after the increase of the iteration counter  $k$ , the update  $m(k) = \min\{m(k-1)+1, M\}$  is required.

The modifications to Algorithm 2 of the feasibility restoration phase are as follows:

- replace the Armijo condition in step 5 by the relaxed condition  $\bar{\theta}_2(x_{k,l}(\alpha_k), ss_{k,l}(\alpha_k)) \leq \max_{0 \leq j \leq m(l)} \bar{\theta}_2(x_{k-j}, ss_{k-j}) + \alpha_k \eta_{\bar{\theta}_2} \nabla \bar{\theta}_2^T \bar{d}_{k,l}$
- include  $m(0)=0$  in step 0 and the update  $m(l) = \min\{m(l-1)+1, M\}$  in step 5 before returning to step 1.

#### 5 Results and Conclusions

To test this SQP framework based on the interior-point strategy with a filter line search method we selected 32 small inequality constrained problems

from the Hock and Schittkowski (HS) collection [7] and other 6 problems, usually known in the literature as difficult problems, named P0-P5 in the Table 1. P0-P4 are from [1] and P5 from [15]. The tests were done in double precision arithmetic with a Pentium 4 and Fortran 90. For the successful termination of the algorithm, the iterative sequence of  $x$ -values must converge and the final point must satisfy the first-order Kuhn-Tucker conditions (see (2)) with a  $10^{-4}$  tolerance.

The chosen values for the constants are similar to the ones used in [14]:  $\bar{\theta}^{\max} = 10^4 \max\{1, \bar{\theta}(x_0, ss_0)\}$ ,  $\bar{\theta}^{\min} = 10^{-4} \max\{1, \bar{\theta}(x_0, ss_0)\}$ ,  $\gamma_{\bar{\theta}} = 10^{-5}$ ,  $\gamma_{\bar{L}} = 10^{-5}$ ,  $\delta = 1$ ,  $\gamma_{\alpha} = 0.05$ ,  $s_{\bar{\theta}} = 1.1$ ,  $s_{\bar{L}} = 2.3$ ,  $\eta_{\bar{L}} = 10^{-4}$ ,  $\eta_{\bar{\theta}_2} = 10^{-4}$  and  $\varepsilon_{\bar{\theta}_2} = 0.1$ , where  $x_0$  is the starting point and  $ss_0 = \max(0, \bar{h}(x_0))$ .

The numerical results concerning our interior-point SQP filter line search method are reported in the first part of Table 1. For comparative purposes, we include in the second part of the table the results obtained when a line search method based on the merit function (7) is used as in [2] instead of the herein proposed filter method. Each set of two columns contains the number of QP subproblems solved ( $N_{QP}$ ) and the number of function evaluations ( $N_{fe}$ ) -  $\bar{L}$  for the *Filter Method* and  $L$  for the *Merit Function* approach.

In most problems the results are similar while in 9 of the 38 problems the filter method requires less function evaluations than the merit function based line search. Slightly better results were obtained with the merit function in 5 problems.

The implementation of the nonmonotone approach, as described in Section 4, for  $M = 5$ , gives similar results for the majority of the tested problems with the following exceptions:

HS1 ( $N_{QP} = 73, N_{fe} = 91$ ), HS2 ( $N_{QP} = 26, N_{fe} = 27$ ),

HS5 ( $N_{QP} = 18, N_{fe} = 28$ ), HS17 ( $N_{QP} = 14, N_{fe} = 15$ ),

HS18 ( $N_{QP} = 6, N_{fe} = 7$ ), HS21 ( $N_{QP} = 2, N_{fe} = 3$ ),

HS34 ( $N_{QP} = 7, N_{fe} = 13$ ), P4 ( $N_{QP} = 11, N_{fe} = 12$ ).

Although the observed differences are not significant a comparative analysis is possible considering the cumulative results. In the monotone filter approach the ratio between the number of function evaluations and the number of QP subproblems solved is 2.8 while in the nonmonotone filter approach the ratio is 2 (Total  $N_{QP} = 379$ , Total  $N_{fe} = 776$ ). Thus a slightly more efficient framework is achieved when the

acceptability criteria for the step size are relaxed. The ratio for the merit function based method is 3.

Table 1: Comparative results

Problem	Filter Method		Merit Function	
	$N_{QP}$	$N_{fe}$	$N_{QP}$	$N_{fe}$
HS1	44	59	44	59
HS2	13	16	13	16
HS3	1	2	1	2
HS4	2	3	2	10
HS5	9	31	9	31
HS15	8	9	10	61
HS16	5	6	5	6
HS17	13	15	12	15
HS18	9	43	6	67
HS19	7	8	7	8
HS20	8	9	7	9
HS21	2	13	2	13
HS23	6	7	6	7
HS24	6	7	6	7
HS30	11	12	11	12
HS31	7	8	7	68
HS32	5	6	5	6
HS33	5	6	5	6
HS34	7	17	8	31
HS35	2	3	2	3
HS36	6	7	6	29
HS37	9	10	9	10
HS38	13	14	13	14
HS41	12	13	12	13
HS44	6	7	6	16
HS45	5	6	5	6
HS53	3	4	3	4
HS55	1	2	1	2
HS60	8	9	12	52
HS63	12	15	9	12
HS64	19	173	19	173
HS65	9	10	10	54
P0	22	197	15	17
P1	5	6	5	6
P2	4	5	4	5
P3	F	-	F	-
P4	11	145	11	96
P5	15	16	15	16
Cum.	330	919	323	962

Cum. - cumulative results; F - failure (unbounded objective)

When solving the problems HS2, HS16, HS20, HS38 and HS44, our algorithms converged to solutions that are different from the ones reported in the literature [7]. Thus, we decided to run well-known nonlinear programming solvers in order to compare the solutions. We also included in this analysis the set P0-P5 of the most difficult problems. This comparative analysis includes two solvers that are available in the NEOS Server (<http://www-neos.mcs.anl.gov/neos/>). This Server provides the

possibility to run problems on powerful machines in a user friendly manner through the internet. The chosen solvers were:

- FILTER, which is an SQP active set trust region algorithm with a filter to promote global convergence [3];
- SNOPT, which is a specific SQP implementation of an active-set method based on a smooth augmented Lagrangian merit function [5].

Table 2 reports the results that we obtained for the above referred problems. For the problems HS2, HS16 and HS20, both solvers also converged to the same solutions that we found. Comparisons with SNOPT and FILTER are not meant to be a rigorous assessment of the performance of our algorithms since the termination criteria are not comparable.

Table 2: FILTER and SNOPT results

Problem	SNOPT		FILTER	
	$N_{OP}$	$N_{fe}$	$N_{OP}$	$N_{fe}$
HS2	18	15	8	9
HS16	1	5	4	5
HS20	1	5	4	5
HS38	160	119	53	54
HS44	2	ni	1	2
P0	1	ni	23	21
P1	3	4	2	3
P2	8	5	4	5
P3	F	-	F	-
P4	16	44	11	12
P5	2	7	33	34

F – failure (unbounded objective); ni – no information is available on the output file sent by NEOS

Despite some similarities, our SQP monotone filter line search method has some differences from the SQP filter line search method proposed and analyzed in [12, 13]. The differences are mainly on the defined  $\alpha_k^{\min}$  and on the feasibility restoration phase. The steps computed from the QP subproblems are in our case obtained by an interior-point method. The implementation of a nonmonotone strategy in a filter line search framework is new. The obtained results seem promising and a global convergence analysis of the proposed method will be carried out in the near future.

#### References:

[1] J.V. Burke and S.-P. Han, A robust sequential quadratic programming method, *Mathematical Programming*, Vol.43, 1989, pp. 277-303.  
 [2] M.F.P. Costa and E.M.G.P. Fernandes, A SQP based on an interior-point strategy for nonlinear inequality constrained optimization problems,

*Proceedings of the VII Congreso Galego de Estatística e Investigación de Operacións*, ISBN 972-99841-0-7 (CD-Rom), 2005.

[3] R. Fletcher and S. Leyffer, Nonlinear programming without a penalty function, *Mathematical Programming*, Vol.91, 2002, pp. 239-269.  
 [4] P.E. Gill, W. Murray and M.A. Saunders, Some theoretical properties of an augmented Lagrangian merit function, *Technical Report SOL 86-6R*, 1986.  
 [5] P.E. Gill, W. Murray and M.A. Saunders, SNOPT: An SQP algorithm for large-scale constrained optimization, *SIAM Journal on Optimization*, Vol.12, 2002, pp. 979-1006.  
 [6] L. Grippo, F. Lampariello and S. Lucidi, A class of nonmonotone stabilization methods in unconstrained optimization, *Numerische Mathematik*, Vol.59, 1991, pp. 779-805.  
 [7] W. Hock and K. Schittkowski, *Test Examples for Nonlinear Programming*, Springer-Verlag, 1981.  
 [8] S. Mehrotra, On the implementation of a (primal-dual) interior point method, *SIAM Journal on Optimization*, Vol.2, 1992, pp. 575-601.  
 [9] J. Nocedal and S.J. Wright, *Numerical Optimization*, Springer-Verlag, 2001.  
 [10] S. Ulbrich, On the superlinear local convergence of a filter-SQP method, *Mathematical Programming*, Vol.100, 2004, pp. 217-245.  
 [11] R.J. Vanderbei, LOQO: An interior-point code for quadratic programming, *Technical Report SOR 94-15*, 1994.  
 [12] A. Wachter and L.T. Biegler, Line search filter methods for nonlinear programming: motivation and global convergence, *SIAM Journal on Optimization*, Vol.16 (1), 2005, pp. 1-31.  
 [13] A. Wachter and L.T. Biegler, Line search filter methods for nonlinear programming: local convergence, *SIAM Journal on Optimization*, Vol.16 (1), 2005, pp. 32-48.  
 [14] A. Wachter and L.T. Biegler, On the implementation of an interior-point filter line search algorithm for large-scale nonlinear programming, *Mathematical Programming*, Vol.106, 2006, pp. 25-57.  
 [15] G. Zhou, A modified SQP method and its global convergence, *Journal of Global Optimization*, Vol.11, 1997, pp. 193-205.