

Reverse Order Law for the Core Inverse in Rings

Honglin Zou^{*}, Jianlong Chen[†], Pedro Patrício[‡]

^{*}College of Mathematics, Yangzhou University, Yangzhou 225002, China

^{*}School of Mathematics and Statistics, Hubei Normal University,
Huangshi 435002, China

[†]School of Mathematics, Southeast University, Nanjing 210096, China

[‡]CMAT-Centro de Matemática and Departamento de Matemática e Aplicações,
Universidade do Minho, Braga 4710-057, Portugal

Abstract: In this paper, necessary and sufficient conditions of the one-sided reverse order law $(ab)^{\oplus} = b^{\oplus}a^{\oplus}$, the two-sided reverse order law $(ab)^{\oplus} = b^{\oplus}a^{\oplus}$ and $(ba)^{\oplus} = a^{\oplus}b^{\oplus}$ for the core inverse are given in rings with involution. In addition, the mixed-type reverse order laws, such as $(ab)^{\#} = b^{\oplus}(abb^{\oplus})^{\oplus}$, $a^{\oplus} = b(ab)^{\#}$ and $(ab)^{\#} = b^{\oplus}a^{\oplus}$, are also considered.

Keywords: core inverse; group inverse; reverse order law

AMS Subject Classifications: 15A09; 16W10; 16B99

1 Introduction

Let $\mathbb{C}_{n \times n}$ denote the set of all $n \times n$ complex matrices. It is well known that $(AB)^{-1} = B^{-1}A^{-1}$, where $A, B \in \mathbb{C}_{n \times n}$ are invertible. The previous equality is called the reverse order law for the ordinary inverse. In general, the equality doesn't hold when the ordinary inverse is replaced by the generalized inverse. In 1966, Greville [13] first gave a necessary and sufficient condition of the reverse order law $(AB)^{\dagger} = B^{\dagger}A^{\dagger}$ for the Moore-Penrose inverse. Since then, many authors studied the reverse order law for various classes of generalized inverses in the setting of complex matrices, operators, and elements of rings with involution. For example, Deng [10] investigated some necessary and sufficient conditions of the reverse order law $(ab)^{\#} = b^{\#}a^{\#}$ for the group inverse of linear bounded operators on Hilbert spaces. In [23], Mosić and Djordjević extended the results of [10] to the ring case, giving some new conditions and providing simpler and more transparent proofs to already existing conditions. Recently, Mary [19] provided equivalent conditions for the two-sided reverse order law $(ab)^{\#} = b^{\#}a^{\#}$ and $(ba)^{\#} = a^{\#}b^{\#}$ for the group inverse in semigroups and rings. In [20, 22, 24], Mosić

^{*}Corresponding author. E-mail: honglinzou@163.com

[†]E-mail: jlchen@seu.edu.cn

[‡]E-mail: pedro@math.uminho.pt

et al. considered the mixed-type reverse order laws in rings, such as $(ab)^\dagger = b^\dagger(a^\dagger abb^\dagger)^\dagger a^\dagger$, $(ab)^\# = b^\dagger(a^\dagger abb^\dagger)^\dagger a^\dagger$, $(ab)^\# = (a^\dagger ab)^\dagger a^\dagger$, and $(ab)^\# = b^\dagger a^\dagger$. More results on the reverse order law for the generalized inverse can be found in [3, 7–9, 11, 16–18, 21, 25, 29].

The following problem on the reverse order law for the core inverse was proposed by Baksalary and Trenkler [2]:

If A^\oplus, B^\oplus , and $(AB)^\oplus$ exist, does it follow that $(AB)^\oplus = B^\oplus A^\oplus$?

This problem attracted researchers' attention. Later, Cohen, Herman and Jayaraman [6] gave several counterexamples for the problem. In [30], Wang and Liu obtained equivalent conditions of the reverse order $(AB)^\oplus = B^\oplus A^\oplus$ by the ranks of matrices. Next, we will continue to consider the reverse law for the core inverse in rings with involution.

The article is motivated by the papers [19, 22, 30]. We present some equivalent conditions for the one-sided reverse order law $(ab)^\oplus = b^\oplus a^\oplus$, the two-sided reverse order law $(ab)^\oplus = b^\oplus a^\oplus$ and $(ba)^\oplus = a^\oplus b^\oplus$ for the core inverse in rings with involution. We also study the mixed-type reverse order laws, such as $(ab)^\# = b^\oplus (abb^\oplus)^\oplus$, $a^\oplus = b(ab)^\#$ and $(ab)^\# = b^\oplus a^\oplus$.

2 Preliminaries

Throughout this paper, R denotes a unital $*$ -ring, that is, a ring with unity 1 and an involution $a \mapsto a^*$ satisfying $(a^*)^* = a$, $(a + b)^* = a^* + b^*$ and $(ab)^* = b^* a^*$ for all $a, b \in R$.

For the readers' convenience, we first recall the definitions of some generalized inverses. An element $a \in R$ is said to be Moore-Penrose invertible with respect to the involution $*$ if the following equations

$$(1) \ axa = a, \quad (2) \ xax = x, \quad (3) \ (ax)^* = ax, \quad (4) \ (xa)^* = xa$$

have a common solution [26]. Such solution is unique if it exists, and is denoted by a^\dagger .

The Drazin inverse [12] of $a \in R$ is the element $x \in R$ which satisfies

$$(1^k) \ a^k = a^{k+1}x, \quad (2) \ xax = x, \quad (5) \ ax = xa, \quad \text{for some } k \geq 1.$$

The element x above is unique if it exists and is denoted by a^D . The smallest such k is called the index of a , and denoted by $\text{ind}(a)$. In particular, when $\text{ind}(a)=1$, the Drazin inverse a^D is called the group inverse of a and it is denoted by $a^\#$.

Baksalary and Trenkler [1] introduced the core inverse for complex matrices. Let $A \in \mathbb{C}_{n \times n}$. A matrix $A^\oplus \in \mathbb{C}_{n \times n}$ is called core inverse of A if

$$(i) \ AA^\oplus = P_A \quad \text{and} \quad (ii) \ \mathcal{R}(A^\oplus) \subseteq \mathcal{R}(A),$$

where P_A is the orthogonal projector onto $\mathcal{R}(A)$, and $\mathcal{R}(A)$ is the column space of A .

Later, the notion of the core inverse for complex matrices was extended to the ring case by Rakić, Dinčić and Djordjević [28]. The core inverse [28] of $a \in R$ is the element $x \in R$ which satisfies

$$(1) \ axa = a, \quad (2) \ xax = x, \quad (3) \ (ax)^* = ax, \quad (6) \ xa^2 = a \quad (7) \ ax^2 = x.$$

The element x above is unique if it exists and is denoted by a^{\oplus} .

Let $\delta = \{1, 2, 3, 4, 5, 6, 7\}$. If $x \in R$ satisfies the equations in (i) for all $i \in \delta$, then x is called a δ -inverse of a . The set of all δ -inverses of a is denoted by $a\{\delta\}$. For example, $a\{1, 2, 5\} = \{a^{\#}\}$. The element $a \in R$ is regular if $a\{1\} \neq \emptyset$. $R^{\#}$, $R^{\{1,3\}}$, R^{\dagger} , R^{\oplus} stand for the set of all group, $\{1,3\}$, Moore-Penrose, core invertible elements of R , respectively. Let $M_n(R)$ denote the ring of $n \times n$ matrices over R .

Given $a \in R$, the following notations will be used:

$$a^0 = \{x \in R : ax = 0\} \text{ and } {}^0a = \{x \in R : xa = 0\}.$$

An element $a \in R$ is said to be EP if $a \in R^{\#} \cap R^{\dagger}$ and $a^{\#} = a^{\dagger}$. An element $a \in R$ satisfying $a^* = a$ is called Hermitian.

Now, we present several known characterizations for some generalized inverses, which play an important role in the sequel.

Lemma 2.1. [31, Theorem 3.1] *Let $a, x \in R$. Then $a \in R^{\oplus}$ with $x = a^{\oplus}$ if and only if*

$$(ax)^* = ax, \quad xa^2 = a \quad \text{and} \quad ax^2 = x.$$

Lemma 2.2. [31, Theorem 2.6] *Let $a \in R$. Then $a \in R^{\oplus}$ if and only if $a \in R^{\#} \cap R^{\{1,3\}}$. In this case, $a^{\oplus} = a^{\#}aa^{(1,3)}$, where $a^{(1,3)} \in a\{1, 3\}$.*

Lemma 2.3. [15, Theorem 1] *Let $a \in R$. Then $a \in R^{\#}$ if and only if $a = a^2x = ya^2$ for some $x, y \in R$. In this case, $a^{\#} = yax = y^2a = ax^2$.*

Lemma 2.4. [14, p. 201] *Let $a, x \in R$. Then x is a $\{1, 3\}$ -inverse of a if and only if $a = x^*a^*a$.*

The following lemmas will also be useful.

Lemma 2.5. [4, Corollary 3.4] *Let $a, x \in R$ with $xa = ax$ and $xa^* = a^*x$. If $a \in R^{\oplus}$, then $xa^{\oplus} = a^{\oplus}x$.*

Lemma 2.6. [5, Cline's formula] *Let $a, b \in R$. If ab is Drazin invertible, then ba is Drazin invertible and $(ba)^D = b((ab)^D)^2a$.*

Lemma 2.7. [28, Theorem 3.1] *Let $a \in R$. Then the following are equivalent:*

- (i) a is EP;
- (ii) $a \in R^{\oplus}$ and $a^{\#} = a^{\oplus}$.
- (iii) $a \in R^{\oplus}$ and $aa^{\oplus} = a^{\oplus}a$.
- (iv) $a \in R^{\#} \cap R^{\dagger}$ and $a^{\dagger} = a^{\oplus}$.

Lemma 2.8. [28, Lemma 2.5, Lemma 2.6] *Let $a, b \in R$.*

- (i) *If $aR \subseteq bR$, then ${}^{\circ}b \subseteq {}^{\circ}a$.*
- (ii) *If b is regular and ${}^{\circ}b \subseteq {}^{\circ}a$, then $aR \subseteq bR$.*
- (iii) *If $Ra \subseteq Rb$, then $b^{\circ} \subseteq a^{\circ}$.*
- (iv) *If b is regular and $b^{\circ} \subseteq a^{\circ}$, then $Ra \subseteq Rb$.*

3 One-sided and Two-sided Reverse Order Laws

In this section, we will give some equivalent conditions of the one-sided $(ab)^\oplus = b^\oplus a^\oplus$, the two-sided reverse order laws $(ab)^\oplus = b^\oplus a^\oplus$ and $(ba)^\oplus = a^\oplus b^\oplus$ for the core inverse in rings.

First, in order to prove our main results, we give new existence criterion for the core inverse as follows:

Theorem 3.1. *Let $a, x \in R$. Then the following are equivalent:*

- (i) $a \in R^\oplus$ and $x = a^\oplus$;
- (ii) $axa = a$, $xR = aR$ and $Rx \subseteq Ra^*$;
- (iii) $axa = a$, ${}^0x = {}^0a$ and $(a^*)^0 \subseteq x^0$;
- (iv) $xax = x$, $xR = aR$ and $Rx = Ra^*$;
- (v) $xax = x$, $xR = aR$ and $Ra^* \subseteq Rx$;
- (vi) $xax = x$, ${}^0x = {}^0a$ and $x^0 \subseteq (a^*)^0$;
- (vii) $a \in R^\#$, $axa = a$, $(ax)^* = ax$, and $xR \subseteq aR$;
- (viii) $a \in R^\#$, $xax = x$, $(ax)^* = ax$, and $aR \subseteq xR$.

Proof. (i) \Rightarrow (ii) and (iv) \Rightarrow (v) are trivial.

(ii) \Rightarrow (iii) and (v) \Rightarrow (vi) follow directly from Lemma 2.8.

(iii) \Rightarrow (iv) Note that a^* is regular. Since $(a^*)^0 \subseteq x^0$, using Lemma 2.8, we have $Rx \subseteq Ra^*$, which implies that $x = t_1 a^*$ for some $t_1 \in R$. Thus, we get

$$x = t_1 a^* = t_1 (axa)^* = t_1 a^* x^* a^* = x x^* a^* = x (ax)^*.$$

Multiplying the previous equality by a from the left side, we obtain $ax = ax(ax)^*$, which gives that $ax = (ax)^*$. Therefore, we get $x = x(ax)^* = xax$.

From $a = axa = (ax)^* a = x^* a^* a$, it follows that $a^* = a^* a x$. So, $Ra^* \subseteq Rx$. Therefore, $Ra^* = Rx$. According to the condition ${}^0x = {}^0a$, we deduce that $xR = aR$ by Lemma 2.8.

(vi) \Rightarrow (i) Clearly, we have $Ra^* \subseteq Rx$, which yields $a = x^* t_2$ for some $t_2 \in R$. So, we have

$$a = x^* t_2 = (xax)^* t_2 = x^* a^* x^* t_2 = x^* a^* a = (ax)^* a.$$

Hence, we get $ax = (ax)^* a x$, which immediately yields $ax = (ax)^*$. Then, $a = (ax)^* a = axa$.

On one hand, we can see that $x = xax = x(ax)^* = x x^* a^*$, which implies $Rx \subseteq Ra^*$, leading to $Rx = Ra^*$, since $Ra^* \subseteq Rx$. On the other hand, it is easy to get $xR = aR$. Finally, by the definition of the core inverse, we claim that $a \in R^\oplus$ and $x = a^\oplus$.

(i) \Rightarrow (vii) and (i) \Rightarrow (viii) can be obtained by Lemma 2.2 and the definition of core inverse.

(vii) \Rightarrow (i) Note that $x \in a\{1, 3\}$ and $x = at_3$ for some $t_3 \in R$. From Lemma 2.2, it follows that $a \in R^\oplus$ and $a^\oplus = a^\# a x = a^\# a a t_3 = a t_3 = x$.

(viii) \Rightarrow (i) This is analogous to the proof of (ii) \Rightarrow (i). Indeed, there exists $r_4 \in R$ such that $a = x t_4$. Then, we have

$$axa = axaaa^\# = a(xax)t_4 a^\# = a(xt_4)a^\# = a^2 a^\# = a,$$

which gives that $x \in a\{1, 3\}$, since $(ax)^* = ax$. Hence, $a \in R^\oplus$ with

$$a^\oplus = a^\#ax = aa^\#x = xt_4a^\#x = xa(xt_4)a^\#x = xa^2a^\#x = xax = x. \quad \square$$

Now, we state a sufficient condition for the reverse order law $(ab)^\oplus = b^\oplus a^\oplus$ to hold, which extends [4, Theorem 3.5].

Theorem 3.2. *Let $a, b \in R^\oplus$, $ab \in R^\#$. If $abb^\oplus = bb^\oplus a$ and $baa^\oplus = aa^\oplus b$, then $ab \in R^\oplus$ and $(ab)^\oplus = b^\oplus a^\oplus$.*

Proof. According to Theorem 3.1 (i) and (ii), it suffices to prove that $abb^\oplus a^\oplus ab = ab$, $b^\oplus a^\oplus R = abR$, and $Rb^\oplus a^\oplus \subseteq R(ab)^*$.

The condition $abb^\oplus = bb^\oplus a$ implies $a^*bb^\oplus = bb^\oplus a^*$. Then, from Lemma 2.5, it follows that $a^\oplus bb^\oplus = bb^\oplus a^\oplus$. Similarly, we obtain $b^\oplus aa^\oplus = aa^\oplus b^\oplus$. Hence, we get $abb^\oplus a^\oplus ab = aa^\oplus abb^\oplus b = ab$.

On one hand, we have

$$\begin{aligned} b^\oplus a^\oplus &= (b^\oplus aa^\oplus)a^\oplus = aa^\oplus b^\oplus a^\oplus = a(a^\oplus bb^\oplus)b^\oplus a^\oplus \\ &= abb^\oplus a^\oplus b^\oplus a^\oplus = ab(b^\oplus a^\oplus)^2, \end{aligned}$$

which gives $b^\oplus a^\oplus R \subseteq abR$. On the other hand, combining $b^\oplus a^\oplus = ab(b^\oplus a^\oplus)^2$ and $abb^\oplus a^\oplus ab = ab$, we get

$$\begin{aligned} ab &= (ab)^\#ab(b^\oplus a^\oplus)(ab)^2 = (ab)^\#abab(b^\oplus a^\oplus)^2(ab)^2 \\ &= (ab(b^\oplus a^\oplus)^2)(ab)^2 = b^\oplus a^\oplus (ab)^2, \end{aligned}$$

which yields $abR \subseteq b^\oplus a^\oplus R$. Thus, we have $b^\oplus a^\oplus R = abR$.

Finally, by the following equalities

$$\begin{aligned} b^\oplus a^\oplus &= b^\oplus (bb^\oplus a^\oplus)aa^\oplus = b^\oplus a^\oplus bb^\oplus aa^\oplus = b^\oplus a^\oplus (aa^\oplus bb^\oplus)^* \\ &= b^\oplus a^\oplus (abb^\oplus a^\oplus)^* = b^\oplus a^\oplus (b^\oplus a^\oplus)^*(ab)^*, \end{aligned}$$

we have $Rb^\oplus a^\oplus \subseteq R(ab)^*$.

The proof is completed. \square

Proposition 3.3. [4, Theorem 3.5] *Let $a, b \in R^\oplus$ with $ab = ba$ and $ab^* = b^*a$. Then $ab \in R^\oplus$ and $(ab)^\oplus = b^\oplus a^\oplus$.*

Proof. From $ab = ba$ and $ab^* = b^*a$, we have $ab^\oplus = b^\oplus a$ and $ba^\oplus = a^\oplus b$ by Lemma 2.5, which imply $abb^\oplus = bb^\oplus a$ and $baa^\oplus = aa^\oplus b$. In addition, we know that $ab \in R^\#$, since $a, b \in R^\#$ with $ab = ba$. Then, $(ab)^\oplus = b^\oplus a^\oplus$ by Theorem 3.2. \square

Remark 3.4. *We have seen that the condition of Proposition 3.3 can imply the condition of Theorem 3.2. But, in general, the condition of Theorem 3.2 does not imply the condition of Proposition 3.3. Indeed, take non-commutative invertible elements $a, b \in R$. Obviously, a and b satisfy the condition of Theorem 3.2. However, $ab \neq ba$.*

Next, we give an equivalent condition which ensures the reverse order law $(ab)^\oplus = b^\oplus a^\oplus$ holds.

Theorem 3.5. *Let $a, b \in R^\oplus$. Then the following are equivalent:*

- (i) $ab \in R^\oplus$ and $(ab)^\oplus = b^\oplus a^\oplus$;
- (ii) $ab \in R^\#$, $ab^\oplus R \subseteq b^\oplus a^\oplus R$, and $b^* a^\#(1 - (abb^\oplus a^\oplus)^*)a = 0$.

Proof. (i) \Rightarrow (ii) Obviously, $ab \in R^\#$. Note that

$$ab^\oplus = ab(b^\oplus)^2 = b^\oplus a^\oplus (ab)^2 (b^\oplus)^2,$$

which implies $ab^\oplus R \subseteq b^\oplus a^\oplus R$. By Lemma 2.2, we have $a^\oplus a = a^\# aa^{(1,3)} a = a^\# a$, together with $b^* = b^* b b^\oplus$, we obtain

$$\begin{aligned} b^* a a^\# &= b^* b b^\oplus a^\oplus a^2 a^\# = b^* b b^\oplus (a^\oplus a) = b^* b (ab)^\oplus a = b^* b (ab)^\oplus ab (ab)^\oplus a \\ &= b^* b b^\oplus a^\oplus ab b^\oplus a^\oplus a = (b^* b b^\oplus) (a^\oplus a) b b^\oplus a^\oplus a = b^* a^\# ab b^\oplus a^\oplus a \\ &= b^* a^\# (ab b^\oplus a^\oplus)^* a. \end{aligned}$$

Therefore, $b^* a^\#(1 - (ab b^\oplus a^\oplus)^*)a = 0$.

(ii) \Rightarrow (i) Observe that $a^\oplus = a^\oplus aa^{(1,3)}$ and $a^\# aa^\oplus = a^\oplus$ by Lemma 2.2. Then, applying the condition $b^* a^\# a = b^* a^\# (ab b^\oplus a^\oplus)^* a$, we obtain

$$\begin{aligned} b^* a^\oplus &= (b^* a^\# a) a^\oplus = b^* a^\# (ab b^\oplus a^\oplus)^* a a^\oplus = b^* a^\# (a^\oplus)^* b b^\oplus a^* a a^\oplus \\ &= b^* a^\# (a^\oplus)^* b b^\oplus a^* = b^* a^\# (a^\oplus aa^{(1,3)})^* b b^\oplus a^* \\ &= b^* a^\# aa^{(1,3)} (a^\oplus)^* b b^\oplus a^* \\ &= b^* a^\oplus (a^\oplus)^* b b^\oplus a^*, \end{aligned}$$

which yields $(a^\oplus)^* b = ab b^\oplus a^\oplus (a^\oplus)^* b$. By the previous equality, we get

$$\begin{aligned} (a^\oplus)^* (b^\oplus)^* &= (a^\oplus)^* (b^\oplus b b^\oplus)^* = ((a^\oplus)^* b) b^\oplus (b^\oplus)^* = ab b^\oplus a^\oplus (a^\oplus)^* b b^\oplus (b^\oplus)^* \\ &= ab b^\oplus a^\oplus (a^\oplus)^* (b^\oplus)^* = ab b^\oplus a^\oplus (b^\oplus a^\oplus)^*, \end{aligned}$$

which gives $b^\oplus a^\oplus = b^\oplus a^\oplus (ab b^\oplus a^\oplus)^*$. Then, $ab b^\oplus a^\oplus = ab b^\oplus a^\oplus (ab b^\oplus a^\oplus)^*$. So, we have $ab b^\oplus a^\oplus = (ab b^\oplus a^\oplus)^*$. This immediately yields $b^\oplus a^\oplus = b^\oplus a^\oplus ab b^\oplus a^\oplus$.

The condition $ab^\oplus R \subseteq b^\oplus a^\oplus R$ ensures that $ab^\oplus = b^\oplus a^\oplus r$ for some $r \in R$. Hence, we get $ab = (ab^\oplus) b^2 = b^\oplus a^\oplus r b^2$, which gives $abR \subseteq b^\oplus a^\oplus R$. From Theorem 3.1, it follows that $ab \in R^\oplus$ and $(ab)^\oplus = b^\oplus a^\oplus$. \square

In order to simplify the proof of the following theorems, we present a useful lemma.

Lemma 3.6. *Let $a, b, ab \in R^\oplus$ with $(ab)^\oplus = b^\oplus a^\oplus$. Then*

- (i) $ab = b b^\oplus ab = b^\oplus b ab$;
- (ii) $abR \subseteq baR$;
- (iii) $ab(ab)^\oplus b b^\oplus = b b^\oplus ab(ab)^\oplus = ab(ab)^\oplus$;
- (iv) $b b^\oplus a^\oplus \in ab b^\oplus \{3, 6\}$.

Proof. (i) Suppose that $(ab)^\oplus = b^\oplus a^\oplus$. Then, we have

$$ab = b^\oplus a^\oplus (ab)^2 = bb^\oplus (b^\oplus a^\oplus (ab)^2) = bb^\oplus ab,$$

and

$$ab = b^\oplus a^\oplus (ab)^2 = b^\oplus b (b^\oplus a^\oplus (ab)^2) = b^\oplus bab.$$

(ii) From (i), it follows that

$$ab = bb^\oplus ab = bb^\oplus a^\oplus a^2 b = b(ab)^\oplus a^2 b = bab((ab)^\oplus)^2 a^2 b.$$

Hence, $abR \subseteq baR$.

(iii) Clearly, we have

$$\begin{aligned} ab(ab)^\oplus bb^\oplus &= (ab(ab)^\oplus)^*(bb^\oplus)^* = (bb^\oplus ab(ab)^\oplus)^* = (bb^\oplus abb^\oplus a^\oplus)^* \\ &= (abb^\oplus a^\oplus)^* = ab(ab)^\oplus, \end{aligned}$$

which implies that $ab(ab)^\oplus bb^\oplus = (ab(ab)^\oplus bb^\oplus)^* = bb^\oplus ab(ab)^\oplus$.

(iv) Since $(ab)^\oplus = b^\oplus a^\oplus$, we have $abb^\oplus bb^\oplus a^\oplus = abb^\oplus a^\oplus = ab(ab)^\oplus$, which gives $bb^\oplus a^\oplus \in abb^\oplus\{3\}$. Also, we have

$$\begin{aligned} bb^\oplus a^\oplus (abb^\oplus)^2 &= b(b^\oplus a^\oplus abb^\oplus a^\oplus) a^2 bb^\oplus = bb^\oplus (a^\oplus a^2) bb^\oplus = (bb^\oplus ab) b^\oplus \\ &= abb^\oplus. \end{aligned}$$

Thus, $bb^\oplus a^\oplus \in abb^\oplus\{3, 6\}$. □

Deng [10] and Mosić [23] studied the reverse order law $(ab)^\# = b^\# a^\#$ for the group inverse under the condition $ba = a^2$. Motivated by this, we will consider the one-sided reverse law for the core inverse under the same condition, using Theorem 3.5.

Theorem 3.7. *Let $a, b \in R^\oplus$ with $ba = a^2$. Then,*

- (i) $ab \in R^\oplus$ and $(ab)^\oplus = b^\oplus a^\oplus$.
- (ii) $abb^\oplus \in R^\oplus$ and $(abb^\oplus)^\oplus = bb^\oplus a^\oplus$.

Proof. (i) First, we prove $ab \in R^\#$. Note that $a \in R^\#$ and $ba = a^2$. Then, we have

$$ab = aa^2 a^2 (a^\#)^4 b = ababa(a^\#)^4 b = (ab)^2 (a^\#)^3 b$$

and

$$ab = a^2 a^\# b = baa^\# b = b(a^\#)^3 a a^2 b = b(a^\#)^3 abab = b(a^\#)^3 (ab)^2,$$

which imply $ab \in R^\#$ by Lemma 2.3.

Next, we see that

$$a = a^2 a^\# = baa^\# = b^\oplus b^2 a a^\# = b^\oplus b (ba) a^\# = b^\oplus b a^2 a^\# = b^\oplus b a = b^\oplus a^2.$$

Hence, $ab^\oplus = b^\oplus a^2 b^\oplus = b^\oplus a^\oplus a^3 b^\oplus$, which yields $ab^\oplus R \subseteq b^\oplus a^\oplus R$. Note that $a = b^\oplus b a$, then we deduce that $a^\oplus = a(a^\oplus)^2 = b^\oplus b a (a^\oplus)^2 = b^\oplus b a^\oplus$. In addition, the assumption $ba = a^2$ ensures that $aa^\oplus = a^2(a^\oplus)^2 = ba(a^\oplus)^2 = ba^\oplus$. Therefore,

$$abb^\oplus a^\oplus = abb^\oplus b^\oplus b a^\oplus = ab^\oplus b a^\oplus = a(b^\oplus a) a^\oplus = a(b^\oplus a^2) a^\oplus = a a^\oplus a^\oplus = aa^\oplus,$$

which implies $(abb^\oplus a^\oplus)^* = abb^\oplus a^\oplus$. So, we conclude that

$$b^* a^\# (1 - (abb^\oplus a^\oplus)^*) a = b^* a^\# a - b^* a^\# (abb^\oplus a^\oplus) a = b^* a^\# a - b^* a^\# a a^\oplus a = 0.$$

By Theorem 3.5, we claim that $ab \in R^\oplus$ and $(ab)^\oplus = b^\oplus a^\oplus$.

(ii) By (i), Lemma 3.6(iv), and Lemma 2.1, we only need to prove $bb^\oplus a^\oplus \in abb^\oplus \{7\}$.

In the proof of (i), we obtain $a^\oplus = b^\oplus b a^\oplus$ and $ba^\oplus = aa^\oplus$ which imply that $bb^\oplus a^\oplus = bb^\oplus b^\oplus (ba^\oplus) = b^\oplus aa^\oplus$. Hence,

$$bb^\oplus a^\oplus (abb^\oplus)^2 = b^\oplus (aa^\oplus a) (bb^\oplus ab) b^\oplus = b^\oplus a ab b^\oplus = (b^\oplus bab) b^\oplus = abb^\oplus. \quad \square$$

Using Lemma 3.6 and Theorem 3.1, we deduce the following result.

Theorem 3.8. *Let $a, b, ab \in R^\oplus$. Then the following are equivalent:*

- (i) $(ab)^\oplus = b^\oplus a^\oplus$;
- (ii) $b(ab)^\oplus = bb^\oplus a^\oplus$ and $abb^\oplus = b^\oplus b ab b^\oplus$.

Proof. (i) \Rightarrow (ii) can be obtained by Lemma 3.6(i).

(ii) \Rightarrow (i) Since $b(ab)^\oplus = bb^\oplus a^\oplus$, we have

$$a(bb^\oplus a^\oplus)ab = ab(ab)^\oplus ab = ab \quad \text{and} \quad abb^\oplus a^\oplus = ab(ab)^\oplus,$$

which implies $b^\oplus a^\oplus \in ab\{1, 3\}$. In addition, observe that

$$\begin{aligned} b^\oplus a^\oplus &= b^\oplus (bb^\oplus a^\oplus) = b^\oplus b (ab)^\oplus = b^\oplus b ab ((ab)^\oplus)^2 \\ &= (b^\oplus b ab b^\oplus) b ((ab)^\oplus)^2 = abb^\oplus b ((ab)^\oplus)^2, \end{aligned}$$

which yields $b^\oplus a^\oplus R \subseteq abR$. Hence, by Theorem 3.1 (i) and (vii), we obtain $(ab)^\oplus = b^\oplus a^\oplus$. \square

In the following theorem, we investigate the one-sided reverse order law for the core inverse under the condition $Rb^*a \subseteq Rab^*$.

Theorem 3.9. *Let $a, b \in R^\oplus$ with $Rb^*a \subseteq Rab^*$. Then the following are equivalent:*

- (i) $ab \in R^\oplus$ and $(ab)^\oplus = b^\oplus a^\oplus$;
- (ii) $ab \in R^\#$, $abR \subseteq baR$, $b^\oplus a^\oplus R \subseteq abR$, and $aa^\oplus bb^\oplus = bb^\oplus aa^\oplus$;
- (iii) $ab, abb^\oplus \in R^\oplus$ with $(abb^\oplus)^\oplus = bb^\oplus a^\oplus$ and $(ab)^\oplus = b^\oplus (abb^\oplus)^\oplus$.

Proof. (i) \Rightarrow (ii) Obviously, $ab \in R^\#$. From $(ab)^\oplus = b^\oplus a^\oplus$ and Lemma 3.6, it follows that $abR \subseteq baR$. The equality $b^\oplus a = b^\oplus a^\oplus a^2 = ab((ab)^\oplus)^2 a^2$ implies $b^\oplus aR \subseteq abR$.

Next, our aim is to prove $aa^\oplus bb^\oplus = bb^\oplus aa^\oplus$. The assumption $Rb^*a \subseteq Rab^*$ guarantees $b^*a = w_1ab^*$ for some $w_1 \in R$. Then, we have

$$b^*a = w_1a(bb^\oplus b)^* = w_1ab^*bb^\oplus = b^*abb^\oplus.$$

Note that $ab = bb^\oplus ab$ by Lemma 3.6(i), we obtain

$$\begin{aligned} abb^\oplus a^\oplus &= bb^\oplus abb^\oplus a^\oplus = (bb^\oplus)^* abb^\oplus a^\oplus = (b^\oplus)^*(b^*abb^\oplus)a^\oplus \\ &= (b^\oplus)^*b^*aa^\oplus = bb^\oplus aa^\oplus. \end{aligned}$$

Therefore, the following equalities hold:

$$abb^\oplus a^\oplus = (abb^\oplus a^\oplus)^* = (bb^\oplus aa^\oplus)^* = aa^\oplus bb^\oplus.$$

So, we have $aa^\oplus bb^\oplus = bb^\oplus aa^\oplus$.

(ii) \Rightarrow (i) Since $abR \subseteq baR$, there exists $w_2 \in R$ such that $ab = baw_2$. Therefore, we obtain $ab = bb^\oplus(baw_2) = bb^\oplus ab$. Then, the equality $abb^\oplus a^\oplus = bb^\oplus aa^\oplus$ can be obtained in a similar way as in the proof of (i) \Rightarrow (ii). Hence, we deduce that

$$\begin{aligned} (b^\oplus a^\oplus)^*(ab)^*ab &= (abb^\oplus a^\oplus)^*ab = (bb^\oplus aa^\oplus)^*ab = aa^\oplus bb^\oplus ab \\ &= bb^\oplus aa^\oplus ab = bb^\oplus ab = ab, \end{aligned}$$

which implies $b^\oplus a^\oplus \in ab\{1, 3\}$ by Lemma 2.4.

From $b^\oplus aR \subseteq abR$, we know that $b^\oplus a = abw_3$ for some $w_3 \in R$. Then, we obtain $b^\oplus a^\oplus = b^\oplus a(a^\oplus)^2 = abw_3(a^\oplus)^2$, which gives that $b^\oplus a^\oplus R \subseteq abR$. Hence, $ab \in R^\oplus$ with $(ab)^\oplus = b^\oplus a^\oplus$ by Theorem 3.1.

(i) \Rightarrow (iii) According to the condition (i) and Lemma 3.6 (iv), we get $bb^\oplus a^\oplus \in abb^\oplus\{3, 6\}$. In addition, from the proof of (i) \Rightarrow (ii), we see that $abb^\oplus a^\oplus = bb^\oplus aa^\oplus = aa^\oplus bb^\oplus$. Then, we obtain

$$\begin{aligned} abb^\oplus(bb^\oplus a^\oplus)^2 &= (abb^\oplus a^\oplus)bb^\oplus a^\oplus = aa^\oplus bb^\oplus bb^\oplus a^\oplus = aa^\oplus bb^\oplus a^\oplus \\ &= bb^\oplus aa^\oplus a^\oplus = bb^\oplus a^\oplus. \end{aligned}$$

By Lemma 2.1, it follows that $abb^\oplus \in R^\oplus$ and $(abb^\oplus)^\oplus = bb^\oplus a^\oplus$. Then, it is easy to see that $(ab)^\oplus = b^\oplus(bb^\oplus a^\oplus) = b^\oplus(abb^\oplus)^\oplus$.

(iii) \Rightarrow (i) Suppose that $(abb^\oplus)^\oplus = bb^\oplus a^\oplus$ and $(ab)^\oplus = b^\oplus(abb^\oplus)^\oplus$. Then, we deduce

$$(ab)^\oplus = b^\oplus(abb^\oplus)^\oplus = b^\oplus bb^\oplus a^\oplus = b^\oplus a^\oplus. \quad \square$$

Remark 3.10. (1) Note that Corollary 3.3 can also be obtained by Theorem 3.9. Indeed, we have $ab^\oplus = b^\oplus a$, $ba^\oplus = a^\oplus b$, and $a^\oplus b^\oplus = b^\oplus a^\oplus$ by Lemma 2.5. Then, $aa^\oplus bb^\oplus = bb^\oplus aa^\oplus$ and $b^\oplus a = ab^\oplus = ab(b^\oplus)^2$, which gives $b^\oplus aR \subseteq abR$. Hence, we have $ab \in R^\oplus$ and $(ab)^\oplus = b^\oplus a^\oplus$ by Theorem 3.9.

(2) In general, any item of (i)–(iii) in Theorem 3.9 does not imply $Rb^*a \subseteq Rab^*$. For example, let \mathbb{Z}_2 be the ring of integers modulo 2, take $R = M_2(\mathbb{Z}_2)$ with the transpose of matrices as involution. Setting $a = b = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \in R$. Then $a^2 = a$, which implies that $a \in R^\#$ with $a^\# = a$. Note that $a = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} a^* a \in Ra^*a$, then $a \in R^{\{1,3\}}$. Hence, $a \in R^\oplus$ with $a^\oplus = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$. Then, it is easy to check that such a and b satisfy any item of (i)–(iii) in Theorem 3.9. However, $ab^* = 0$ and $b^*a \neq 0$, so $Rb^*a \not\subseteq Rab^*$.

(3) If we replace $Rb^*a \subseteq Rab^*$ with $ba = a^2$ in Theorem 3.9 then the items (i), (ii), and (iii) hold. Indeed, by Theorem 3.7 and Lemma 3.6, we only need to show that $b^\oplus a R \subseteq abR$ and $aa^\oplus bb^\oplus = bb^\oplus aa^\oplus$. Since $(ab)^\oplus = b^\oplus a^\oplus$ by Theorem 3.7, then $b^\oplus a R = (b^\oplus a^\oplus) a^2 R = (ab)^\oplus a^2 R = ab((ab)^\oplus)^2 a^2 R \subseteq abR$. In addition, from the proof of Theorem 3.7, we see that $a^\oplus = b^\oplus b a^\oplus$ and $aa^\oplus = b a^\oplus$, which gives $b(b^\oplus a a^\oplus) = b a^\oplus = a a^\oplus$ is Hermitian. Thus, we get $bb^\oplus a a^\oplus = (bb^\oplus a a^\oplus)^* = (a a^\oplus)^* (b b^\oplus)^* = a a^\oplus b b^\oplus$.

Next, we continue to consider the one-sided reverse law for the core inverse under certain conditions.

Theorem 3.11. *Let $a, b \in R^\oplus$ with $Rb \subseteq Rab$ (or $aR \subseteq abR$). If a is EP, then the following are equivalent:*

- (i) $ab \in R^\oplus$ and $(ab)^\oplus = b^\oplus a^\oplus$;
- (ii) $(a^\oplus)^* b \in R^\oplus$ and $((a^\oplus)^* b)^\oplus = b^\oplus a^*$.

Proof. (i) \Rightarrow (ii) Since $(ab)^\oplus = b^\oplus a^\oplus$, then we have that

$$(a^\oplus)^* b b^\oplus a^* = (a^\oplus)^* (b b^\oplus)^* a^* = (a b b^\oplus a^\oplus)^* = a b b^\oplus a^\oplus$$

is Hermitian, i.e., $b^\oplus a^* \in (a^\oplus)^* b \{3\}$.

Note that a is EP, then $a^\oplus a = a a^\oplus$ is Hermitian by Lemma 2.7. So, $a^* = (a(a^\oplus a))^* = a^\oplus a a^*$. Thus, we have

$$\begin{aligned} (a^\oplus)^* b (b^\oplus a^*)^2 &= ((a^\oplus)^* (b b^\oplus)^* a^*) b^\oplus a^* = (a b b^\oplus a^\oplus) b^\oplus a^* \\ &= (a b b^\oplus a^\oplus b^\oplus a^\oplus) a a^* = b^\oplus a^\oplus a a^* = b^\oplus a^*. \end{aligned}$$

If $Rb \subseteq Rab$, then $b = xab$ for some $x \in R$. Therefore, we can deduce that

$$\begin{aligned} b^\oplus a^* ((a^\oplus)^* b)^2 &= b^\oplus a^* (a^\oplus)^* b (a^\oplus)^* b = b^\oplus (a^\oplus a)^* b (a^\oplus)^* b \\ &= b^\oplus a^\oplus a b (a^\oplus)^* (b b^\oplus)^* b = b^\oplus a^\oplus a b (b b^\oplus a^\oplus)^* b \\ &= b^\oplus a^\oplus a b (x a b b^\oplus a^\oplus)^* b = (b^\oplus a^\oplus a b a b) b^\oplus a^\oplus x^* b \\ &= a b b^\oplus a^\oplus x^* b = (a b b^\oplus a^\oplus)^* x^* b \\ &= (x a b b^\oplus a^\oplus)^* b = (b b^\oplus a^\oplus)^* b \\ &= (a^\oplus)^* (b b^\oplus)^* b = (a^\oplus)^* b. \end{aligned}$$

If $aR \subseteq abR$, then $a = aby$ for some $y \in R$. Thus, we get

$$\begin{aligned} b^\oplus a^* ((a^\oplus)^* b)^2 &= b^\oplus a^\oplus a b (a^\oplus)^* b = b^\oplus a^\oplus a b (a^\oplus a a^\oplus)^* b \\ &= b^\oplus a^\oplus a b a a^\oplus (a^\oplus)^* b = (b^\oplus a^\oplus a b a b) y a^\oplus (a^\oplus)^* b \\ &= a b y a^\oplus (a^\oplus)^* b = a a^\oplus (a^\oplus)^* b = (a^\oplus)^* b. \end{aligned}$$

Finally, from Lemma 2.1, it follows that $(a^\oplus)^*b \in R^\oplus$ and $((a^\oplus)^*b)^\oplus = b^\oplus a^*$.

(ii) \Rightarrow (i) By Lemma 2.7, we have $a^\# = a^\oplus$. Let $c = (a^\oplus)^*$, then $c = (a^\#)^* = (a^*)^\#$, which implies that $a^* = c^\#$. Note that c is EP. Thus, $a^* = c^\oplus$. According to the condition (ii), we obtain that $cb \in R^\oplus$ and $(cb)^\oplus = b^\oplus c^\oplus$.

If $Rb \subseteq Rab$, then $Rb \subseteq Rab = Raa^\oplus ab = Ra(a^\oplus a)^*b = Raa^*(a^\oplus)^*b \subseteq R(a^\oplus)^*b = Rcb$.

If $aR \subseteq abR$, then $a = aby$ for some $y \in R$. Since $((a^\oplus)^*b)^\oplus = b^\oplus a^*$, then $b^\oplus a^*R = (a^\oplus)^*bR$, which yields $(a^\oplus)^*b = b^\oplus a^*z$ for some $z \in R$. Note that $a^* = a^\oplus aa^*$. Then, we obtain

$$\begin{aligned} (a^\oplus)^* &= (aa^\oplus a^\oplus)^* = (a^\oplus)^*(a^\oplus)^*a^* = (a^\oplus)^*(a^\oplus)^*a^\oplus aa^* \\ &= (a^\oplus)^*(a^\oplus)^*a^\oplus aby a^* = (a^\oplus)^*((a^\oplus)^*(a^\oplus a)^*)by a^* \\ &= (a^\oplus)^*((a^\oplus)^*b)ya^* = (a^\oplus)^*b^\oplus a^*zy a^* \\ &= (a^\oplus)^*b(b^\oplus)^2 a^*zy a^*, \end{aligned}$$

which gives $(a^\oplus)^*R \subseteq (a^\oplus)^*bR$, i.e., $cR \subseteq cbR$.

Now, replacing a with c in the proof of (i) \Rightarrow (ii), we can obtain that (i) holds. \square

If we replace the assumption $Rb \subseteq Rab$ (or $aR \subseteq abR$) of Theorem 3.11 by $Rb \subseteq Rab$ and $aR \subseteq abR$, then we obtain the following result.

Theorem 3.12. *Let $a, b \in R^\oplus$ with $Rb \subseteq Rab$ and $aR \subseteq abR$. If a is EP, then the following are equivalent:*

- (i) $ab \in R^\oplus$ and $(ab)^\oplus = b^\oplus a^\oplus$;
- (ii) $ab, abb^\oplus \in R^\oplus$ with $(ab)^\oplus = b^\oplus(abb^\oplus)^\oplus$ and $(abb^\oplus)^\oplus = bb^\oplus a^\oplus$.

Proof. (i) \Rightarrow (ii) In order to prove $(abb^\oplus)^\oplus = bb^\oplus a^\oplus$, it suffices to prove that $abb^\oplus(bb^\oplus a^\oplus)^2 = bb^\oplus a^\oplus$ by Lemma 2.6 (iv).

Since $Rb \subseteq Rab$ and $aR \subseteq abR$, there exist t_1, t_2 such that $b = t_1 ab$ and $a = abt_2$. Note that $ab = bb^\oplus ab$ and $aa^\oplus = a^\oplus a$. Thus, we get

$$\begin{aligned} abb^\oplus(bb^\oplus a^\oplus)^2 &= abb^\oplus a^\oplus bb^\oplus a^\oplus = bb^\oplus abb^\oplus a^\oplus aa^\oplus bb^\oplus a^\oplus \\ &= bb^\oplus(abb^\oplus a^\oplus ab)t_2 a^\oplus bb^\oplus a^\oplus = t_1 abb^\oplus(abt_2)a^\oplus bb^\oplus a^\oplus \\ &= t_1 abb^\oplus aa^\oplus bb^\oplus a^\oplus = t_1(abb^\oplus a^\oplus ab)b^\oplus a^\oplus \\ &= (t_1 ab)b^\oplus a^\oplus = bb^\oplus a^\oplus. \end{aligned}$$

(ii) \Rightarrow (i) It is obvious. \square

Now, necessary and sufficient conditions of the two-sided reverse laws for the core inverse are stated as follows.

Theorem 3.13. *Let $a, b \in R^\oplus$. Then the following are equivalent:*

- (i) $ab, ba \in R^\oplus$ with $(ab)^\oplus = b^\oplus a^\oplus$, $(ba)^\oplus = a^\oplus b^\oplus$;
- (ii) $abb^\oplus, baa^\oplus \in R^\oplus$ with $(abb^\oplus)^\oplus = bb^\oplus a^\oplus$, $(baa^\oplus)^\oplus = aa^\oplus b^\oplus$, $abR = (ab)^2R$, and $baR = (ba)^2R$.

Proof. (i) \Rightarrow (ii) Obviously, $abR = (ab)^2R$ and $baR = (ba)^2R$. By symmetry, we only need to prove $abb^\oplus \in R^\oplus$ with $(abb^\oplus)^\oplus = bb^\oplus a^\oplus$. Note that $bb^\oplus a^\oplus \in abb^\oplus\{3, 6\}$. Applying the hypothesis $(ba)^\oplus = a^\oplus b^\oplus$ and Lemma 3.6(ii), we have $baR = abR$, i.e., $ba = abu$ for some $u \in R$. Then,

$$\begin{aligned} abb^\oplus(bb^\oplus a^\oplus)^2 &= abb^\oplus a^\oplus bb^\oplus a^\oplus = ab(ab)^\oplus b(ab)^\oplus = ab(ab)^\oplus bab((ab)^\oplus)^2 \\ &= (ab(ab)^\oplus ab)ub((ab)^\oplus)^2 = abub((ab)^\oplus)^2 = bab((ab)^\oplus)^2 \\ &= b(ab)^\oplus = bb^\oplus a^\oplus. \end{aligned}$$

Using Lemma 2.1, we get $abb^\oplus \in R^\oplus$ with $(abb^\oplus)^\oplus = bb^\oplus a^\oplus$.

(ii) \Rightarrow (i) Since $(abb^\oplus)^\oplus = bb^\oplus a^\oplus$, we claim that

$$b^\oplus a^\oplus abb^\oplus a^\oplus = b^\oplus (bb^\oplus a^\oplus abb^\oplus bb^\oplus a^\oplus) = b^\oplus bb^\oplus a^\oplus = b^\oplus a^\oplus$$

and

$$abb^\oplus a^\oplus = abb^\oplus bb^\oplus a^\oplus = abb^\oplus (abb^\oplus)^\oplus,$$

which imply $b^\oplus a^\oplus \in ab\{2, 3\}$. In addition, observe that

$$abb^\oplus = bb^\oplus a^\oplus (abb^\oplus)^2 = b^\oplus b (bb^\oplus a^\oplus (abb^\oplus)^2) = b^\oplus babb^\oplus.$$

Multiplying the previous equality by b from the right side, we get $ab = b^\oplus bab$. Symmetrically, $ba = a^\oplus aba$. Hence, we have $ab = b^\oplus (ba)b = b^\oplus a^\oplus (ab)^2$, which gives $abR \subseteq b^\oplus a^\oplus R$ and $Rab = R(ab)^2$. Note that $abR = (ab)^2R$, then $ab \in R^\#$ by Lemma 2.3. Finally, from Theorem 3.1, it follows that $ab \in R^\oplus$ with $(ab)^\oplus = b^\oplus a^\oplus$. The statement for ba can be obtained by symmetry. \square

Recall that a ring R (with unit 1) is a Dedekind-finite ring if $ab = 1$ is sufficient for $ba = 1$. Let $p = q = 1 \in R$ in [27, Theorem 1]. Then we can see the fact: If $a \in R$ is regular, $a^- \in a\{1\}$, then $u = a^2 a^- + 1 - aa^-$ is right (resp. left) invertible if and only if $aR = a^2 R$ (resp. $Ra = Ra^2$). Form the previous fact, it immediately yields that $aR = a^2 R$ if and only if $Ra = Ra^2$, where R is a Dedekind-finite ring, and a is regular in R . Thus, by Theorem 3.13, we get

Corollary 3.14. *Let R be a Dedekind-finite ring and $a, b \in R^\oplus$. Then the following are equivalent:*

- (i) $ab, ba \in R^\oplus$ with $(ab)^\oplus = b^\oplus a^\oplus$, $(ba)^\oplus = a^\oplus b^\oplus$;
- (ii) $abb^\oplus, baa^\oplus \in R^\oplus$ with $(abb^\oplus)^\oplus = bb^\oplus a^\oplus$, $(baa^\oplus)^\oplus = aa^\oplus b^\oplus$.

Proof. According to the proof of Theorem 3.13, we only need to prove that ab and ba are regular. Suppose that $(abb^\oplus)^\oplus = bb^\oplus a^\oplus$, then we have

$$abb^\oplus = bb^\oplus a^\oplus (abb^\oplus)^2 = bb^\oplus (bb^\oplus a^\oplus (abb^\oplus)^2) = bb^\oplus abb^\oplus,$$

which yields

$$ab = (abb^\oplus)b = bb^\oplus abb^\oplus b = bb^\oplus ab = (bb^\oplus a^\oplus) a^2 b = abb^\oplus (bb^\oplus a^\oplus)^2 aab.$$

Therefore, ab is regular. Similarly, by the condition $(baa^\oplus)^\oplus = aa^\oplus b^\oplus$, we can obtain that ba is regular. \square

4 Mixed-type Reverse Order Laws

In this section, we will consider necessary and sufficient conditions for the mixed-type reverse laws: $(ab)^\# = b^\oplus(abb^\oplus)^\oplus$, $a^\oplus = b(ab)^\#$ and $(ab)^\# = b^\oplus a^\oplus$ to hold in rings.

Theorem 4.1. *Let $b \in R^\oplus$, $abb^\oplus \in R^\oplus$. Then the following are equivalent:*

- (i) $ab \in R^\oplus$ and $(ab)^\oplus = (ab)^\# = b^\oplus(abb^\oplus)^\oplus$;
- (ii) $b^\oplus(abb^\oplus)^\oplus \in ab\{5\}$;
- (iii) $b^\oplus bab = ab = bb^\oplus ab$ and $ba(abb^\oplus)^\oplus = (abb^\oplus)^\oplus ab$;
- (iv) $ab \in R^\#$ and $(abb^\oplus)^\oplus = b(ab)^\#$.

Proof. (i) \Rightarrow (ii) It is obvious.

(ii) \Rightarrow (iii) Note that $abb^\oplus(abb^\oplus)^\oplus ab = (abb^\oplus(abb^\oplus)^\oplus abb^\oplus)b = abb^\oplus b = ab$. Also, since $b^\oplus(abb^\oplus)^\oplus \in ab\{5\}$, we have $ab = b^\oplus(abb^\oplus)^\oplus(ab)^2$. From the previous equality, we get

$$ab = bb^\oplus(b^\oplus(abb^\oplus)^\oplus(ab)^2) = bb^\oplus ab$$

and

$$ab = b^\oplus b(b^\oplus(abb^\oplus)^\oplus(ab)^2) = b^\oplus bab.$$

Thus, we obtain $abb^\oplus = bb^\oplus abb^\oplus$. From the definition of the core inverse, it follows that $(abb^\oplus)^\oplus R = abb^\oplus R$, i.e., $(abb^\oplus)^\oplus = abb^\oplus u$ for some $u \in R$. Hence, we deduce

$$(abb^\oplus)^\oplus = bb^\oplus abb^\oplus u = bb^\oplus(abb^\oplus)^\oplus.$$

Thus, the following equations hold:

$$ba(abb^\oplus)^\oplus = babb^\oplus(abb^\oplus)^\oplus = bb^\oplus(abb^\oplus)^\oplus ab = (abb^\oplus)^\oplus ab.$$

(iii) \Rightarrow (iv) Clearly, $b^\oplus(abb^\oplus)^\oplus \in ab\{1, 2\}$.

Now, we prove $b^\oplus(abb^\oplus)^\oplus \in ab\{5\}$. Since $ab = bb^\oplus ab$, from the proof of (ii) \Rightarrow (iii), we can get $(abb^\oplus)^\oplus = bb^\oplus(abb^\oplus)^\oplus$. Then,

$$(ab)b^\oplus(abb^\oplus)^\oplus = b^\oplus babb^\oplus(abb^\oplus)^\oplus = b^\oplus ba(abb^\oplus)^\oplus = b^\oplus(abb^\oplus)^\oplus(ab),$$

which implies that $(ab)^\# = b^\oplus(abb^\oplus)^\oplus$. Thus, we get

$$(abb^\oplus)^\oplus = abb^\oplus((abb^\oplus)^\oplus)^2 = bb^\oplus abb^\oplus((abb^\oplus)^\oplus)^2 = bb^\oplus(abb^\oplus)^\oplus = b(ab)^\#.$$

(iv) \Rightarrow (i) Since $(abb^\oplus)^\oplus = b(ab)^\#$, we have

$$\begin{aligned} ab &= abb^\oplus b = (abb^\oplus)^\oplus(abb^\oplus)^2 b = b(ab)^\#(abb^\oplus)^2 b \\ &= b^\oplus bb(ab)^\#(abb^\oplus)^2 b = b^\oplus b(abb^\oplus)^\oplus(abb^\oplus)^2 b \\ &= b^\oplus babb^\oplus b = b^\oplus bab. \end{aligned}$$

Then, we get

$$(ab)^\# = ab((ab)^\#)^2 = b^\oplus bab((ab)^\#)^2 = b^\oplus b(ab)^\# = b^\oplus(abb^\oplus)^\oplus.$$

Our next aim is to prove $(ab)^\oplus = b^\oplus(abb^\oplus)^\oplus$. Obviously, $b^\oplus(abb^\oplus)^\oplus \in ab\{3\}$. Note that

$$b^\oplus(abb^\oplus)^\oplus(ab)^2 = (ab)^\#(ab)^2 = ab$$

and

$$ab(b^\oplus(abb^\oplus)^\oplus)^2 = ab((ab)^\#)^2 = (ab)^\# = b^\oplus(abb^\oplus)^\oplus,$$

which imply $ab \in R^\oplus$ and $(ab)^\oplus = b^\oplus(abb^\oplus)^\oplus$ by Lemma 2.1. \square

Remark 4.2. Any item of (i)–(iv) in Theorem 4.1 can imply that ab is EP. Indeed, this can be obtained by Lemma 2.7.

Next, characterizations for the equality $a^\oplus = b(ab)^\#$ are presented. In particular, we will see that such equivalent conditions also ensure that ab is EP.

Theorem 4.3. Let $a, b \in R^\oplus$, $ab \in R^\#$. Then the following are equivalent:

- (i) $a^\oplus = b(ab)^\#$;
- (ii) $a^\oplus ab = baa^\oplus$ and $aR \subseteq abR$;
- (iii) $ab \in R^\oplus$ with $(ab)^\# = (ab)^\oplus = b^\oplus a^\oplus$, and $aR \subseteq bR$;
- (iv) $abb^\oplus \in R^\oplus$ with $(abb^\oplus)^\oplus = bb^\oplus a^\oplus$, $(ab)^\# = b^\oplus(abb^\oplus)^\oplus$, and $aR \subseteq bR$;
- (v) $abb^\oplus \in R^\oplus$ with $(abb^\oplus)^\oplus = bb^\oplus a^\oplus = b(ab)^\#$, and $aR \subseteq bR$.

Proof. (i) \Rightarrow (ii) Suppose that $a^\oplus = b(ab)^\#$, then

$$a^\oplus ab = b(ab)^\# ab = ba(b(ab)^\#) = baa^\oplus.$$

Also, we have $a = aa^\oplus a = ab(ab)^\# a$, which implies $aR \subseteq abR$.

(ii) \Rightarrow (iii) There exists $r_1 \in R$ such that $a = abr_1$, since $aR \subseteq abR$. Then

$$a = a^\oplus a^2 = (a^\oplus ab)r_1 a = baa^\oplus r_1 a.$$

Let $r_2 = aa^\oplus r_1 a$. Then, $a = br_2$, which gives $aR \subseteq bR$.

Since

$$b^\oplus(a^\oplus ab) = b^\oplus baa^\oplus = b^\oplus bbr_2 a^\oplus = br_2 a^\oplus = aa^\oplus$$

and

$$abb^\oplus a^\oplus = abb^\oplus a(a^\oplus)^2 = abb^\oplus br_2(a^\oplus)^2 = abr_2(a^\oplus)^2 = a^2(a^\oplus)^2 = aa^\oplus,$$

we have $b^\oplus a^\oplus \in ab\{3, 5\}$.

Note that $(abb^\oplus a^\oplus)ab = aa^\oplus ab = ab$ and $b^\oplus a^\oplus(abb^\oplus a^\oplus) = b^\oplus a^\oplus aa^\oplus = b^\oplus a^\oplus$, which gives $b^\oplus a^\oplus \in ab\{1, 2\}$. Hence, $ab \in R^\#$ with $(ab)^\# = b^\oplus a^\oplus$. In addition, it is easy to see that $(ab)^\oplus = b^\oplus a^\oplus$. In fact,

$$b^\oplus a^\oplus (ab)^2 = (ab)^\#(ab)^2 = ab \quad \text{and} \quad ab(b^\oplus a^\oplus)^2 = ab((ab)^\#)^2 = (ab)^\# = b^\oplus a^\oplus.$$

(iii) \Rightarrow (i) Obviously, $b(ab)^\# \in a\{2, 3\}$. Observe that $a = br_3$ for some $r_3 \in R$. Then, we have

$$\begin{aligned} a &= a(a^\oplus)^2 a^2 = br_3(a^\oplus)^2 a^2 = bb^\oplus br_3(a^\oplus)^2 a^2 = bb^\oplus a(a^\oplus)^2 a^2 \\ &= bb^\oplus a^\oplus a^2 = b(ab)^\# a^2, \end{aligned}$$

which implies that $aR \subseteq b(ab)^\# R$. Therefore, $a \in R^\oplus$ and $a^\oplus = b(ab)^\#$ by Theorem 3.1.

(iii) \Rightarrow (iv) Since $a = br_4$ for some $r_4 \in R$, we get

$$bb^\oplus a^\oplus = bb^\oplus a(a^\oplus)^2 = bb^\oplus br_4(a^\oplus)^2 = br_4(a^\oplus)^2 = a(a^\oplus)^2 = a^\oplus.$$

Therefore, we only need to prove $abb^\oplus \in R^\oplus$ with $(abb^\oplus)^\oplus = a^\oplus$. From $a(bb^\oplus a^\oplus) = aa^\oplus$, it follows that $a^\oplus \in abb^\oplus\{3\}$. Also, we have

$$abb^\oplus(a^\oplus)^2 = (abb^\oplus a^\oplus)a^\oplus = aa^\oplus a^\oplus = a^\oplus$$

and

$$a^\oplus(abb^\oplus)^2 = a^\oplus(abb^\oplus a^\oplus)a^2 bb^\oplus = a^\oplus aa^\oplus a^2 bb^\oplus = abb^\oplus.$$

Hence, we have $(abb^\oplus)^\oplus = a^\oplus$ by Lemma 2.1. Then, it is easy to see that $(ab)^\# = b^\oplus(abb^\oplus)^\oplus$.

(iv) \Rightarrow (v) It is obvious.

(v) \Rightarrow (i) From the proof of (iii) \Rightarrow (iv), we know $a^\oplus = bb^\oplus a^\oplus$, which gives $a^\oplus = b(ab)^\#$. \square

The next theorem gives the result related to $a^\oplus = b(ab)^\oplus$.

Theorem 4.4. *Let $a \in R^\#$, $ab \in R^\oplus$. Then the following are equivalent:*

- (i) $a \in R^\oplus$ and $a^\oplus = b(ab)^\oplus$;
- (ii) $aR \subseteq babR$.

Proof. (i) \Rightarrow (ii) Suppose that $a^\oplus = b(ab)^\oplus$. Then, we have

$$a = a^\oplus a^2 = b(ab)^\oplus a^2 = bab((ab)^\oplus)^2 a^2,$$

which implies that $aR \subseteq babR$.

(ii) \Rightarrow (i) Since $aR \subseteq babR$, there exists $w \in R$ such that $a = babw$. Then, we obtain $a = b(ab)^\oplus(ab)^2 w$, which gives $aR \subseteq b(ab)^\oplus R$. In addition, it is easy to verify $b(ab)^\oplus \in a\{2, 3\}$. Hence, by Theorem 3.1, it follows that $a \in R^\oplus$ with $a^\oplus = b(ab)^\oplus$. \square

Finally, we characterize the mixed-type reverse order laws $(ab)^\# = b^\oplus a^\oplus$ and $(ba)^\# = a^\oplus b^\oplus$ by the following necessary and sufficient conditions.

Theorem 4.5. *Let $a, b \in R^\oplus$. Then the following are equivalent:*

- (i) $ab, ba \in R^\#$ with $(ab)^\# = b^\oplus a^\oplus$, $(ba)^\# = a^\oplus b^\oplus$;
- (ii) $a^\oplus ab = baa^\oplus$, $b^\oplus ba = abb^\oplus$, and $aa^\oplus b^\oplus = b^\oplus a^\oplus a$.

Proof. (i) \Rightarrow (ii) Since $(ab)^\# = b^\oplus a^\oplus$, we have

$$ab = b^\oplus a^\oplus (ab)^2 = b^\oplus b (b^\oplus a^\oplus (ab)^2) = b^\oplus bab$$

and

$$ab = (ab)^2 b^\oplus a^\oplus = ((ab)^2 b^\oplus a^\oplus) a a^\oplus = ab a a^\oplus.$$

Similarly, from $(ba)^\# = a^\oplus b^\oplus$, it follows that $ba = a^\oplus aba = b a b b^\oplus$. Hence,

$$a^\oplus ab = (a^\oplus aba) a^\oplus = b a a^\oplus \text{ and } b^\oplus ba = (b^\oplus bab) b^\oplus = a b b^\oplus.$$

Since $ab, ba \in R^\#$, from Lemma 2.6, it follows that $a(ba)^\# = ab((ab)^\#)^2 a = (ab)^\# a$, which immediately yields $aa^\oplus b^\oplus = b^\oplus a^\oplus a$.

(ii) \Rightarrow (i) By the hypotheses, we deduce

$$a b b^\oplus (a^\oplus ab) = a b b^\oplus b a a^\oplus = a (b a a^\oplus) = a a^\oplus ab = ab,$$

$$(b^\oplus a^\oplus a) b b^\oplus a^\oplus = a a^\oplus b^\oplus b b^\oplus a^\oplus = (a a^\oplus b^\oplus) a^\oplus = b^\oplus a^\oplus a a^\oplus = b^\oplus a^\oplus$$

and

$$b^\oplus (a^\oplus ab) = (b^\oplus ba) a^\oplus = a b b^\oplus a^\oplus,$$

which imply $b^\oplus a^\oplus \in ab\{1, 2, 5\}$, i.e., $(ab)^\# = b^\oplus a^\oplus$. Similarly, we can obtain $(ba)^\# = a^\oplus b^\oplus$. \square

Acknowledgements.

This research was supported by China Postdoctoral Science Foundation (No. 2018M632385), the National Natural Science Foundation of China (No. 11771076), the Natural Science Foundation of Jiangsu Province (No. BK20141327), the Portuguese Funds through FCT-“Fundação para a Ciência e a Tecnologia”, within the project UID/MAT/00013/2013.

References

- [1] O.M. Baksalary, G. Trenkler, Core inverse of matrices, *Linear Multilinear Algebra* 58 (2010) 681-697.
- [2] O.M. Baksalary, G. Trenkler, Problem 48-1: reverse order law for the core inverse, *Image* 48 (2012) 40.
- [3] J. Benítez, X.J. Liu, J. Zhong, Some results on matrix partial orderings and reverse order law, *Electron. J. Linear Algebra* 20 (2010) 254-273.
- [4] J.L. Chen, H.H. Zhu, P. Patrício, Y.L. Zhang, Characterizations and representations of core and dual core inverses, *Canad. Math. Bull.* 60 (2017) 269-282.

- [5] R.E. Cline, An application of representation for the generalized inverse of a matrix, MRC Technical Report 592, 1965.
- [6] N. Cohen, E.A. Herman, S. Jayaraman, Solution to problem 48-1: reverse order law for the core inverse, *Image* 49 (2012) 46-47.
- [7] D.S. Cvetković-Ilić, M. Djikić, Various solutions to reverse order law problems, *Linear Multilinear Algebra* 64 (2016) 1207-1219.
- [8] D.S. Cvetković-Ilić, R. Harte, Reverse order laws in C^* -algebras, *Linear Algebra Appl.* 434 (2011) 1388-1394.
- [9] D.S. Cvetković-Ilić, V. Pavlović, A comment on some recent results concerning the reverse order law for $\{1, 3, 4\}$ -inverses, *Appl. Math. Comput.* 217 (2010) 105-109.
- [10] C.Y. Deng, Reverse order law for the group inverses, *J. Math. Anal. Appl.* 382 (2011) 663-671.
- [11] N.Č. Dinčić, D.S. Djordjević, Hartwig's triple reverse order law revisited, *Linear Multilinear Algebra* 62 (2014) 918-924.
- [12] M.P. Drazin, Pseudo-inverses in associative rings and semigroups, *Amer. Math. Monthly* 65 (1958) 506-514.
- [13] T.N.E. Greville, Note on the generalized inverse of a matrix product, *SIAM Rev.* 8 (1966) 518-521.
- [14] R.E. Hartwig, Block generalized inverses, *Arch. Ration. Mech. Anal.* 61 (1976) 197-251.
- [15] R.E. Hartwig, J. Luh, A note on the group structure of unit regular ring elements, *Pacific J. Math.* 71 (1977) 449-461.
- [16] S. Izumino, The product of operators with closed range and an extension of the reverse order law, *Tohoku Math. J.* 34 (1982) 43-52.
- [17] X.J. Liu, S.Q. Fu, Y.M. Yu, An invariance property of operator products related to the mixed-type reverse order laws, *Linear Multilinear Algebra* 64 (2016) 885-896.
- [18] X.J. Liu, M. Zhang, J. Benítez, Further results on the reverse order law for the group inverse in rings, *Appl. Math. Comput.* 229 (2014) 316-326.
- [19] X. Mary, Reverse order law for the group inverse in semigroups and rings, *Comm. Algebra* 43 (2015) 2492-2508.
- [20] D. Mosić, N.Č. Dinčić, Reverse order law $(ab)^\dagger = b^\dagger(a^\dagger abb^\dagger)^\dagger a^\dagger$ in rings with involution, *Filomat* 28 (2014) 1791-1815.

- [21] D. Mosić, D.S. Djordjević, Reverse order law for the Moore-Penrose inverse in C^* -algebras, *Electron. J. Linear Algebra* 22 (2011) 92-111.
- [22] D. Mosić, D.S. Djordjević, Some results on the reverse order law in rings with involution, *Aequat. Math.* 83 (2012) 271-282.
- [23] D. Mosić, D.S. Djordjević, Reverse order law for the group inverse in rings, *Appl. Math. Comput.* 219 (2012) 2526-2534.
- [24] D. Mosić, D.S. Djordjević, The reverse order law $(ab)^\# = b^\dagger(a^\dagger abb^\dagger)^\dagger a^\dagger$ in rings with involution, *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas* 109 (2015) 257-265.
- [25] V. Pavlović, D.S. Cvetković-Ilić, Applications of completions of operator matrices to reverse order law for $\{1\}$ -inverses of operators on Hilbert spaces, *Linear Algebra Appl.* 484 (2015) 219-236.
- [26] R. Penrose, A generalized inverse for matrices, *Proc. Cambridge Philos. Soc.* 51 (1955) 406-413.
- [27] P. Puystjens, R.E. Hartwig, The group inverse of a companion matrix, *Linear Multilinear Algebra* 43 (1997) 137-150.
- [28] D.S. Rakić, N.Č. Dinčić, D.S. Djordjević, Group, Moore-Penrose, core and dual core inverse in rings with involution, *Linear Algebra Appl.* 463 (2014) 115-133.
- [29] Y. Tian, On mixed-type reverse-order laws for the Moore-Penrose inverse of a matrix product, *Int. J. Math. Math. Sci.* 58 (2004) 3103-3116.
- [30] H.X. Wang, X.J. Liu, Characterizations of the core inverse and the core ordering, *Linear Multilinear Algebra* 63 (2015) 1829-1836.
- [31] S.Z. Xu, J.L. Chen, X.X. Zhang, New characterizations for core inverses in rings with involution, *Front. Math. China* 12 (2017) 231-246.