

Group-regular rings

Xavier Mary · Pedro Patrício

Abstract We propose different generalizations of unit-regularity of elements in general rings (non necessarily unital rings). We then study general rings for which all elements have these properties. We notably compare them with unit-regular ideals and general rings with stable range one. We also prove that these rings are morphic rings.

Keywords unit-regular elements · strongly regular (group) elements · general rings · adjoint semigroup

Mathematics Subject Classification (2000) 16U99 · 16E50 · 15A09

1 Introduction

In this paper, R denotes a general ring (that may or may not contain an identity), and $E(R)$ its set of idempotents. To emphasize the distinction between rings with or without an identity, we call the former unital rings, the latter non-unital rings and by a ring we always mean a general ring (without necessarily an identity). While every unital ring may be regarded as the endomorphism ring of a module (thus making all the module-theoretical statements available), non-unital rings cannot be endomorphism rings. Thus new approaches (generally elementwise statements with more “elementary” proofs) have to be introduced.

We say that a is (von Neumann) regular in R if $a \in aRa$. A particular solution to $axa = a$ is called an inner inverse of a . A solution to $xax = a$ is called a weak (or outer) inverse. Finally, an element that satisfies $axa = a$ and $xax = x$ is called an inverse (or reflexive inverse, or relative inverse) of a . A commuting inverse, if it exists, is unique and denoted by $a^\#$. It is the unique solution to:

$$ax = xa, axa = a, xax = x.$$

It is usually called the group inverse of a , for a is group invertible if and only if it belongs to a subgroup G_a of the multiplicative semigroup (R, \cdot) (for instance the commutative subgroup

Xavier Mary
Laboratoire Modal'X, Université Paris Nanterre
E-mail: xavier.mary@parisnanterre.fr

Pedro Patrício,
CMAT – Centro de Matemática and Departamento de Matemática e Aplicações, Universidade do Minho
E-mail: pedro@math.uminho.pt

$G_a = \{a, a^\#, aa^\#\}$, which $aa^\#$ is the identity of the group). We let $R^\#$ denote the set of group invertible elements. These are exactly the strongly regular elements of R , where $a \in R$ is strongly regular if $a \in a^2R \cap Ra^2$.

We will use without further comment that for a group element $x \in R$, $xR = x^\#R = eR$ with $e = xx^\# = x^\#x \in E(R)$, and dually.

If R is unital, we say a is unit-regular if $a \in aU(R)a$, where $U(R)$ denotes the set of invertible elements (units). A ring is regular if all its elements are regular, and a unital ring is unit-regular if all its elements are unit-regular.

Many papers have recently focus on the extension of notions defined for unital rings to general rings, see for instance [3], [29], where the authors define exchange elements and clean elements in general rings. They use the adjoint semigroup of the general ring $(R, \cdot, +)$, defined as the set R with adjoint operation $x * y = x + y + xy$. The semigroup $(R, *)$ is actually a monoid with identity 0, and we define $Q(R) = \{q \in R \mid \exists q' \in R, q + q' + qq' = q + q' + q'q = 0\}$, group of units of $(R, *)$. Elements of $Q(R)$ are called quasiregular elements of R , and play a prominent role in the theory of Jacobson radicals. If R is unital, then the monoids (R, \cdot) and $(R, *)$ are isomorphic through the map $x \mapsto x - 1$, and so are $U(R)$ and $Q(R)$. Generalizations of unit-regular elements have not been defined in this manner. We do so in Section 2. In this section we also propose a second method, inspired by semigroup theory ([13], [16], [24]) that uses group-invertible elements instead of solely the group of units, that may even not exist. Section 3 relates these notions of regularity with unit-regularity in ideal extensions. In Section 4 we study general rings where all elements have such properties, and show that they define a unique class of general rings (called group-regular rings afterwards). We then characterize these rings by means of unitizations. This allows us to compare group-regular rings with two other classes of rings: rings with stable range 1 [33] and unit-regular ideals [9]. An example shows that, in lack of an identity, group-regular rings need not have stable range 1. Finally in Section 5 we give a characterization of group-regular rings in terms of isomorphic idempotents. We then deduce that group-regular rings are morphic (see [28] for the notion), thus recovering one half of the equivalence of Ehrlich [11]: a unital ring is unit-regular if and only if it is regular and (left) morphic.

We will make use of the natural partial order (minus partial order) on a regular semigroup, defined independently by Hartwig[18] and Nambooripad[27] and extended by Mitsch[26] to non-regular semigroups. It is defined for a regular $b \in R$ by $a \leq b \Leftrightarrow \exists e, f \in E(R), a = eb = bf$. For idempotents, this reduces to $e \leq f \Leftrightarrow ef = fe = e$. We will also use corner rings eRe ($e \in E(R)$).

Finally, we recall the following definitions and results about stable range 1, regular rings, unit-regular rings and corner rings.

A unital ring R has stable range 1 if for all $a, b \in R$, $aR + bR = R$ implies that $(a + bc)R = R$ for some $c \in R$ (equivalently, $a + bc \in U(R)$ by [33] Theorem 2.6). A (general) ring R has stable range 1 if for all $a \in R, b \in \hat{R}$, $(1 + a)\hat{R} + b\hat{R} = \hat{R}$ implies that $(1 + a + bc)\hat{R} = \hat{R}$ for some $c \in \hat{R}$ and some (all [33] Theorem 3.6) unitization \hat{R} of R .

Lemma 1 ([12] Lemma 2) *A regular ring is the directed union of its corner rings.*

Lemma 2 ([14] Theorem) *A regular ring admits a regular unitization.*

The next lemma is well-known in the literature. A proof based on module cancellation is due to Ehrlich[11] and Handelman[17]. The first ring theoretical (and element wise) proof is probably due to Kaplansky, as explained in [20]. The link between unit-regularity in a ring and unit-regularity in a corner ring is then precisely studied in [23].

Lemma 3 ([20] Proposition 8) *Corner rings of a unit-regular ring are unit-regular.*

Proposition 1 (Fuchs and Kaplansky [15] Proposition 4.12) *A unital ring R is unit-regular iff it is regular with stable range 1.*

Lemma 4 ([25] Lemma 1.4, [8] Lemma 1) *Let R be a regular ring. Then the following are equivalent:*

1. *For each idempotent $e \in E(R)$ the corner ring eRe is unit-regular.*
2. *R admits a unit-regular unitization.*
3. *R has stable range 1.*

2 Group-regular, group-dominated and Q -unit-regular elements

There are many equivalent characterizations of a unit-regular element a in R unital ring. By definition, $aua = a$ for some unit $u \in U(R)$. But also $a = eu = vf$ for idempotents e, f and units u, v . One can also characterize unit-regularity by means of an inequality: a is unit-regular if and only if $a \leq u$ for some unit u ([32] Theorem 4.3, see also Proposition 2 below), that is we can use the same unit in the previous statement. Finally, since $u \mapsto u - 1$ is a group isomorphism from $U(R)$ to $Q(R)$, $a(1 + q)a = a$ for a quasi-regular element $q \in Q(R)$.

Definition 1 Let $a \in R$ general ring. We say that:

1. a is group-regular if $a = axa$ for some $x \in R^\#$;
2. a is intra group-regular if there exists $x \in R^\#$ such that $axa = a$ and $a^2 = axx^\#a$;
3. a is group-dominated if $a \leq x^\#$ for some $x \in R^\#$;
4. a is Q -unit-regular if $a^2 + aqa = a$ for some $q \in Q(R)$.

R is group-regular (resp. intra group-regular, group-dominated, Q -unit-regular) if every element of R is group-regular (resp. intra group-regular, group-dominated, Q -unit-regular).

We have the following characterizations of group-domination. The third one is a generalization of [7] and [6] Theorem 5. The fifth one goes back to [19] for unit-regular elements.

Proposition 2 *Let $a, x \in R$ with $x \in R^\#$. The following statements are equivalent:*

- (1) $a \leq x^\#$;
- (2) $Ra \subseteq Rx$, $aR \subseteq xR$ and $axa = a$;
- (3) $Ra \subseteq Rx$, $aR \subseteq xR$ and $aR \cap (a - x^\#)R = \{0\}$ (or $Ra \cap R(a - x^\#) = \{0\}$);
- (4) a is unit-regular in the corner ring $xx^\#Rxx^\#$;
- (5) $a = ex^\#$ for some $e \in E(R)$ such that $e \leq xx^\#$ ($a = x^\#f$ for some $f \in E(R)$ such that $f \leq xx^\#$).

Proof Let $a \in R, x \in R^\#$.

- (1) \Rightarrow (2) Assume (1) and let $e, f \in E(R)$ such that $a = ex^\# = x^\#f$. Then $Ra = Re(x^\#)^2x \subseteq Rx$, $aR = x(x^\#)^2fR \subseteq xR$, and $axa = ex^\#xx^\#f = ex^\#f = eex^\# = a$.
- (2) \Rightarrow (3) Assume (2) and let $b = ac = (a - x^\#)d$ for some $c, d \in R$. First $ax \in E(R)$, and as $Ra \subseteq Rx = Rxx^\#$ then $a = axa = yxx^\#$ for some $y \in R$, and $a = axx^\#$ since $xx^\# \in E(R)$. Then $b = axx^\#c = ax = axaxx^\#c = axb = ax(a - xx^\#) = axa - a = 0$. The other statement is dual.

- (3) \Rightarrow (4) Assume (3) and pose $e = xx^\# \in E(R)$. As $Rx = Re$ and $xR = eR$ then $a \in eRe$ (in particular $axx^\# = a = xx^\#a$). Also $xx^\# = x^\#x = e$ by definition and x is a unit of eRe . We now compute $axa - a = ax(a - x^\#) = (a - x^\# + x^\#)x(a - x^\#) = (a - x^\#)x(a - x^\#) + a - x^\# = (a - x^\#)[x(a - x^\#) + xx^\#] \in aR \cap (a - x^\#)R = \{0\}$.
- (4) \Rightarrow (5) Assume (4). Then $axx^\# = a = x^\#xa$ and $axax = ax$. Thus $e = ax \in E(R)$, and $(ax)(xx^\#) = ax = (xx^\#)(ax)$ ($ax \leq xx^\#$). The other statement is dual.
- (5) \Rightarrow (1) Assume (5) and let $a = ex^\#$ with $e \in E(R)$, $e \leq xx^\#$. Pose $f = xex^\#$. Then $f^2 = xex^\#xex^\# = xex^\# = f \in E(R)$ and $x^\#f = x^\#xex^\# = ex^\# = a$.

Proposition 2 shows that group-domination is a localized version of unit-regularity, that proved important in the theory of Leavitt path algebras [1], [31]. It claims in particular that every group-dominated element a of a ring R lies in some corner ring eRe in which it is unit-regular. It does not claim however that if $a \in fRf$, $f \in E(R)$, it is unit-regular in this corner ring. Indeed [23] gives an example of a element $a \in eRe \subseteq R$ ring of two by two matrices over a special ring S such that a is unit-regular in R but not in eRe .

We have the following implications:

Corollary 1 *Let $a \in R$. Then*

$$\{a \text{ group-dominated}\} \Rightarrow \{a \text{ intra group-regular}\} \Rightarrow \{a \text{ group-regular}\} \Rightarrow \{a \text{ } Q\text{-unit-regular}\}.$$

Proof The first implication follows from Proposition 2, and the second is tautological. We now prove the third one.

Let a be group-regular with $x \in R^\#$ such that $axa = a$ and pose $q = x - xx^\# - xa + xaxx^\#$. Pose also $q' = x^\# - xx^\# + xx^\#a - xx^\#axx^\#$. Then $qx^\# = xx^\# - x^\#$, $qxx^\# = x - xx^\#$ and computations give $qq' = xx^\# - x^\# - x + xx^\# + xa - xx^\#a - xaxx^\# + xx^\#axx^\# = q + q'$. On the other hand, $x^\#q = q'$ and $xx^\#q = q$ and $(xx^\#a)q = (xx^\#a)(xx^\#q)$ and $q'q = q' + q$. This proves that q is quasiregular. We finally compute $a^2 + aqa = a^2 + a(x - xx^\# - xa + xaxx^\#)a = a^2 + axa - axx^\#a - axa^2 + axaxx^\#a$ and since $axa = a$, then $a^2 + aqa = a$. Thus a is Q -unit-regular.

In the next example, we construct a ring R with an element $a \in R$ group-regular but not group-dominated, so that (some) of the reverse implications do not hold in general.

Example 1 This example is inspired by Example 4.8 and Remark 4.9 in [30]. Let F be a field. Consider the semigroup S quotient of the free semigroup with two generators a, e by the relations $aea = a$, $e^2 = e$ and let $R = F[S]$ be the semigroup ring of S over F . Equivalently, R is the non-unital free algebra on two indeterminates a, e (excluding the empty word) over F quotiented by the above relations. These relations form a reduction system, whose ambiguities are all resolvable in the sense of [4]. Equivalently, by Theorem 1.2 of [4], every element of R is reduction unique (it has a unique canonical form, where all occurrences of the left-hand side of the relations are replaced by their right-hand side). First of all, a (as an element of R , $a = 1a$) is group-regular with inner inverse $e = 1e$. We now prove that a is not group-dominated because it does not belong to any corner ring. Contrary to [30], we cannot work only on monomials inner inverses of a here, because we would need to consider all (not just one) inner inverses. And some inner inverses are not monomials: for instance $e + a - ea^2$ satisfies $a(e + a - ea^2)a = a + a^3 - aea^3 = a$.

So assume that there exists an idempotent $f \in E(R)$ such that $a \in fRf$, that is $fa = af = a$. As ea is reduced then $\lambda ea \neq a$ for any $\lambda \in F$ and f is not a multiple of e . By applying the reductions above, we thus see that f can be written as

$$f = \lambda e + \sum_{k=1}^n \left(\lambda_k^0 a^k + \lambda_k^l e a^k + \lambda_k^r a^k e + \lambda_k^{lr} e a^k e \right)$$

for some $n \geq 1$, with all coefficients in F , and one of $\lambda_n^0, \lambda_n^l, \lambda_n^r, \lambda_n^{lr}$ non zero. After applying the reductions, we get that

$$fa = \lambda_1^r a + (\lambda + \lambda_1^{lr})ea + \sum_{k=2}^n [(\lambda_{k-1}^0 + \lambda_k^r)a^k + (\lambda_{k-1}^l + \lambda_k^{lr})ea^k]$$

and dually

$$af = \lambda_1^l a + (\lambda + \lambda_1^{lr})ae + \sum_{k=2}^n [(\lambda_{k-1}^0 + \lambda_k^l)a^k + (\lambda_{k-1}^r + \lambda_k^{lr})a^k e].$$

We deduce that the coefficients of f satisfy:

$$\begin{cases} \lambda + \lambda_1^{lr} = 0 \\ \lambda_1^l = \lambda_1^r = 1 \\ \lambda_k^l = \lambda_k^r = -\lambda_{k-1}^0 \quad (\forall k \geq 2) \\ \lambda_k^{lr} = -\lambda_{k-1}^r = -\lambda_{k-1}^l \quad (\forall k \geq 2) \end{cases}$$

We finally consider the coefficients of rank $n \geq 1$ of f . If $\lambda_n^0 \neq 0$ then $f^2 = f$ contains a^{2n} hence $2n \geq n < 2n$ whence a contradiction. Thus $\lambda_n^0 = 0$. Also if $\lambda_n^l \neq 0$ then as $\lambda_n^l = \lambda_n^r$ then $f^2 = f$ contains $ea^{2n}e$ whence a contradiction. Thus $\lambda_n^l = \lambda_n^r = 0$. It follows that $\lambda_n^{lr} \neq 0$, and $f^2 = f$ contains $ea^{2n-1}e$. Thus $2n-1 \leq n$ and $n = 1$. But we then get a contradiction since $\lambda_1^l = 1 = 0$.

Finally such an f does not exist, and a cannot be group-dominated.

In Example 1, the ring is not unital. This is actually a necessary condition because it happens that in unital rings, all the previous concepts are equivalent elementwise. Example 1 thus show that the situation is drastically different for non-unital rings than for unital ones.

Proposition 3 *Let $a \in R$ unital ring. Then the following statements are equivalent:*

- (1) a is unit-regular;
- (2) a is group-dominated (locally unit-regular);
- (3) a is intra group-regular;
- (4) a is Q -unit-regular;
- (5) a is group-regular.

Proof That (1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4) is straightforward by Corollary 1. (4) \Rightarrow (5) because if $a = a^2 + aqa$ for some $q \in Q(R)$, then $a = axa$ where $x = 1 + q \in U(R) \subseteq R^\#$. We finally prove that (5) \Rightarrow (1). Let $a \in R, x \in R^\#$ such that $axa = a$. Pose $y = 1 + x - xx^\# - xa(1 - xx^\#)$. Then $y' = 1 + x^\# - xx^\# + xx^\#a(1 - xx^\#)$ satisfies $yy' = y'y = 1$ hence $y \in U(R)$, and $aya = a$.

In particular, we recover directly Arens-Kaplansky's result that completely regular rings are unit-regular. The particular case of group-dominated elements can actually be deduced from Proposition 2 and results from two other papers: the main Theorem in [23] and Lemma 3 in [31].

Corner rings of general rings provide unital rings. We will use without further comments that if a has any of the previous properties in a corner ring $eRe, e \in E(R)$ then it has the property in R , because of the equalities $(eRe)^\# = R^\# \cap eRe$ and $Q(eRe) = Q(R) \cap eRe$.

Also, we deduce from Proposition 3 a converse to Corollary 1 in the special case of semicentral idempotents. Recall that an idempotent $e \in E(R)$ is right semicentral [5] if one of the following equivalent conditions hold: 1) $eRe = eR$, 2) $(\forall r \in R) ere = er$, 3) Re is an ideal.

Proposition 4 *Let $e \in E(R)$ be a right semicentral idempotent of R and $a \in eRe = eR$. Then the following implication holds:
if a is Q -unit-regular (in R), then it is unit-regular in eRe , and in particular group-dominated (in R).*

Proof Let $e \in E(R)$ such that $eRe = eR$, and $a \in eRe$. Assume that a is Q -unit-regular and let $q \in Q(R)$ such that $a^2 + aqa = a$. Let q' be the inverse of q in the group $Q(R)$. As $ae = ea = a$ then $a(e + eq)a = a$. As $eq, eq' \in eR = eRe$, then $eqe = eq$ and $eq'e = eq'$. Pose $x = e + eq$ and $x' = e + eq'$. As $q + q' - qq' = q' + q - q'q = 0$ then $xx' = x'x = e$ and $xx'x = x, x'xx' = x$. It follows that x, x' are inverses in $(eRe)^{-1}$ and a is unit-regular in the unital corner ring eRe since $axa = a$.

3 Ideal extensions

Our next results deal with ideal extensions.

Lemma 5 *Let $a \in R$. Consider the following statements:*

- (1) *a is Q -unit-regular.*
- (2) *Whenever $R \triangleleft T$ where T is a unital ring, a is unit-regular in T .*
- (3) *$(0, a)$ is unit-regular in $\mathbb{Z} \oplus R$ (Dorroh unitization of R).*
- (4) *a or $-a$ is Q -unit-regular.*

Then (1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4).

Proof Let $a \in R$.

- (1) \Rightarrow (2) Assume a is Q -unit-regular, with $a^2 + aqa = a$ for $q \in Q(R)$. Then $1 + q$ is a unit of T and $a(1 + q)a = a$.
- (2) \Rightarrow (3) Follows from $R \triangleleft T = \mathbb{Z} \oplus R$.
- (3) \Rightarrow (4) Assume $(0, a)$ is unit-regular in $\mathbb{Z} \oplus R$. Then exists $u = (n, r) \in (\mathbb{Z} \oplus R)^{-1}$, $(0, a)(n, r)(0, a) = (0, a)$. First, as u is a unit then $n = 1$ or $n = -1$ and $na^2 + ara = a$. If $n = 1$ then $q = u - 1 = (0, r) \in Q(\mathbb{Z} \oplus R)$. Thus exists $q' = (m, r')$, $q + q' + qq' = q + q' + q'q = 0$ and it follows that $m = 0$ and $r + r' + rr' = r + r' + r'r = 0$ in R , so that $r \in Q(R)$. Finally $a = a^2 + ara$ is Q -unit-regular. If $n = -1$ then $(0, -a)(1, -r)(0, -a) = -(0, a) = (0, -a)$ and $(0, -a)$ is unit-regular with unit $-u = (1, -r)$. Thus $q = -u - 1 = (0, -r) \in Q(\mathbb{Z} \oplus R)$ and as previously $-r \in Q(R)$. Finally from $-a^2 + ara = a$ we deduce $(-a)^2 + (-a)(-r)(-a) = -a$ and $-a$ is Q -unit-regular.

As in a unital ring T , $a \in T$ is unit-regular (with $aua = a, u \in T^{-1}$) if and only if $-a$ is unit-regular ($-a = (-a)(-u)(-a), -u \in T^{-1}$) we deduce:

Corollary 2 *Let $a \in R$. Then the following statements are equivalent:*

- (1) *a or $-a$ is Q -unit-regular.*
- (2) *Whenever $R \triangleleft T$ where T is a unital ring, a and $-a$ are unit-regular in T .*
- (3) *$(0, a)$ and $(0, -a)$ are unit-regular in $\mathbb{Z} \oplus R$.*

It is actually not clear whether, in lack of a identity, a Q -unit-regular implies $-a$ Q -unit-regular.

We now consider group-domination.

Lemma 6 *Let $a \in R$. Then the following statements are equivalent:*

- (1) a is group-dominated.
(2) There exists $e \in E(R)$ such that $a \in eRe$ and whenever $R \triangleleft T$ where T is a ring, $1 + a - e$ is unit-regular in T .
(3) There exists $e \in E(R)$ such that $a \in eRe$ and $(1, a - e)$ is unit-regular in $\mathbb{Z} \oplus R$.
(4) There exists $e \in E(R)$ such that $a \in eRe$, and there exists a unital ring S such that R is a left and right module over S and $(1, a - e)$ is unit-regular in $S \oplus R$.

Proof Let $a \in R$.

- (1) \Rightarrow (2) Assume $a \leq x^\#$ for some $x \in R^\#$ and pose $xx^\# = e$. Then $1 + x - xx^\# = (1 - e) + x$ is a unit of T and $(1 - e + a)(1 - e + x)(1 - e + a) = (1 - e)^3 + axa = 1 - e + a$.
(2) \Rightarrow (3) Follows from $R \triangleleft T = \mathbb{Z} \oplus R$.
(3) \Rightarrow (4) Take $S = \mathbb{Z}$.
(4) \Rightarrow (1) Let $e \in E(R)$ such that $a \in eRe$ and S be a ring such that R is a left and right module over S and $(1, a - e)$ is unit-regular in $T = S \oplus R$. Then $(0, a) + [1 - (0, e)]$ is unit-regular with $(0, a) \in (0, e)T(0, e)$ and by [23], it is unit-regular in the corner ring $(0, e)T(0, e) = (0, eRe)$. Finally a is locally unit-regular, that is group-dominated.

We finally consider the case R has a unit-regular unitization:

Lemma 7 *Let R be a left (right, two-sided) ideal of T where T is a unit-regular ring, and $a \in R$. Then a is group-dominated.*

Proof First, as T is regular then R is regular. Indeed, let $a \in R, x \in T$ such that $axa = a$. Then $a = a(xax)a$ is regular in R . Let now $a \in R$. As R is regular, it is the directed union of its corner rings and exists $e \in E(R)$ such that $a \in eRe$. As corner rings of unit-regular rings are unit-regular, then a is unit-regular in $eTe = eRe$ since R is a left (right, two-sided) ideal and $e \in R$. Denote by $x \in U(eRe)$ its inverse and pose $x^\# = x_{eRe}^{-1}$ (inverse in the corner ring). Then $a \leq x^\#$ and a is group-dominated.

4 Group-regular rings

If R is a ring without identity, Example 1 shows that the previous concepts are different elementwise. It happens however that they provide a unique global characterization, that is group-regular rings, group-dominated rings and Q -unit-regular rings are the same. This is based on the following observation. If an element is group-regular or Q -unit-regular in a corner ring, then it is actually unit-regular in this corner ring hence group-dominated in the whole ring. This happens for instance if any finite set of elements of the ring lies in a corner ring. Such rings are usually called rings with ‘‘local units’’ ([2, Definition 1]), but the terminology may however have other meanings. By Lemma 1 this is the case for von Neumann regular general rings. Thus we deduce:

Theorem 1 *Let R be a (general) ring. Then the following statements are equivalent:*

- (1) R is group-dominated ($\forall a \in R, a$ is group-dominated);
(2) R is group-regular ($\forall a \in R, a$ is group-regular);
(3) R is Q -unit-regular ($\forall a \in R, a$ is Q -unit-regular).

Proof Let R be a ring.

- (1) \Rightarrow (2) Straightforward;

- (2) \Rightarrow (3) Follows directly from the last implication of Corollary 1.
 (3) \Rightarrow (1) Assume (3) and let $a \in R$. Then exists $q \in Q(R)$ such that $a^2 + aqa = a$ and $a = a(a + aq + q)a$ is regular. Thus the whole ring R is regular and exists $e \in E(R)$ such that $a \in eRe$. Also exists $f \in E(R)$ such that $e, a, q \in fRf$. As a is Q -unit-regular in the unital ring fRf it is unit-regular in fRf , hence group-dominated in R by Proposition 2.

Obviously, this is also equivalent with being an intra-group-regular ring by Corollary 1.

Example 2 Let R be the ideal of all linear operators of finite rank on a vector space (over a field F) of countably infinite dimension E , and T be the ring of all linear operators on E . It is known that T is regular but not unit-regular [10]. Thus R is regular, but we cannot use Lemma 7 directly to show that R is group-regular. Let $a \in R$. As R is regular, then $a \in eRe = eTe$ with $e \in E(R)$ of finite rank n . But eRe is isomorphic to $\mathcal{M}_n(\mathbb{K})$ which is unit-regular, and a is group-regular.

In the example, all corner rings of R are unit-regular so that R has stable range 1 by Lemma 4. And if R has stable range 1, then it admits a unit-regular unitization (still by Lemma 4) and by Lemma 7, it is group-regular. Thus we may wonder whether the two concepts are equivalent. This is not the case, as shows next example.

Example 3 Let T_0 be a regular non unit-regular unital ring. We define iteratively $T_{n+1} = \mathcal{M}_2(T_n)$ for all $n \in \mathbb{N}$, and embed each $T_n, n \in \mathbb{N}$ as the 1-1 corner of T_{n+1} . We then define $R = \varinjlim T_n$, direct limit of T_n . First, we deduce by induction that each $T_n, n \in \mathbb{N}$ is regular since matrix rings over regular rings are regular (Theorem 24 in [21]). Also R has not stable range 1 since some corner rings are not unit-regular: for instance, T_0 is a non unit-regular corner ring by assumption. Let now $a \in R$. Then $a \in T_n$ for some $n \in \mathbb{N}$, and as T_n is regular we there exist $b \in T_n$ an inner inverse of a . Pose $B = \begin{pmatrix} b & b-1 \\ 1 & 1 \end{pmatrix} \in T_{n+1}$. Then B is group-invertible in R with group inverse $B^\# = \begin{pmatrix} 1 & 1-b \\ -1 & b \end{pmatrix} \in T_{n+1}$ and $\begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix} B \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix}$. It follows that a is group-regular, and R is group-regular.

Group-regular rings share many properties with unit-regular rings. For instance, they are dependent, a result due to Ehrlich ([10] Theorem 6) for unit-regular rings.

Corollary 3 *Let R be a group-regular ring, and $a, b \in R$. Then there exists $s, t \in R$ with s or t non zero such that $sa + tb = 0$.*

Proof Let $a, b \in R$. As R is group-regular it is group dominated by Theorem 1 and there exists $x, y \in R^\#$ such that $a \leq x^\#$ and $b \leq y^\#$, and in particular x and y are inner inverses of a and b respectively. Also, as R is Von Neumann regular, then $a, b, x, x^\#, y$ and $y^\#$ lie in a common corner ring eRe , $e \in E(R)$. If $e = xx^\# = yy^\#$, then by Proposition 2, a and b are unit-regular in the unital ring eRe , and by [10] Theorem 6 a and b are dependent. So assume that $e \neq xx^\#$ (the other case is symmetric). Then $s = e - xx^\# \neq 0$. But as $a \leq x^\#$ then $sa = (e - xx^\#)a = ea - xx^\#a = a - a = 0$. Letting $t = 0$ gives the desired property.

Using Proposition 3, Corollary 1, Lemma 5 on ideal extensions and Theorem 1 we also deduce a characterization of group-regular rings in terms of their unitizations.

Corollary 4 *Let R be a ring. Then the following statements are equivalent:*

1. R is group-regular;

2. For any unitization T of R , all elements of R are unit-regular in T ;
3. All elements of R are unit-regular in the Dorroh extension $\hat{R} = \mathbb{Z} \oplus R$ of R .

Proof Let R be a ring.

- (1) \Rightarrow (2) Assume that R is group-regular, and let T be a unitization of R . Let also $a \in R$. As a is group-regular in R it is group-regular in T and by Proposition 3, a is unit-regular in T .
- (2) \Rightarrow (3) The Dorroh extension $\hat{R} = \mathbb{Z} \oplus R$ is a unitization of R .
- (3) \Rightarrow (1) Assume (3) and let $a \in R$. By (3) there exists $u \in \hat{R}$ such that $aua = a$, and $a = a(uau)a$ is regular in R since $uau \in R$ ideal of \hat{R} . Thus the whole ring R is regular. We now prove that a is Q -regular in R . By (3) $a = (0, a)$ is unit-regular in \hat{R} and by Lemma 5, a or $-a$ is Q -unit-regular in R . Assume that $-a$ is Q -unit-regular. As R is regular then exists $f \in E(R)$ such that a lies in the (unital) corner ring fRf . By Proposition 3, $-a$ is group-regular. Let $x \in R^\#$ be an inner of $-a$. Then $-x \in R^\#$ is an inner inverse of a , and a is group-regular hence Q -unit regular by Corollary 1. Finally in both cases a is Q -unit-regular, and the whole ring is Q -unit-regular. By Theorem 1, R is a group-regular ring.

In [9], Chen et al. define an ideal R of a unital ring T to be a unit-regular ideal if all elements of R are unit-regular in T . Corollary 4 thus claims that R is group-regular if and only if it is a unit-regular ideal of any of its unitizations, if and only if it is a unit-regular ideal of its Dorroh extension $\hat{R} = \mathbb{Z} \oplus R$. In particular, Lemma 1 and Theorem 5 in [9] together with Corollary 4 give the following equivalences:

Corollary 5 *Let R be a regular ring, and $\hat{R} = \mathbb{Z} \oplus R$ be the Dorroh extension of R . Then the following statements are equivalent:*

1. R is group-regular;
2. For any unitization T of R and any $a \in R, b \in T$, if $aT + bT = T$ then exists $c \in T$ such that $a + bc \in U(T)$;
3. For any $a \in R, b \in \hat{R}$, if $a\hat{R} + b\hat{R} = \hat{R}$ then exists $c \in \hat{R}$ such that $a + bc \in U(\hat{R})$;
4. For any unitization T of R and any $a, b \in R$, if $aR \cong bR$ then a and b are equivalent in T ($(\exists u, v \in U(T)) a = ubv$);
5. For any $a, b \in R$, if $aR \cong bR$ then a and b are equivalent in \hat{R} ($(\exists u, v \in U(\hat{R})) a = ubv$).

We close this section by an open question. It is known that unit-regular elements may not be clean [22], but that a unit-regular ring is clean [7].

Question: Are group-regular rings clean general rings (every element is the sum of an idempotent and a quasiregular element [29]) ?

5 Group-regular rings, isomorphic idempotents and the morpnic property

From the results of Ehrlich [11] and Handelman [17] regarding module cancellation in unit-regular rings, it was deduced that unit-regular rings can be characterized in terms of isomorphic idempotents: a unital ring is unit-regular if and only if it is regular and complementary idempotents of isomorphic idempotents are isomorphic. Recall that two idempotents $e, f \in E(R)$ are isomorphic (or Kaplansky equivalent, or Murray-von Neumann equivalent), denoted by $e \sim f$, if $Re \cong Rf$ as left modules, or equivalently if $e = ab$ and $f = ba$ for some $a, b \in R$. We can always choose a and b to be reflexive inverses, $aba = a, bab = b$. Then Calugareanu [6] gave a purely ring theoretical, elementwise proof of the result. We can restate his elementwise results as follows.

Theorem 2 (Ehrlich, Handelman, Calugareanu)

Let $e = ab, f = ba$ be isomorphic idempotents with $aba = a, bab = b$ in a unital ring R . Then:

1. if a is unit-regular then $1 - e \sim 1 - f$;
2. if $1 - e \sim 1 - f$ then a is unit-regular.

We deduce from all the previous results the following characterization of group-regular rings in terms of isomorphic idempotents.

Theorem 3 Let R be a regular ring. Then the following statements are equivalent:

1. R is group-regular;
2. for all $e, f \in E(R)$, if $e \sim f$ then $1 - e \sim 1 - f$ in any unitization T of R ;
3. for all $e, f \in E(R)$, if $e \sim f$ then $1 - e \sim 1 - f$ in the Dorroh extension $\hat{R} = \mathbb{Z} \oplus R$.

Proof

- (1) \Rightarrow (2) Let T be a unitization of R group-regular ring, and let $e = ab, f = ba$ be isomorphic idempotents with $aba = a, bab = b$. As R is group-regular a is actually group-dominated by Theorem 1, and exists $x \in R^\#$, $a \leq x^\#$. Then $1 + x - xx^\#$ is a unit in T , and an inverse of a , so that a is unit-regular in T . It then follows by Theorem 2 that $1 - e \sim 1 - f$ in T .
- (2) \Rightarrow (3) Straightforward.
- (3) \Rightarrow (1) Let $a \in R$. As a is regular, it admits a reflexive inverse x , and $e = ax \sim xa = f$ are isomorphic idempotents. By assumption, $1 - e \sim 1 - f$ in the Dorroh extension $\hat{R} = \mathbb{Z} \oplus R$ and by Theorem 2, a is unit regular in \hat{R} . By Lemma 5, a or $-a$ is Q -unit-regular in R . Assume that $-a$ is Q -unit-regular with inverse q . As R is regular then exists $g \in E(R)$, $a, q \in gRg$ and by Proposition 3, $-a$ is group-dominated in the unital ring gRg . But then $a = -(-a)$ is also group-dominated, hence Q -unit-regular in R by Corollary 1. Finally, in the two-cases a is Q -unit-regular in R . Thus R is a Q -unit-regular ring hence a group-regular ring by Theorem 1.

Our proof relies heavily on Theorem 2. However, we note here that Theorem 3 can be proved by another road, that combines Corollary 5 and a lemma on generalized inverses due to Hartwig and Luh ([19] Lemma 3).

The isomorphism theorem states that $R/l(a) \cong Ra$ (as left modules) for any element a of a ring R (whether or not it has an identity), where as usual $l(a)$ denotes the left annihilator of a : $l(a) = \{r \in R, ra = 0\}$. A fundamental result due to Ehrlich[11], see also [28], states that R is a unit-regular rings if and only if it is regular and the dual isomorphism theorem $R/Ra \cong l(a)$ holds for all $a \in R$. In [28], Nicholson and Sánchez Campos call an element with the property $R/Ra \cong l(a)$ left morphic (The ring itself is called a left morphic ring if every element is left morphic), and give a short proof of the result elementwise:

Proposition 5 (Ehrlich, Nicholson, Sánchez Campos) Let R be a unital ring. Then a is unit-regular if and only if it is regular and morphic.

We conclude the article by proving that group-regular rings are morphic. We first note that idempotents are always left morphic:

Lemma 8 Let $e \in E(R)$, R general ring. Then e is morphic.

Proof Define $\phi : R \setminus Re \rightarrow l(e)$ by $\phi : x + Re \mapsto (x - xe)$. Then ϕ is well-defined and invertible ($x - y \in Re \Leftrightarrow x - y = (x - y)e \Leftrightarrow x - xe = y - ye$). Thus e is left morphic. It is right morphic by duality.

Theorem 4 *Let R be a group-regular ring. Then:*

1. *for all $e, f \in E(R)$, if $e \sim f$ then $l(e) \cong l(f)$;*
2. *for all $a \in R$, a is morphic.*

Proof

1. Let $e, f \in E(R)$ be isomorphic idempotents, and let T be a unitization of R . Then $1 - e \sim 1 - f$ by Theorem 3 and exists $c, d \in T$ such that $1 - e = cd, 1 - f = dc, cdc = c, dcd = d$. We consider the map $\phi : l(e) \rightarrow l(f)$ (left annihilators in R) defined by $x \mapsto xc$. It is well defined since R is a ideal of T , and $xe = 0 \Rightarrow x = xcd \Rightarrow xcf = xc - xc(1 - f) = xc - xcdc = 0$. It is bijective with reciprocal $y \mapsto yd$ and a morphism of left modules since it acts by right multiplication. Thus $l(e) \cong l(f)$.
2. Let now $a \in R$ and assume that a is group-dominated by $x^\# \in R^\#$. Then $e = ax$ and $f = xa$ are isomorphic idempotents such that $Ra = Rf$ and $l(e) = l(a)$. Indeed, as $a = ex^\# = x^\#f$ then $Rf = Rxx^\#f \subseteq Ra = Rx^\#f \subseteq Rf$, and dually ($\forall z \in R$) $za = 0 \Leftrightarrow zex^\# = 0 \Leftrightarrow ze = 0$ (since $ex^\#x = e$). Finally

$$R/Ra = R/Rf \cong l(f) \cong l(e) = l(a)$$

and a is left morphic. It is right morphic by dual arguments.

References

1. G. Abrams and K.M. Rangaswamy. Regularity conditions for arbitrary Leavitt path algebras. *Algebr. Represent. Theory*, 13(3):319–334, 2010.
2. PN Anh and L Márki. Morita equivalence for rings without identity. *Tsukuba journal of mathematics*, 11(1):1–16, 1987.
3. P. Ara. Extensions of exchange rings. *J. Algebra*, 197(2):409–423, 1997.
4. G. M. Bergman. The diamond lemma for ring theory. *Adv. Math.*, 29(2):178–218, 1978.
5. Gary F Birkenmeir. Idempotents and completely semiprime ideals. *Communications in Algebra*, 11(6):567–580, 1983.
6. G. Calugareanu. On unit-regular rings. *Preprint*, 2010.
7. V.P. Camillo and D. Khurana. A characterization of unit regular rings. *Comm. Algebra*, 29(5):2293–2295, 2001.
8. H. Chen. Extensions of Unit-Regular rings. *Comm. Algebra*, 32(6):2359–2364, 2004.
9. Huanyin Chen and Miaosen Chen. On unit-regular ideals. *New York J. Math*, 9:295–302, 2003.
10. G. Ehrlich. Unit-regular rings. *Port. Math.*, 27(4):209–212, 1968.
11. G. Ehrlich. Units and one-sided units in regular rings. *Trans. Amer. Math. Soc.*, 216:81–90, 1976.
12. C. Faith and Y. Utumi. On a new proof of Litoff's theorem. *Acta Math. Hungar.*, 14(3-4):369–371, 1963.
13. J. Fountain and M. Petrich. Completely 0-simple semigroups of quotients. *J. Algebra*, 101(2):365–402, 1986.
14. N. Funayama. Imbedding a regular ring in a regular ring with identity. *Nagoya Math. J.*, 27(1):61–64, 1966.
15. K.R. Goodearl. *Von Neumann regular rings*. Krieger Pub Co, 1991.
16. V. Gould. Semigroups of left quotients: existence, straightness and locality. *J. Algebra*, 267(2):514–541, 2003.
17. D. Handelman. Perspectivity and cancellation in regular rings. *J. Algebra*, 48(1):1–16, 1977.
18. R.E. Hartwig. How to partially order regular elements. *Math. Japon.*, 25:1–13, 1980.
19. R.E. Hartwig and J. Luh. A note on the group structure of unit regular ring elements. *Pacific J. Math.*, 71(2):449–461, 1977.

20. R.E. Hartwig and J. Luh. On finite regular rings. *Pacific J. Math.*, 69(1):73–95, 1977.
21. Irving Kaplansky. *Fields and rings*. University of Chicago Press, 1972.
22. D. Khurana and T.Y. Lam. Clean matrices and unit-regular matrices. *J. Algebra*, 280(2):683–698, 2004.
23. T.Y. Lam and W. Murray. Unit regular elements in corner rings. *Bull. HongKong Math. Soc.*, 1(1):61–65, 1997.
24. X. Mary. On (E, \tilde{H}_E) -abundant semigroups and their subclasses. *Semigroup Forum*, 94(3):738–776, 2017.
25. P. Menal and J. Moncasi. Lifting units in self-injective rings and an index theory for Rickart c^* -algebras. *Pacific J. Math.*, 126(2):295–329, 1987.
26. H. Mitsch. A natural partial order for semigroups. *Proc. Amer. Math. Soc.*, 97(3):384–388, 1986.
27. K.S.S. Nambooripad. The natural partial order on a regular semigroup. *Proc. Edinb. Math. Soc.*, 23(3):249–260, 1980.
28. WK Nicholson and E Sánchez Campos. Rings with the dual of the isomorphism theorem. *Journal of Algebra*, 271(1):391–406, 2004.
29. W.K. Nicholson and Y. Zhou. Clean general rings. *J. Algebra*, 291(1):297–311, 2005.
30. P.P. Nielsen and J. Ster. Connections between unit-regularity, regularity, cleanness, and strong cleanness of elements and rings. *Trans. Amer. Math. Soc.*, 2017.
31. T. Özdin. On locally unit-regular Leavitt path algebras. *Commun. Fac. Sci. Univ. Ank.*, 67(2):11–18, 2018.
32. Vladimir Rakočević. On harte’s theorem for regular boundary elements. *Filomat*, pages 899–910, 1995.
33. L.N. Vaserstein. Bass’s first stable range condition. *J. Pure Appl. Algebra*, 34(2-3):319–330, 1984.