







BOOK of ABSTRACTS

ISBN: 978-88-8080-224-2 - DOI: https://doi.org/10.48257/ACBA01



Published by

Institute of Sciences of Food Production-National Research Council (CNR-ISPA)

Editor in Chief

Dr. Antonio F. Logrieco

Editorial and Organising Board

Dr. Antonio Moretti
Institute of Science of Food Production (CNR-ISPA), Italy
Dr. Giancarlo Perrone
Institute of Science of Food Production (CNR-ISPA), Italy
Dr. Hayrettin Ozer
TUBITAK MRC Food Institute, Turkey
Dr. Giuseppina Avantaggiato
Institute of Science of Food Production (CNR-ISPA), Italy
Dr. Vincenzo Lippolis
Institute of Science of Food Production (CNR-ISPA), Italy

Technical editors

Domenico GENCHI, Martina LOI, Dalia LEVITA, Alessandra VILLANI Institute of Sciences of Food Production, (CNR-ISPA), Italy

Digital book by

Domenico GENCHI Institute of Sciences of Food Production, (CNR-ISPA), Italy

ISBN digital version: 978-88-8080-221-1 ISBN printed version: 978-88-8080-224-2

DOI: https://doi.org/10.48257/ACBA01

Session 1 Oral 4

BIODIVERSITY OF TOXIGENIC FUNGI IN THE FOOD CHAIN AND THE CO-OCCURRENCE OF MYCOTOXINS.

<u>VENÂNCIO A.^{1*}</u>, VALE DIAS T.¹, GUIMARÃES A.¹, CALADO T.¹ and RODRIGUES P.².

¹CEB - Centre of Biological Engineering, University of Minho, 4710-057 Braga, Portugal; ²Centro de Investigação de Montanha (CIMO); Instituto Politécnico de Bragança, Campus de Santa Apolónia, 5300-253 Bragança, Portugal avenan@deb.uminho.pt

Filamentous fungi are ubiquitous in Nature and may be found in any food crop, either in the field or during storage. With the exponential growth of the population, there is a worldwide challenge to reduce food losses. Food contamination by fungi causes great economic costs and several health threats due to the toxicity and pathogenicity of some species. The biodiversity of this community, and the dynamics of the water activity, temperature and availability of carbon sources evolution, during the growing season, as well as in storage, determines the competitiveness of each species against other co-occurring fungal species.

Some of these fungi are capable of producing a wide range of secondary metabolites – mycotoxins, which may accumulate in food chain, be resilient during food processing, and persist in the final food product. Contamination of food products with fungi is frequent, affecting food security and food safety. The ability of the different fungal species to compete under available conditions will influence the cocktail of mycotoxin that may occur.

The study of the fungal biodiversity have been mainly carried out by surveying culturable strains (culture-dependent approaches), but recently metagenomics approaches have been used, enabling the possibility of spotting mycotoxigenic fungal strains, that are not easy to detect and isolate by conventional means.

In this presentation, the biodiversity of fungal species in the food chain will be discussed, under a perspective of the co-occurrence of mycotoxins.

This study was supported by the Portuguese Foundation for Science and Technology (FCT) under the scope of the strategic funding of UIDB/04469/2020 (CEB) and UID/AGR/00690/2019 (CIMO) units. We also thank the FCT for the Ph.D. scholarship given to Teresa Vale Dias (2020.05849.BD).