

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

06 January 2022 (06.01.2022)



(10) International Publication Number

WO 2022/003655 A1

(51) International Patent Classification:

A61K 8/64 (2006.01) A61Q 15/00 (2006.01)

A61K 8/11 (2006.01)

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))
- in black and white; the international application as filed contained color or greyscale and is available for download from PATENTSCOPE

(21) International Application Number:

PCT/IB2021/056011

(22) International Filing Date:

05 July 2021 (05.07.2021)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

116561 03 July 2020 (03.07.2020) PT

20206292.3 06 November 2020 (06.11.2020) EP

(71) Applicant: UNIVERSIDADE DO MINHO [PT/PT];

Largo Do Paço, 4704-553 Braga (PT).

(72) Inventors: CAVACO PAULO, Artur Manuel; Rua Garcia De Orta, 49, 4715 - 191 Braga (PT). GOMES

GONÇALVES, Filipa Daniela; Rua Cândido De Sousa N° 11, 2°dir, 4710-503 Braga (PT). ARAÚJO MAGAL-

HÃES RIBEIRO, Artur Jorge; Estrada De Dadim N°14, 4715-266 Braga (PT). PEREIRA MARINHO DA SILVA,

Carla Manuela; Rua Da Fé, N° 488 5° Andar, Guimarães (PT).

(74) Agent: PATENTREE; Association 669, Rua de Salazares,

842, Edf. NET, 4149-002 Porto (PT).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: FRAGRANCE RELEASE MECHANISM, METHOD AND USES THEREOF

(57) Abstract: The present disclosure relates to a release composition comprising a protein and an active agent, wherein the active agent is released when in presence of an electrolyte solution. A kit and an article comprising the release composition of the present-subject matter are also encompassed.



WO 2022/003655 A1

FRAGRANCE RELEASE MECHANISM, METHOD AND USES THEREOF

TECHNICAL FIELD

[0001] The present disclosure relates to a mechanism of adsorption and dissociation involving fragrance molecule and an odorant-binding protein (OBP-I) regulated simultaneously by human sweat and temperature. This system can be used in cosmetic formulations where the fragrances are released upon presence of sweat at body temperature.

BACKGROUND

[0002] Odorant-binding proteins (OBPs) are small water-soluble proteins, belonging to lipocalins superfamily.¹⁻² They are responsible to transport hydrophobic odorous molecules, called odorants, in their calyx-shaped cavity, across the aqueous mucus barrier towards the olfactory receptors, where a cascade of transduction signal is traduced in the brain's interpretation.³⁻⁴ These proteins are also described as involved in removing odorants from the olfactory receptors after their stimulation.⁵⁻⁶

[0003] Several mammalian odorant-binding proteins have been identified and some of them isolated from nasal mucus such as bovine, pig, boar, panda, mice, rats and humans.⁷⁻¹³

[0004] DNA sequences of mammalian OBPs present low similarity: porcine OBP (OBP-I) and human OBP (hOBP_{Ila}) present only 13.9% of DNA sequence similarity; OBP-I and bovine OBP present 42.7%.⁴ Despite wide genetic variability between OBPs from different mammalian species, lipocalin members present few characteristic signatures that allow their identification as the case of the conserved tertiary structure, presenting a β -barrel structure composed by eight β -strands linked by seven loops and connected to a short α -helix close to the C-terminus and a ninth β -strand followed by the

disordered C-terminal tail.¹⁴⁻¹⁶ The structure of OBPs is highly stable and resistant to degradation by temperature, organic solvents, pH variation, or proteolytic digestion.¹⁷⁻¹⁸ ¹⁹ The FT-IR spectra for porcine OBP revealed a structure exceptionally stable to thermal denaturation (up to 80 °C), particularly in the presence of a ligand.²⁰ Furthermore, vertebrate OBPs show capacity to reverse the unfolding of protein even after denaturation.²⁰

[0005] Human body produces unpleasant odours associated with stress, anxiousness, nervousness and physical exercise.²¹ To prevent or reduce their occurrence, antibacterial agents and fragrances are commonly added to cosmetic formulations. However, drawbacks related with the limited effect against different odours and with the residual amount of these deodorants are detected in clothing and skin.²²

[0006] OBPs have affinity for several molecules associated with odorific feeling. All those molecules are volatile and detected by OBPs at very low concentrations, being a system highly sensitive. The fast responsive time of the OBPs and the high stability of these proteins create an excellent biological element as biosensor for detection of the dangerous substances and pathogens, pesticides and drugs present in food or water¹⁸,²³ as well as the potential use as deodorizer and medical diagnostics.²⁴⁻²⁵

[0007] Odorific molecules can be associated with pleasant or unpleasant feelings. OBP have affinities for all molecules associated with odors.³⁻⁴ Fragrances are compositions containing odorific molecules with pleasant feeling.

[0008] The use of 1-aminoanthracene (1-AMA) as odour model molecule provides a capacity to measure the binding capacity of odorant-binding protein, by fluorescence assay. The free 1-AMA and 1-AMA bound to pig OBP-I can be quantified measuring the fluorescence with excitation wavelength at 295 nm. The maximum wavelength of 1-AMA bound to OBP-I is shifted from 537 nm to 481 nm.²⁰ The non-fluorescence odorant can be measured by competitive assays or by gas chromatography–mass spectrometry.

[0009] The following works already reported the interaction between OBPs and odour model molecules, as well as with lipidic structures such as liposomes.

[0010] Filipa Gonçalves *et al* (2018) “Two Engineered OBPs with opposite temperature-dependent affinities towards 1-aminoanthracene” mentions two recombinant proteins based on pig OBP sequence (i) truncated OBP (tOBP-F44A/F66A) obtained from the deletion of the first 16 residues of the N-terminal and the replacement of two phenylalanine residues at the binding pocket by alanine residues (F44A and F66A), and (ii) OBP::GQ₂₀::SP-DS3 resulted of the fusion of OBP-I with a spacer of 20 repetitions of glycine-glutamine residues and the anchor peptide SP-DS3.³⁴ Experimental and molecular modelling data showed that 1-AMA model ligand binds preferentially to tOBP-F44A/F66A at 25 °C while ligand binds to OBP::GQ₂₀::SP-DS3 favourably at 37 °C.³⁴

[0011] Filipa Gonçalves *et al* (2018) “OBP fused with cell-penetrating peptides promotes liposomal transduction” report the fusion of porcine OBP with cell-penetrating peptides (CPPs, e.g. TAT, Pep-1 and pVEC). The study revealed different efficiencies on 1-AMA transduction into liposomes.³⁰

[0012] Filipa Gonçalves *et al* (2018) “1-Aminoanthracene Transduction into Liposomes Driven by Odorant-Binding Protein Proximity” discloses the design of two fusion proteins based on pig OBP fused with anchor peptide SP-DS3³² in absence and presence of a spacer (GQ₂₀). This work demonstrated that the 1-AMA transduction into liposomes is driven by proximity of protein anchored to liposomal membrane (advantage for absence of spacer).³³

[0013] Filipa Gonçalves *et al* (2019) “Release of Fragrances from Cotton Functionalized with Carbohydrate-Binding Module Proteins” discloses the design of fusion protein based on pig OBP fused with a spacer (GQ₂₀) and a carbohydrate binding module (CBM). The work demonstrated the affinity of protein to one fragrance (β -citronellol) and the release of this fragrance from cotton in presence of sweat.²⁵ Regardless, the release capacity in the presence of sweat is inferior as compared to the native OBP.

[0014] Alessandro Capo *et al* (2018) “The porcine odorant-binding protein as molecular probe for benzene detection” discloses pig odorant-binding protein to be used as probe for benzene detection in atmosphere, through fluorescence assay.²⁸

[0015] Nunzio Cennamo *et al* (2015) “Easy to Use Plastic Optical Fiber-Based Biosensor for Detection of Butanal” reports the detection of butanal (20-1000 μM) by porcine odorant-binding protein through competitive assay. This aldehyde is very toxic, exhibiting high risks for human health like cytotoxicity and cancer. The authors describe an optical biosensor to detect butanol in liquid samples.²⁷

[0016] Carla Silva *et al* (2014) “Odorant binding proteins: a biotechnological tool for odour control” discloses the application of porcine odorant-binding protein for release of fragrances from a cotton fabric to mask smoke odour. The authors confirmed the functionalization of OBP on fabrics. They tested only the release of one fragrance from textile. Contrary to the work of Silva *et al.*, the present disclosure includes the release mechanism of different fragrances or other molecules as response of human perspiration with upper efficiency. Additionally, the subject-matter of the present disclosure is suitable for use in textile and cosmetic fields.

[0017] Paolo Pelosi *et al* (2014) “Structure and biotechnological applications of odorant-binding proteins” discloses the possibility of OBPs to be used as a sensor to detect volatile and slow release of odorant molecules.

[0018] Lei Han *et al* (2014) “Operating Mechanism and Molecular Dynamics of Pheromone-Binding Protein ASP1 as Influenced by pH” indicates pheromone-binding protein ASP1 as binds odorants at low pH and the dissociation respond to pH change. The authors describe the benefit of this research in biotechnology and agriculture. The results were determined by molecular docking and dynamics simulations.

[0019] Alberto Mazzini *et al* (2007) “Dissociation and unfolding of bovine odorant binding protein at acidic pH” discloses the structure of bovine OBP at neutral and acidic pH, by molecular simulation.

[0020] Mariella Parisi *et al* (2003) “Unfolding and refolding of porcine odorant binding protein in guanidinium hydrochloride: equilibrium studies at neutral pH” discloses the denaturant effect of guanidinium hydrochloride, a well-known chaotropic agent, in

folding/unfolding of protein. The aim of this fundamental study was to understand the structure of the OBP protein, in particular its unfolding and refolding process.

[0021] Document WO 0123890A1 discloses a detector array based on sensing elements within a solid support with use for clinical samples or cell extracts, in gaseous state. It is an immunoassay utilizing viral peptides.

[0022] Document EP0335654A3 discloses the method for gene isolation of odorant-binding protein from rat and a protein production method.

[0023] Document WO2001012806A3 described OBPII as a fixer of hydrophobic ligands such as odours that can be used for personal hygiene, agri-food system and nutritional and therapeutic uses.

[0024] These facts are disclosed in order to illustrate the technical problem addressed by the present disclosure.

GENERAL DESCRIPTION

[0025] The present disclosure is related to adsorption and dissociation mechanism of native odorant-binding protein (OBP-I), in particular pig OBP-I (SEQ. ID NO. 1), and OBP fused with linker GQ 20x and KP peptide (OBP::GQ₂₀::KP, SEQ. ID NO. 21). These proteins have a negative charge of approximately -20 (pH 7.4) due to the high content in aspartic acid and glutamic acid residues. The isoelectric point value of these proteins are between 4.08 and 4.65.²⁶

[0026] In an embodiment, the solution here disclosed can have a high impact in human social life that are associated with perspiration issues. The system has several advantages including the use of bioinspired cosmetic bioingredients (green solutions) without damaging for the ecosystems.

[0027] The mechanism here disclosed divulge that odorant-binding protein, in particular porcine odorant-binding protein (OBP-I), presents high affinity (adsorption) to fragrances in 50 mM Tris-HCl pH 7.5 buffer, at 37 °C. The affinity constant (K_a) of OBP-I

was of $4.00 \pm 0.03 \mu\text{M}$. On the other hand, OBP-I presents a reverse mechanism, i.e., the dissociation of fragrance from its binding pocket with reduced K_a ($0.20 \pm 0.02 \mu\text{M}$) when in exposition of perspiration (sweat), even at different pH (range of 4.0-8.5, **Table 2**). Similarity, OBP::GQ₂₀::KP presents high affinity to fragrances in buffer, at 37 °C ($K_a = 4.00 \pm 0.04 \mu\text{M}$) that is reduced in presence of an electrolyte solution, such as sweat ($K_a = 0.59 \pm 0.01 \mu\text{M}$). Therefore, OBPs presents reduced affinity when in contact with perspiration, releasing the fragrance in this condition.

[0028] Surprisingly, OBP::GQ₂₀::KP (SEQ ID NO. 21) showed 6.8x more fragrance release in presence of sweat versus the presence of buffer. These values are superior to the values reported in state of art, in particular to the values reported for OBP::GQ₂₀::CBM (SEQ ID NO. 22), where the release mechanism showed 1.3x release of fragrance in presence of sweat.²⁵

[0029] The adsorption and dissociation mechanism of porcine odorant-binding protein can be done in a repeated manner.

[0030] Human sweat can be used as a trigger to release/dissociate a fragrance from OBP-I. Therefore, the subject-matter of the present disclosure can be used in skin care products as well as in textile items, in particular clothes.

[0031] In an embodiment, the present disclosure relates to a protein with an amino acid sequence similar to mammals odorant-binding proteins to be incorporated in formulations for cosmetic or textile applications.

[0032] In an embodiment, the native form of odorant-binding protein may be from pig, human, dog, cat, rat, mouse, cow, boar, panda, Chinese hamster, Meishan pig, Guinea pig, Tibetan pig, horse, dolphin and chimpanzee.

[0033] In an aspect of the present disclosure, applications of the present subject-matter may be based on the release of odorant molecules from odorant-binding protein, triggered by electrolyte solutions at body temperature.

[0034] In an embodiment, the electrolyte solution refers to a solution with a NaCl concentration higher than 9.5 grams/L, in particular to a solution with a NaCl concentration ranging from 9.5 to 45 grams/L. Preferably, the electrolyte solution is human sweat, pet sweat, salty water or micellar water.

[0035] In an embodiment, human sweat may comprise water, lactic acid, urea and minerals, such as sodium, potassium, calcium, and magnesium.

[0036] In an embodiment, cosmetic applications may be for skin and hair care. Skin care applications may be related with OBPs formulated in specialty formulations for skin creams, lipsticks, lips creams and face mask powders, face and body creams, skin clarifiers, primers and foundations.

[0037] In a further embodiment, hair care applications may be related with OBPs formulated in eyelash mascaras, hair shampoos, hair serum, hairs masks, hair conditioners, or hair coloration creams.

[0038] The present disclosure relates to a release composition comprising an isolated or artificial protein with at least 90% of identity with an amino acid sequence selected from the following list: SEQ. ID NO. 1, SEQ. ID NO. 2, SEQ. ID NO. 3, SEQ. ID NO. 4, SEQ. ID NO. 5, SEQ. ID NO. 6, SEQ. ID NO. 7, SEQ. ID NO. 8, SEQ. ID NO. 9, SEQ. ID NO. 10, SEQ. ID NO. 11, SEQ. ID NO. 12, SEQ. ID NO. 13, SEQ. ID NO. 14, SEQ. ID NO. 15, SEQ. ID NO. 16, SEQ. ID NO. 17, SEQ. ID NO. 18, SEQ. ID NO. 19, SEQ. ID NO. 20, SEQ. ID NO. 21, or mixtures thereof, preferably 91% identical, 92% identical, 93% identical, 94% identical, 95% identical, 96% identical, 97% identical, 98% identical, 99% identical or identical; an active agent selected from a list comprising a deodorizing agent, a natural essence, a fragrance, a moisturizing agent, or mixtures thereof; wherein the active agent is bounded to the protein; and wherein the protein releases the active agent in the presence of an electrolyte solution, at a temperature between 10-60 °C.

[0039] In an embodiment, the release composition comprises an isolated or artificial unmodified protein with at least 90% of identity with an amino acid sequence selected from the following list: SEQ. ID NO. 1, SEQ. ID NO. 2, SEQ. ID NO. 3, SEQ. ID NO. 4, SEQ.

ID NO. 5, SEQ. ID NO. 6, SEQ. ID NO. 7, SEQ. ID NO. 8, SEQ. ID NO. 9, SEQ. ID NO. 10, SEQ. ID NO. 11, SEQ. ID NO. 12, SEQ. ID NO. 13, SEQ. ID NO. 14, SEQ. ID NO. 15, SEQ. ID NO. 16, SEQ. ID NO. 17, SEQ. ID NO. 18, SEQ. ID NO. 19, SEQ. ID NO. 20, SEQ. ID NO. 21, or mixtures thereof, preferably 91% identical, 92% identical, 93% identical, 94% identical, 95% identical, 96% identical, 97% identical, 98% identical, 99% identical or identical.

[0040] Methods for the alignment of sequences for comparison are well known in the art, such methods include GAP, BESTFIT, BLAST, FASTA and TFASTA. GAP uses the algorithm of Needleman and Wunsch ((1970) J Mol Biol 48: 443-453) to find the global (over the whole the sequence) alignment of two sequences that maximizes the number of matches and minimizes the number of gaps. The BLAST algorithm (Altschul et al. (1990) J Mol Biol 215: 403-10) calculates percent sequence identity and performs a statistical analysis of the similarity between the two sequences. The software for performing BLAST analysis is publicly available through the National Centre for Biotechnology Information (NCBI). Global percentages of similarity and identity may also be determined using one of the methods available in the MatGAT software package (Campanella et al., BMC Bioinformatics. 2003 Jul 10; 4:29. MatGAT: an application that generates similarity/identity matrices using protein or DNA sequences). Minor manual editing may be performed to optimise alignment between conserved motifs, as would be apparent to a person skilled in the art. The sequence identity values, which are indicated in the present subject matter as a percentage were determined over the entire amino acid sequence, using BLAST with the default parameters.

[0041] In an embodiment, the composition of the present subject-matter comprises 0.01 to 5000 μM of the unmodified protein.

[0042] In an embodiment, the composition of the present subject-matter comprises 0.01 to 5000 μM of the protein.

[0043] In another embodiment, the release composition comprises 0.1 μM to 2 M of active agent, preferably 0.2 μM to 1 M.

[0044] In an embodiment, the protein has an affinity constant of 1-4.5 μM to the active agent, in water or buffer solutions, preferably Tris-HCl, phosphate solution, and/or phosphate buffered saline. In a further embodiment, the affinity constant of the protein to the active agent ranges between 0.1-0.99 μM in the electrolyte solution, preferably in sweat.

[0045] The present disclosure relates to a fragrance release composition comprising: 0.01 to 5000 μM of an isolated or artificial protein with at least 90% of identity with an amino acid sequence selected from the following list: SEQ. ID NO. 1, SEQ. ID NO. 2, SEQ. ID NO. 3, SEQ. ID NO. 4, SEQ. ID NO. 5, SEQ. ID NO. 6, SEQ. ID NO. 7, SEQ. ID NO. 8, SEQ. ID NO. 9, SEQ. ID NO. 10, SEQ. ID NO. 11, SEQ. ID NO. 12, SEQ. ID NO. 13, SEQ. ID NO. 14, SEQ. ID NO. 15, SEQ. ID NO. 16, SEQ. ID NO. 17, SEQ. ID NO. 18, SEQ. ID NO. 19, SEQ. ID NO. 20, SEQ. ID NO. 21 or mixtures thereof; 0.1 μM to 2 M of an active agent selected from a list comprising a deodorizing agent, a natural essence, a fragrance, a moisturizing agent, or mixtures thereof; wherein the active agent is bounded and/or entrapped to the protein; and wherein the protein releases the active agent in the presence of an electrolyte solution, at a temperature between 10-70 $^{\circ}\text{C}$, preferably 10-60 $^{\circ}\text{C}$; wherein the affinity constant of the protein to the active agent, in water, ranges between 1-4.5 μM ; wherein the active agent has a molecular weight from 20 to 1000 g/mol; wherein the electrolyte solution is sweat, salty water or micellar water.

[0046] In an embodiment, the active agent has a molecular weight between 20 to 1000 g/mol, preferably 75-300 g/mol. In a further embodiment, the active agent is a fragrance molecule. In a yet further embodiment, the bioactive agent comprises a functional group selected from aromatic, aldehyde or alcohols. In a yet further embodiment, the active agent is a fragrance molecule, selected from a list comprising the molecules listed in Table 1.

[0047] Table 1 - List of fragrances and their properties.

CAS#	Name	Odor description	Chemical Family	MW (g/mol)
85213-22-5	2-acetyl-1-pyrroline	roasted/bread	ketone	111,14
8000-41-7	α -terpineol	lilac	alcohol/terpene	154,25
502-99-8	β -ocimene	sweet herbal	hydrocarbon/terpene	136,24
140-11-4	benzyl acetate	strawberry/pear	ester	150,17
123-86-4	butyl acetate	banana	ester	116,16
76-22-2	camphor	camphora	ketone/terpene	152,24
6485-40-1	carvone	mint	ketone/terpene	150,22
5392-40-5	citral	lemon/citrus	aldehyde/terpene	152,24
106-22-9	citronellol	citronella/rose-like	alcohol/terpene	156,27
91-64-5	coumarin	sweet vanilla/pleasant	lactone/aromatic	146,15
431-03-8	diacetyl	buttery	ketone	86,09
97-53-0	eugenol	cloves	aromatic alcohol	164,20
6413-10-1	fructone	apple	ester	174,19
706-14-9	gamma decalactone	coconut	lactone	170,25
104-61-0	gamma nonalactone	peach/fruity	lactone	156,23
106-24-1	geraniol	floral/sweet rose	terpene	154,24
24851-98-7	hedione	floral/jasmine	ester	226,32
123-92-2	isoamyl acetate	pear/banana	ester	130,19
67920-63-2	lilac aldehyde	floral/lilac	aldehyde/terpene	168,24
5989-27-5	limonene	citric	terpene	136,23
126-91-0	linalool	lavender/bergamot	terpene	154,25
55066-48-3	mefrosol	floral/rose	alcohol	178,27
2216-51-5	menthol	peppermint	alcohol	156,26
623-42-7	methyl butyrate	apple/pineapple	ester	102,13
123-35-3	myrcene	herbal/woody	terpene	136,24
80-56-8	pinene	pine	terpene	136,24
357650-26-1	pomarose	plums/apples rose	ketone	166,26
89-82-7	pulegone	peppermint	ketone/terpene	152,24
65113-99-7	sandalore	sandalwood	alcohol	210,36
121-33-5	vanilin	vanilla	aldehyde/aromatic	152,15

[0048] In an embodiment, the release of the active agent occurs during 30 seconds to 24 h.

[0049] In an embodiment, electrolyte solution has a pH between 4.0-8.5. In a further embodiment, the electrolyte solution is sweat, salty water or micellar water, preferably human sweat or pet sweat.

[0050] In an embodiment, the unmodified protein is stable in polar and non-polar solvents, including methanol, butanol, benzene, ethanol and undecanol as well as buffer solutions, preferably Tris-HCl, phosphate, or phosphate buffered saline (PBS)). In a further embodiment, the unmodified protein is also stable in temperatures between 18-70 °C, preferably 18-60 °C, and in a pH range of 4.0-10.0.

[0051] In an embodiment, the protein is stable in polar and non-polar solvents, including methanol, butanol, benzene and undecanol as well as buffer solutions (Tris-HCl, phosphate, PBS). In a further embodiment, the protein is also stable in temperatures between 18-70 °C, preferably 18-60 °C, and in a pH range of 4.0-10.0.

[0052] In an embodiment, the release of the active agent from the protein occurs at 20-40 °C.

[0053] In an embodiment, the composition further comprises glycerol, erythritol, arabitol, sorbitol, mannitol, xylitol, mannisdomannitol, glucosylglycerol, glucose, fructose, sucrose, trehalose, isofluoroside, dextrans, levans, polyethylene glycol, salts of chloride, citrate, sulfates, acetate or phosphates, or mixtures thereof.

[0054] An aspect of the present disclosure comprises a kit or article comprising the composition described in the present subject-matter. In an embodiment, the article comprising said composition can be selected from a list comprising fabric, textiles, fibers, clothes, scarfs, hats, gloves, socks and turbans, shoes, insoles, bags, handbags, detergents, creams, lotions, foams, perfumes, softeners, aerosols, deodorants, lipsticks, lip creams, face mask powders, face and body creams, skin clarifiers, primers, foundations, hair shampoos, hair serum, hairs masks, hair conditioners or hair coloration creams.

[0055] The present disclosure also relates to the use of the fragrance release composition described in the present subject-matter in cosmetics, preferably as a cosmetic agent/composition, more preferably skin care or hair care; as well as the use of said composition in the textile industry.

[0056] The present disclosure also relates to the use of the fragrance release composition as a deodorant agent/composition.

BRIEF DESCRIPTION OF THE DRAWINGS

[0057] The following figures provide preferred embodiments for illustrating the disclosure and should not be seen as limiting the scope of invention.

[0058] **Figure 1:** Schematic representation Sodium dodecyl sulfate and polyacrylamide (SDS-PAGE) gel electrophoresis under reducing conditions (A), and circular dichroism (CD) spectrum (B) of native pig odorant-binding protein. Mw: Precision Plus Protein™ standards (BioRad).

DETAILED DESCRIPTION

[0059] The present disclosure concerns the method of manufacture of a fragrance/protein complex and the adsorption and dissociation mechanism of native odorant-binding protein (OBP-I), in particular pig OBP-I, and native protein fused with GQ₂₀ and KP peptide. This mechanism involves the release of active agents, preferably odorific molecules (or fragrances), in response to human perspiration.

[0060] The present disclosure presents high impact in textile and cosmetic fields, particularly in the release of fragrances from lotion and cream base products as well as textile fabrics, awarding a green solution.

[0061] In an embodiment, the produced odorant-binding protein and fragrance/protein complex are very soluble and stable at different solvents (like methanol, butanol, benzene and undecanol) and different range of temperature (18-60 °C) and different range of pH (4.0-10.0).^{12, 27-29}

[0062] In a further embodiment, the mixture of odorific molecules and the OBP-I or OBP::GQ₂₀::KP proteins showed improved results regarding the affinity constants in a buffer solution and in sweat (**Table 2**). In particular, the lower K_a in sweat enforces the use of OBP proteins when sweat-triggered responses are envisaged. Notably, the adsorption and dissociation mechanism by the proteins is reversible. In response to human perspiration at different range of pH (pH 4.0-8.5) the disclosed proteins released more than 20x to 6.8x (OBP-I and OBP::GQ₂₀::KP, respectively) of odorific molecules compared with other fusion proteins based on porcine odorant-binding protein (Table 2). Thus, the release of fragrance occurs in response of perspiration and it is independent of pH of sweat of each human. Additionally, the fragrance release occurs at least during 0.5-24 h.

[0063] In an embodiment, a variety of active agents, including deodorizing agents, natural essences, fragrance agents, moisturizing agents, or mixtures thereof, can be used in the complex formation, allowing a wide range of cosmetic uses. In particular, pleasant odorific molecules (of different size and shape) showed great affinity to the odorant-binding protein, in particular porcine odorant-binding protein.

[0064] In an embodiment, the odorific molecules used in the present disclosure belong to different functional groups (aromatic, aldehyde, alcohols). In a further embodiment the odorific molecules comprise molecules with a molecular weight ranging from 20,00 to 1000,00 g/mol, and the concentration varies from 0.2-2000 μ M.

[0065] In an embodiment, odorific molecules have a molecular weight between 75 to 300 g/mol.

[0066] In an embodiment, the system efficiently responded to human perspiration, releasing fragrances over time. Importantly, the system did not respond against the water existent in human body, giving specificity and robustness to the subject-matter presented in this disclosure. Without this selectivity, i.e., if the system responded to water, the OBP protein would immediately release the odorific molecules when in contact with the skin.

Example:

[0067] In an embodiment, native porcine odorant-binding protein (OBP-I) and OBP-I fused with a spacer glycine-glutamine, repeated 20x (GQ₂₀) and with a carbohydrate-binding module (OBP::GQ₂₀::CBM) were cloned in plasmid pET28a and transformed into *Escherichia coli* BL21(DE3). The proteins were expressed and purified through Nickel magnetic beads with specificity to His-tag present in the protein's N-terminal. 10 µM of protein were loaded on sodium dodecyl sulfate and polyacrylamide (SDS-PAGE) gel electrophoresis under reducing conditions. The same concentration was used to determine the structure of proteins by circular dichroism (CD) spectroscopy.

[0068] The protein purified in laboratory reveals a high level of purity (observed by SDS-PAGE gel, Figure 1-A) and secondary structure in barrel, a consequence of the presence of a high number of β-sheets, as analysed by circular dichroism spectroscopy (Figure 1-B).

[0069] In an embodiment, the pure odorant-binding protein was lyophilized and used in further procedures. To determine the affinity of the odorant-binding protein to odour molecules, 1-aminoanthracene (1-AMA) was used as an odour model molecule. Increased concentrations of fluorescent ligand model 1-aminoanthracene (1-AMA) were added to 1 µM of protein, and the formation of ligand/protein complex was quantified after 1 h at 37 °C by fluorescence emission at 481 nm (excitation at 295 nm).³⁰ Dissociation constant (K_d) was determined from a plot of fluorescence intensity versus concentration of ligand, obtained with a standard non-linear regression method, described in Malpeli et al. (1998).³¹ The affinity behaviour of protein in presence of sweat solution was performed through a competitive fluorescence assay. Here, 1 µM of protein was mixed with 2 µM of 1-AMA and incubated at 37 °C for 1 h. After this period, increased volumes of sweat solution were added to the complex and incubated at same conditions.

[0070] In an embodiment, the sweat solution was prepared as indicated in AATCC method 15-2009 "Colorfastness to Perspiration". The pH of the prepared solution varied

between 4.0 and 8.5. In accordance with its composition, the sweat solution is also regarded as electrolyte solution.

[0071] In a further embodiment, fluorescence emission at 481 nm (excitation at 295 nm) was recorded and the dissociation constant (K_d) calculated. The association constant (K_a) was calculated by formula $K_d = 1/K_a$.

[0072] In an embodiment, the release of fragrance was quantified by gas chromatography-mass spectrometry (GC-MS). Increased concentrations of fragrance were used in different vials and the fragrance in headspace quantified performing the calibration curve (area of peak vs fragrance concentration). The fragrance was incubated with the odorant-binding protein at 37 °C. Sweat solution was added and the fragrance release determined after several periods of time (0.5-24 h) of perspiration exposition.

[0073] In an embodiment, as comparative data, porcine odorant-binding protein was fused with a spacer GQ₂₀ and a carbohydrate-binding module (CBM), as previously reported.²⁵ Through the fusion of CBM_{N1} (PDB ID 1ULP) of endoglucanase C from *Cellulomonas fimi*, the OBP has a specific affinity to cotton. The modified protein showed a high association constant ($K_a = 4.17 \pm 0.05 \mu\text{M}$) that decreased for $K_a = 3.16 \pm 0.02 \mu\text{M}$ when a sweat solution was added. Native OBP-I (SEQ ID No. 1) and OBP::GQ₂₀::KP (SEQ ID NO. 21) showed an association constant very similar ($K_a = 4.00 \pm 0.03 \mu\text{M}$) to the value obtained for OBP::GQ₂₀::CBM (SEQ ID NO. 22, $K_a = 4.17 \pm 0.05$). However, the addition of the sweat solution had a remarkable effect in the constant of affinity of the native protein ($K_a = 0.20 \pm 0.02 \mu\text{M}$) and of the OBP::GQ₂₀::KP ($K_a = 0.59 \pm 0.01 \mu\text{M}$). A high reduction of affinity was observed, 20x using native OBP (SEQ ID NO. 1) and 6.8x using OBP::GQ₂₀::KP (SEQ ID NO. 21), as compared with the value quantified for protein fused with CBM (SEQ ID NO 22), where a reduction of only 1.3x was verified ($K_a = 3.16 \pm 0.02$ **Table 2**). Thus, the release of fragrance by OBP-I and OBP fused with KP is evident in response of perspiration.

Table 2. Affinity constant (Ka) of native OBP-I (SEQ ID NO. 1) and fusion proteins based on OBP, SEQ ID NO. 21 and SEQ ID NO 22. SEQ ID NO 22 was used as comparative data. Values are the mean of 2 independent experiments at 37 °C.

Protein	Ka in buffer (μM)	Ka in sweat (μM)
Native porcine OBP (OBP-I) – SEQ ID NO 1	4.00 \pm 0.03	0.20 \pm 0.02
OBP::GQ ₂₀ ::KP – SEQ ID NO 21	4.00 \pm 0.04	0.59 \pm 0.01
OBP::GQ ₂₀ ::CBM (fusion protein) ²⁵ – SEQ ID NO 22	4.17 \pm 0.05	3.16 \pm 0.02

[0074] In an embodiment, the following protein sequences can be incorporated in different substrates in order to release fragrances in the presence of sweat. In this regard, substrates can be selected from a list comprising textiles, fabrics, skin care products, hair care products, among others.

List of protein sequences

[0075] The sequences of protein are described by one letter code of amino acids. The code is as follows:

One letter code	Amino acid
A	Alanine
C	Cysteine
D	Aspartic acid
E	Glutamic acid
F	Phenylalanine
G	Glycine
H	Histidine
I	Isoleucine
K	Lysine
L	Leucine
M	Methionine
N	Asparagine
P	Proline
Q	Glutamine

R	Arginine
S	Serine
T	Threonine
V	Valine
W	Tryptophan
Y	Tyrosine

Pig OBP (PDB ID 1DZK) – SEQ ID NO. 1

QEPQPEQDPFELSGKWITSYIGSSDLEKIGENAPFQVFMRSIEFDDKESKVYLNFFSKENGICEEFSLI
 GTKQEGNTYDVNYAGNNKFVVSYASETALIISNINVDEEGDKTIMTGLLGKGTDIEDQDLEKFKEVT
 RENGIPEENIVNIIERDDCPA

Human OBP_{Ila} (UniProt ID Q9NY56) – SEQ ID NO. 2

MKTLFLGVTLGLAAALSFTLEEEDITGTWYVKAMVVVDKDFPEDRRRPRKVSPVKVTALGGGNLEATF
 TFMREDRCIQKKILMRKTEEPGKFSAYGGRKLIYLQELPGTDDYVFYCKDQRRGGLRYMGKLVGR
 NPNTNLEALEEFKKLQVQHKGLSEEDIFMPLQTGSCVLEH

Human OBP_{Iib} (UniProt ID Q9NPH6) – SEQ ID NO. 3

MKTLFLGVTLGLAAALSFTLEEEDITGTWYVKAMVVVDKDFPEDRRRPRKVSPVKVTALGGGKLEATF
 TFMREDRCIQKKILMRKTEEPGKYSAYGGRKLMYQLPRRDHYIFYCKDQHHGGLLHMGLVGR
 NSDTNREALEEFKKLQVQHKGLSEEDIFPLQTGSCVPEH

Mouse OBP (UniProt ID OBP1A) – SEQ ID NO. 4

MAKFLLLALTFGLAHAAMEGPWKTVAIAADRVDKIERGGELRIYCRSLTCEKECKEMKVTFYVNNEN
 GQCSLTTITGYLQEDGKTYKTQFQGNTRYKLVDESPENLTFYSENVDRADRKTLLFILGHGPLTSE
 QKEKFAELAEKGIKIPAGNIREVLITDYCPE

Mouse OBP2A (UniProt ID Q8K1H9) – SEQ ID NO. 5

MKSLLLTILLGLVAVLKAQEAPPDDLVDYSGIWIYAKAMVHNGTLPSHKIPSIVFPVRIIALEEGDLE
 TTVVFWNNGHCREFKFVMKKTEEPGKYTAFHNTKVIHVEKTSVNEHYIFYCEGRHNGTSSFGMG
 KLMGRDSGENPEAMEEFKNFIKRMNLRLENMFVPEIGDKCVESD

Mouse OBP1B (UniProt ID A2AEP0) – SEQ ID NO. 6

MMVKFLLLALVFGLAHVHAHDHPELQGGQWKTAIMADNIDKIETSGPLELFFVREITCDEGCQKM
 KVTFYVKQNGQCSLTTVTGYKQEDGKTFKNQYEGENNYKLLKATSENLFYDENVDRASRKTLLY
 ILGKGEALTHEQKERLTELATQKGIKIPAGNLRELAHEDTCPE

Rat OBP (PDB ID 3FIQ) – SEQ ID NO. 7

HHENLDISPSEVNGDWRTLYIVADNVEKVAEGGSLRAYFQHMECGDECQELKIIFNVKLDSECQT
HTVVGQKHEDGRYTTDYSGRNYFHVLKKTDDIIFHNVNVDSESGKETNVILVAGKREDLNKAQKQ
ELRKLAEEYNIPNENTQHLVPTDTCNQ

Rat OBP (PDB ID 3ZQ3) – SEQ ID NO. 8

MRGSHHHHHHTDPEEASFERGNLDVDKLNWDWFSIVVASDKREKIEENGSMRVFVQHIDVLENS
LGFTFRIKENGVCTEFSLVADKTAKDGEYFVEYDGENFTTILKTDYDNYVMFHLVNVNNGETFQLM
ELYGRTKDLSSDIKEKFAKLCVAHGITRDNIIDLTKTDRCLQA

Rat OBP (UniProt ID P08937) – SEQ ID NO. 9

MVKFLLIVLALGVSCAHHENLDISPSEVNGDWRTLYIVADNVEKVAEGGSLRAYFQHMECGDECQ
ELKIIFNVKLDSECQHTVVGQKHEDGRYTTDYSGRNYFHVLKKTDDIIFHNVNVDSESGRRQCDL
VAGKREDLNKAQKQELRKLAEEYNIPNENTQHLVPTDTCNQ

Bovine OBP (PDB ID 1OBP) – SEQ ID NO. 10

AQEEEEAEQNLSSESGPWRTVYIGSTNPEKIQENGPFRTYFRELVFDDEKGTVDYFYSVKRDGKWK
NVHVKATKQDDGTYVADYEGQNVFKIVSLSRTHLVAHNINVDKHGQTTELTGLFVKNLVEDEDLE
KFWKLTEDKGIDKKNVVNFLENEDHPHPE

Boar OBP (PDB ID 1GM6) – SEQ ID NO. 11

HKEAGQDVVTSNFDASKIAGEWYSILLASDAKENIEENGSMRVFVEHIRVLDNSSLAFKFQRKVN
ECTDFYAVCDKVGDVYTVAYYGENKFRLLVNYSDYVILHLVDVNGDKTFQLMEFYGRKPDVEP
KLKDKFVEICQYGIKENIIDLTKIDRCFQLRGSGGVQESSAE

Panda OBP (PDB ID 5NGH) – SEQ ID NO. 12

HEEGNDVRRNFDVSKISGYWYSVLLASDVREKTEENSSMRVFNHIEVLSNSSLLFNMHKVDGKC
TEIALVSDKTEKDGEYSVEYDGYNVFRIVETDYTDYIIFHLVNFKEKDSFQMMELSAREPDTSEEVRK
RFVEYCQKHGIVKENIFDLTEVDRCLQARGSEKA

Chinese hamster OBP (Ensembl ID ENSGRP00015014591.1) – SEQ ID NO. 13

MVKFLLLAFALSVSCAHHKIPEISPSEVDGKWRTLYIGADNTEKVIQGGPLRAYFRHMECSDECQTL
TITFNTKEEGKCQHTVVGKDEDGQYKTGFSGNDFHVVEKADGIIIFHNVNVDSSGKKTNVILV
AGKGKLSKEQKERLENIAKEFDISKENIQHLVPTDTCNQ

Meishan pig OBP (Ensembl ID ENSSSCP00040041163.1) – SEQ ID NO. 14

MKSLLLSLVLGLVCAQEPQPEQDPFVLSGKWITSYIGSSDLEKIGENAPFQVFMRSIEFDDKESKVYL
 NFFSKENGICEEFLIGTKQEGNTYDVNYAGNNKFVVSYASETALIISNINVDEEGDKTMTGLLGKG
 TDIEDQDLEKFKEVTRENGIPEENIVNIIERDDCPAK

Horse OBP (Ensembl ID ENSECAP0000000103.2) – SEQ ID NO. 15

MQILLLSLVLGVVCAVQEPQSETDYSLFSGEWNTIYIGSSNIEKISENGPFRILLRRLDLSAGDRIIYT
 FFLKVNQCTKISSLAIKTEENTYVCHYAGKNKFEILHLSKTAIIIDIVNEDEGLVTKMVALVGMGLG
 DIQKEDIEKFKEVAKEKEIPEENIVNIIINIDDCPTSE

Guinea pig OBP (Ensembl ID ENSCPOP00000016393.2) – SEQ ID NO. 16

MQILLLALTIGLAYAHQTLDPSEINGQWHTISIAADNVEKIGEGGPLRGYFHNHLHCYDGGCKNIGLTF
 YVKLDGNCQRFVDVLGAKQEDSDVYVAQYSGTNHFVIGKKEDAIAFYNHNTDETGKETKMIVVVA
 RRDSLTEEEQKQLQEVAGEKGIKDNIRYFRERDTCQAQ

Dog OBP (Ensembl ID ENSCAFP00040020992.1) – SEQ ID NO. 17

MKILLLCLLVLACDAHLPLPNVLTQVSGPWKTLVSSNLDKIAENGPFRIYIRINVDIPRLKILFSF
 FVKVDGECVEKSVEASIGQDNLINAHYAGGNYHQILDVTPNALIGYIVNVDDKGRITKLASLVGRG
 AHVNEEDIAKFKKLSREKGIPEENIYLGDTDNCPNHE

Tibetan pig OBP (Ensembl ID ENSSSCP00015013912.1) – SEQ ID NO. 18

MKSLLLSLVLGLVCAQEPQPEQDPFELSGKWITSYIGSSDLEKIGENAPFQVFMRSIEFDDKESKVYL
 NFFSKENGICEEFLTGTKQEGNTYDVNYAGNNKFVVSYASETALIANINVDEEGDKTMTGLLGK
 GTDIEDQDLEKFKEVTRENGIPEENIVNIIERDDCPAK

Cat felis OBP (Ensembl ID ENSFCAP00000053707.1) – SEQ ID NO. 19

RSCVIHLQCLPTGCLFSALHNGLPDGRPLPDGRLPLPDGRLPLPDSRLPLPDGRLPLPDGRLPLPDG
 RLPLPDGRLPLPDGRLPLPDGRLPLPDGRLPLPDGRLPLPEGRPLPLPDSHPPLQDNLTQLSGEWNTL
 LVAATNVDKISNGPFHGYICKVDVDVTNGTVVFNFSVMMNGRCTEKSAVGTIGRDKFINIGSMN
 QNLFNLSVTSNTIAINVNTRRNTTKAFALDNTGNIFNIGYDSLGLIIHTANVDTAGQTTQV FALL
 GKRLHPDDNDFAKFRELMRENNIPEENLIDMSKTEKCPKKEKGTNPS

Chimpanzee OBP (Ensembl ID ENSPTRP00000048681.3) – SEQ ID NO. 20

MALLLSLGLSLITAQEFDPNRNVMQRNYNMARVSGVWYSIFMADDLNRIKENGDLRVFVQNIIEHL
 KNGSLKFD FEYMVQGECAVVVVCEKTEKNGEYSIN YEGQNTVAVSETDYRLFITFHLQNFRTNGTE
 THTLALYETCKKYGLGSQNIINLTNKDPCYSKHYRSPRPPMRE

OBP::GQ₂₀::KP (recombinant protein) – SEQ ID NO. 21

QEPQPEQDPFELSGKWITSYIGSSDLEKIGENAPFQVFMRSIEFDDKESKVYLNFFSKENGICEEFSLI
 GTKQEGNTYDVNYAGNNKFVVSYASETALIISNINVDEEGDKTGMTGLLGKGTDIEDQDLEKFKEVT
 RENGIPEENIVNIIERDDCPAGQG
 QGGVCGPSPPCITT

OBP::GQ₂₀::CBM (recombinant protein) – SEQ ID NO. 22

QEPQPEQDPFELSGKWITSYIGSSDLEKIGENAPFQVFMRSIEFDDKESKVYLNFFSKENGICEEFSLI
 GTKQEGNTYDVNYAGNNKFVVSYASETALIISNINVDEEGDKTGMTGLLGKGTDIEDQDLEKFKEVT
 RENGIPEENIVNIIERDDCPAGQG
 GQASPIGEGTFDDGPEGWVAYGTDGPLDTSTGALCVAVPAGSAQYGVGVVLNGVAIEEGTTYTL
 RYTATASTDVTRALVGQNGAPYGTVLDTSPALTSEPRQVTETFTASATYPATPAADDPEGQIAFQ
 LGGFSADAWTLCLDDVALDSEVEL

References

- (1) Flower, D. R. The lipocalin protein family: structure and function. *Biochem. J.* 1996, 318, 1-14.
- (2) Flower, D. R. Beyond the superfamily: the lipocalin receptors. *Biochimica et biophysica acta* 2000, 1482, 327-336.
- (3) Breer, H. Olfactory receptors: molecular basis for recognition and discrimination of odors. *Analytical and bioanalytical chemistry* 2003, 377 (3), 427-33, DOI: 10.1007/s00216-003-2113-9.
- (4) Tegoni, M.; Pelosi, P.; Vincent, F.; Spinelli, S.; Campanacci, V.; Grolli, S.; Ramoni, R.; Cambillau, C. Mammalian odorant binding proteins. *Biochimica et biophysica acta* 2000, 1482, 229-240.
- (5) Bignetti, E.; Cattaneo, P.; Cavaggioni, A.; Damiani, G. The pyrazine-binding protein and olfaction. *Comp. Biochem. Physiol.* 1988, 90B, 1-5.
- (6) Pevsner, J.; Hou, V.; Snowman, A. M.; Snyder, S. H. Odorant-binding protein: characterization of ligand binding. *The Journal of biological chemistry* 1990, 265 (11), 6118-6125.
- (7) Bignetti, E.; Cavaggioni, A.; Pelosi, P.; Persaud, K. C.; Sorbi, R. T.; Tirindelli, R. Purification and characterisation of an odorant-binding protein from cow nasal tissue. *Eur. J. Biochem.* 1985, 149, 227-231.
- (8) Dal Monte, M.; Andreini, I.; Revoltella, R.; Pelosi, P. Purification and characterization of two odorant-binding proteins from nasal tissue of rabbit and pig. *Comp Biochem Physiol* 1991, 99B (2), 445-451.
- (9) Garibotti, M.; Navarrini, A.; Pisanelli, A. M.; Pelosi, P. Three Odorant-binding Proteins from Rabbit Nasal Mucosa. *Chemical senses* 1997, 22 (4), 383-390.
- (10) Lazar, J.; Greenwood, D. R.; Rasmussen, L. E. L.; Prestwich, G. D. Molecular and Functional Characterization of an Odorant Binding Protein of the Asian Elephant,

Elephas maximus: Implications for the Role of Lipocalins in Mammalian Olfaction. *Biochemistry* 2002, 41, 11786-11794.

(11) Pes, D.; Dal Monte, M.; Ganni, M.; Pelosi, P. Isolation of two odorant-binding proteins from mouse nasal tissue. *Comp. Biochem. Physiol.* 1992, 103B (4), 1011-1017.

(12) Lobel, D.; Jacob, M.; Volkner, M.; Breer, H. Odorant of different chemical classes interact with distinct odorant binding protein subtypes. *Chemical senses* 2002, 27, 39-44.

(13) Briand, L.; Eloit, C.; Nespoulous, C.; Bezirard, V.; Huet, J.-C.; Henry, C.; Blon, F.; Trotier, D.; Permollet, J.-C. Evidence of an Odorant-Binding Protein in the Human Olfactory Mucus: Location, Structural Characterization, and Odorant-Binding Properties. *Biochemistry* 2002, 41, 7241-7252.

(14) Pelosi, P. Odorant-Binding Proteins: Structural Aspects. In *Annals New York academy of sciences; Olfaction and Taste XII: an international symposium, 1998*; pp 281-293.

(15) Spinelli, S.; Ramoni, R.; Grolli, S.; Bonicel, J.; Cambillau, C.; Tegoni, M. The Structure of the Monomeric Porcine Odorant Binding Protein Sheds Light on the Domain Swapping Mechanism. *Biochemistry* 1998, 37, 7913-7918.

(16) Perduca, M.; Mancia, F.; Del Giorgio, R.; Monaco, H. L. Crystal Structure of a Truncated Form of Porcine Odorant-Binding Protein. *Proteins: Structure, Function, and Genetics* 2001, 42, 201-209.

(17) Cave, J. W.; Wickiser, J. K.; Mitropoulos, A. N. Progress in the development of olfactory-based bioelectronic chemosensors. *Biosensors & bioelectronics* 2019, 123, 211-222, DOI: 10.1016/j.bios.2018.08.063.

(18) Pelosi, P.; Mastrogiacomo, R.; Iovinella, I.; Tuccori, E.; Persaud, K. C. Structure and biotechnological applications of odorant-binding proteins. *Applied microbiology and biotechnology* 2014, 98 (1), 61-70, DOI: 10.1007/s00253-013-5383-y.

(19) Mulla, M. Y.; Tuccori, E.; Magliulo, M.; Lattanzi, G.; Palazzo, G.; Persaud, K.; Torsi, L. Capacitance-modulated transistor detects odorant binding protein chiral interactions. *Nature communications* 2015, 6, 6010, DOI: 10.1038/ncomms7010.

(20) Paolini, S.; Tanfani, F.; Fini, C.; Bertoli, E.; Pelosi, P. Porcine odorant-binding protein: structural stability and ligand affinities measured by Fourier-transform infrared spectroscopy and fluorescence spectroscopy. *Biochimica et biophysica acta* 1999, 1431, 179-188.

(21) Sorokowska, A.; Sorokowski, P.; Szmajke, A. Does Personality Smell? Accuracy of Personality Assessments Based on Body Odour. *European Journal of Personality* 2012, 26 (5), 496-503, DOI: 10.1002/per.848.

(22) Ozeki, C.; Moro, O. A study of the suppression of body odour in elderly subjects by anti-fungal agents. *International journal of cosmetic science* 2016, 38 (3), 312-8, DOI: 10.1111/ics.12295.

(23) Di Pietrantonio, F.; Cannata, D.; Benetti, M.; Verona, E.; Varriale, A.; Staiano, M.; D'Auria, S. Detection of odorant molecules via surface acoustic wave biosensor array

based on odorant-binding proteins. *Biosensors & bioelectronics* 2013, 41, 328-34, DOI: 10.1016/j.bios.2012.08.046.

(24) Sankaran, S.; Khot, L. R.; Panigrahi, S. Biology and applications of olfactory sensing system: A review. *Sensors and Actuators B: Chemical* 2012, 171-172, 1-17, DOI: 10.1016/j.snb.2012.03.029.

(25) Goncalves, F.; Ribeiro, A.; Silva, C.; Cavaco-Paulo, A. Release of Fragrances from Cotton Functionalized with Carbohydrate-Binding Module Proteins. *ACS Appl Mater Interfaces* 2019, DOI: 10.1021/acsami.9b08191.

(26) Kozlowski, L. P. IPC - Isoelectric Point Calculator. *Biology direct* 2016, 11 (1), 55, DOI: 10.1186/s13062-016-0159-9.

(27) Cennamo, N.; Di Giovanni, S.; Varriale, A.; Staiano, M.; Di Pietrantonio, F.; Notargiacomo, A.; Zeni, L.; D'Auria, S. Easy to Use Plastic Optical Fiber-Based Biosensor for Detection of Butanal. *PloS one* 2015, 10 (3), e0116770, DOI: 10.1371/journal.pone.0116770.

(28) Capo, A.; Pennacchio, A.; Varriale, A.; D'Auria, S.; Staiano, M. The porcine odorant-binding protein as molecular probe for benzene detection. *PloS one* 2018, 13 (9), e0202630, DOI: 10.1371/journal.pone.0202630.

(29) Vincent, F.; Ramoni, R.; Spinelli, S.; Grolli, S.; Tegoni, M.; Cambillau, C. Crystal structures of bovine odorant-binding protein in complex with odorant molecules. *European journal of biochemistry* 2004, 271 (19), 3832-42, DOI: 10.1111/j.1432-1033.2004.04315.x.

(30) Goncalves, F.; Castro, T. G.; Nogueira, E.; Pires, R.; Silva, C.; Ribeiro, A.; Cavaco-Paulo, A. OBP fused with cell-penetrating peptides promotes liposomal transduction. *Colloids and surfaces. B, Biointerfaces* 2018, 161, 645-653, DOI: 10.1016/j.colsurfb.2017.11.026.

(31) Malpeli, G.; Folli, C.; Cavazzini, D.; Sartori, G.; Berti, R. Purification and Fluorescent Titration of Cellular Retinol-Binding Protein. In *Methods in Molecular Biology*; Redfern, C. P. F., Ed.; 1998; pp 111-122.

(32) Nogueira, E.; Mangialavori, I. C.; Loureiro, A.; Azoia, N. G.; Sárria, M. P.; Nogueira, P.; Freitas, J.; Harmark, J.; Shimanovich, U.; Rollet, A.; Lacroix, G.; Bernardes, G. J. L.; Guebitz, G.; Hebert, H.; Moreira, A.; Carmo, A. M.; Rossi, J. P. F. C.; Gomes, A. C.; Preto, A.; Cavaco-Paulo, A. Peptide anchor for folate-targeted liposomal delivery. *Biomacromolecules* 2015, 16 (9), 2904-2910, DOI: 10.1021/acs.biomac.5b00823.

(33) Gonçalves, F.; Silva, C.; Ribeiro, A.; Cavaco-Paulo, A. 1-Aminoanthracene Transduction into Liposomes Driven by Odorant-Binding Protein Proximity. *ACS Applied Materials & Interfaces* 2018, DOI: 10.1021/acsami.8b10158.

(34) Goncalves, F.; Castro, T. G.; Azoia, N. G.; Ribeiro, A.; Silva, C.; Cavaco-Paulo, A. Two Engineered OBPs with opposite temperature-dependent affinities towards 1-aminoanthracene. *Scientific reports* 2018, 8 (1), 14844, DOI: 10.1038/s41598-018-33085-8.

[0076] The term "comprising" whenever used in this document is intended to indicate the presence of stated features, integers, steps, components, but not to preclude the presence or addition of one or more other features, integers, steps, components or groups thereof.

[0077] The disclosure should not be seen in any way restricted to the embodiments described and a person with ordinary skill in the art will foresee many possibilities to modifications thereof.

[0078] The above described embodiments are combinable.

[0079] The following claims further set out particular embodiments of the disclosure.

C L A I M S

1. Fragrance release composition comprising:
0.01 to 5000 μM of an isolated or artificial protein with at least 90% of identity with an amino acid sequence selected from the following list: SEQ. ID NO. 1, SEQ. ID NO. 2, SEQ. ID NO. 3, SEQ. ID NO. 4, SEQ. ID NO. 5, SEQ. ID NO. 6, SEQ. ID NO. 7, SEQ. ID NO. 8, SEQ. ID NO. 9, SEQ. ID NO. 10, SEQ. ID NO. 11, SEQ. ID NO. 12, SEQ. ID NO. 13, SEQ. ID NO. 14, SEQ. ID NO. 15. SEQ. ID NO. 16, SEQ. ID NO. 17, SEQ. ID NO. 18, SEQ. ID NO. 19, SEQ. ID NO. 20, SEQ. ID NO. 21, or mixtures thereof;
0.1 μM to 2 M of an active agent selected from a list comprising a deodorizing agent, a natural essence, a fragrance, a moisturizing agent, or mixtures thereof;
wherein the active agent is bounded and/or entrapped to the protein; and
wherein the protein releases the active agent in the presence of an electrolyte solution, at a temperature between 10-70 $^{\circ}\text{C}$;
wherein the active agent has a molecular weight from 20 to 1000 g/mol;
wherein the electrolyte solution is sweat, salty water or micellar water.
2. Composition according to the previous claim wherein the protein has at least 95% of identity with an amino acid sequence selected from the following list: SEQ. ID NO. 1, SEQ. ID NO. 2, SEQ. ID NO. 3, SEQ. ID NO. 4, SEQ. ID NO. 5, SEQ. ID NO. 6, SEQ. ID NO. 7, SEQ. ID NO. 8, SEQ. ID NO. 9, SEQ. ID NO. 10, SEQ. ID NO. 11, SEQ. ID NO. 12, SEQ. ID NO. 13, SEQ. ID NO. 14, SEQ. ID NO. 15. SEQ. ID NO. 16, SEQ. ID NO. 17, SEQ. ID NO. 18, SEQ. ID NO. 19, SEQ. ID NO. 20, SEQ. ID NO. 21, or mixtures thereof.
3. Composition according to any of the previous claims wherein the protein is identical to an amino acid sequence selected from the following list: SEQ. ID NO. 1, SEQ. ID NO. 2, SEQ. ID NO. 3, SEQ. ID NO. 4, SEQ. ID NO. 5, SEQ. ID NO. 6, SEQ. ID NO. 7, SEQ. ID NO. 8, SEQ. ID NO. 9, SEQ. ID NO. 10, SEQ. ID NO. 11, SEQ. ID NO. 12, SEQ. ID NO. 13, SEQ. ID NO. 14, SEQ. ID NO. 15. SEQ. ID NO. 16, SEQ. ID NO. 17, SEQ. ID NO. 18, SEQ. ID NO. 19, SEQ. ID NO. 20, SEQ. ID NO. 21, or mixtures thereof.

4. Composition according to any of the previous claims comprising 0.2 μM to 1 M of active agent.
5. Composition according to any of the previous claims wherein the affinity constant of the protein to the active agent in water or a buffer solution ranges between 1-4.5 μM .
6. Composition according to any of the previous claims wherein the affinity constant of the protein to the active agent in the electrolyte solution ranges between 0.1-0.99 μM .
7. Composition according to any of the previous claims wherein the active agent has a molecular weight from 75-300 g/mol.
8. Composition according to any of the previous claims wherein the active agent is a fragrance molecule.
9. Composition according to any of the previous claims wherein the active agent comprises a functional group selected from aromatic, aldehyde or alcohols.
10. Composition according to any of the previous claims wherein the active agent is a fragrance molecule selected from a list comprising: 2-acetyl-1-pyrroline, α -terpineol, β -ocimene, benzyl acetate, butyl acetate, camphor, carvone, citral, citronellol, coumarin, diacetyl, eugenol, fructose, gamma decalactone, gamma nonalactone, geraniol, hedione, isoamyl acetate, lilac aldehyde, limonene, linalool, mefrosol, menthol, methyl butyrate, myrcene, pinene, pomarose, pulegone, sandalore, vanillin, or mixtures thereof.
11. Composition according to any of the previous claims wherein the release of the active agent occurs during 30 seconds to 24 h.
12. Composition according to any of the previous claims wherein the electrolyte solution has a pH between 4.0-8.5.

13. Composition according to any of the previous claims wherein sweat is human sweat or pet sweat.
14. Composition according to any of the previous claims wherein the protein is stable in polar and non-polar solvents, preferably, ethanol, methanol, butanol, benzene and undecanol.
15. Composition according to any of the previous claims wherein the protein is stable in temperatures between 18-70 °C.
16. Composition according to any of the previous claims wherein the protein is stable in a pH range of 4.0-10.0.
17. Composition according to any of the previous claims wherein the release of the active agent from the protein occurs at 20-40 °C.
18. Composition according to any of the previous claims further comprising glycerol, erythritol, arabitol, sorbitol, mannitol, xylitol, mannitol, glucosylglycerol, glucose, fructose, sucrose, trehalose, isofluoroside, dextrans, levans, polyethylene glycol, salts of chloride, citrate, sulfates, acetate, or phosphates, or mixtures thereof.
19. Use of the composition described in the previous claims as a cosmetic agent/composition.
20. Use of the composition according to the previous claim as a deodorant agent/composition.
21. Kit or article comprising the composition described in any of the previous claims.
22. Kit or article according to the previous claim wherein the article is selected from a list comprising fabric, textiles, fibers, clothes, scarfs, hats, gloves, socks and turbans, shoes, insoles, bags, handbags, detergents, creams, lotions, foams, perfumes, softeners, aerosols, deodorants, lipsticks, lip creams, face mask powders, face and

body creams, skin clarifiers, primers, foundations, eyelash mascaras, hair shampoos, hair serum, hairs masks, hair conditioners or hair coloration creams.




Fig. 1

INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2021/056011

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61K8/64 A61K8/11 A61Q15/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K A61Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>SILVA CARLA ET AL: "Odorant binding proteins: a biotechnological tool for odour control", APPLIED MICROBIOLOGY AND BIOTECHNOLOGY, SPRINGER BERLIN HEIDELBERG, BERLIN/HEIDELBERG, vol. 98, no. 8, 5 October 2013 (2013-10-05), pages 3629-3638, XP035328978, ISSN: 0175-7598, DOI: 10.1007/S00253-013-5243-9 [retrieved on 2013-10-05] the whole document</p> <p style="text-align: center;">----- -/--</p>	1-22



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

30 September 2021

Date of mailing of the international search report

06/10/2021

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Bader, Karl Günther

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2021/056011

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>PARISI M ET AL: "Unfolding and refolding of porcine odorant binding protein in guanidinium hydrochloride: equilibrium studies at neutral pH", BIOCHIMICA ET BIOPHYSICA ACTA (BBA) - PROTEINS & PROTEOMICS, ELSEVIER, NETHERLANDS, vol. 1652, no. 2, 1 December 2003 (2003-12-01), pages 115-125, XP004476493, ISSN: 1570-9639, DOI: 10.1016/J.BBAPAP.2003.08.009 the whole document -----</p>	1-22

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB2021/056011

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:
 - a. forming part of the international application as filed:
 - in the form of an Annex C/ST.25 text file.
 - on paper or in the form of an image file.
 - b. furnished together with the international application under PCT Rule 13~~ter~~.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.
 - c. furnished subsequent to the international filing date for the purposes of international search only:
 - in the form of an Annex C/ST.25 text file (Rule 13~~ter~~.1(a)).
 - on paper or in the form of an image file (Rule 13~~ter~~.1(b) and Administrative Instructions, Section 713).
2. In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments: