


Graphical Abstract

Labelling of amino acid esters followed by investigation of the conditions of cleavage of the chromophore by nucleophiles or electrolysis



Development of a temporary marker for peptides

M. Sameiro T. Gonçalves and Hernâni L.S. Maia*

Department of Chemistry, University of Minho, Gualtar, P-4710-057 Braga,
Portugal, hlsmaia@quimica.uminho.pt

3-[(*N,N*-dimethylaminophenyl)-4'-diazenyl]-benzoic acid was coupled with several amino acid esters and the product further acylated with Boc. The material thus obtained was then submitted to cleavage by electrolysis and nucleophilic attack in order to evaluate the possibility to use this chromophore as a temporary marker.

Introduction

In recent years, the use of dyes, or dye-like molecules in biomedical applications, has seen a remarkable growth in research interest and technical importance, and at present it is probably the fastest expanding area of dye chemistry. This, can be illustrated by the use of dyes in many diagnostic applications, often to allow qualitative and quantitative determinations to be performed easily and reliably by rapid and economic methods.¹ Such applications range from simple organic reactions for spectroscopic detection and measurement of body fluid analytes to high definition imaging technology for tumor detection.

Diazo coupling is particularly useful in methods for identification of proteins and for determining enzyme activity.² Although the procedures do not employ dyes as such, the end result is an azo dye chromophore. Diazo coupling has long been employed in protein chemistry, and as early as 1915 Pauly first used diazotised sulfanilic acid (Pauly reagent) for coupling with tyrosine and histidine residues to form coloured products.^{3,4} The resulting azo compounds are coloured and several spectrophotometric methods have since been developed for various applications, such as protein labelling, detection of drug abuse, diagnosing diseases, immunological assays and cancer treatment. Later, Kozaki *et al.*⁵ improved Pauly's method for quantitative analysis of L-histidine.

Meanwhile, new methods and also improvements in existing methods for the determination of enzyme activity have been developed.^{6,7} The search for photoresponsive conformational and biological properties led Behrendt *et al.*⁸ to design small cyclic peptides containing azobenzene moieties in the backbone. Sebestyén *et al.*⁹ reported the synthesis and some properties of free peptides and peptide libraries labelled with chromophores and studied the effect of colour labelling on the biological activity of a model peptide. It is now clear that dye chemistry will continue to attract the biochemist or clinician into becoming involved with such materials and, thus, they should retain an awareness of classical dye chemistry.

Having this in mind, we have acylated several amino acid derivatives with an azo dye to test its use as a marker for possible application in biological assays. The acylating reagent was a reactive azo dye we had developed for textile applications and that was used with good results in dyeing wool and polyamide fibres.¹⁰ The coloured products were submitted to different cleavage tests in order to investigate their use as a temporary marker.

Results and discussion

One equivalent of a carboxyl azo dye (**1**) obtained from 3-aminobenzoic acid and *N,N*-dimethylaniline was reacted with amino acid methyl or ethyl esters in

DMF by a DCC/ HOBt coupling (Scheme 1). After purification by chromatography (dry or flash) on silica gel followed by recrystallisation, the corresponding orange 3-[(*N,N*-dimethylaminophenyl)-4'-diazenyl]-benzoyl derivatives (**2 a-h**) were obtained as solid materials in yields ranging within 56 and 99% (Table 1); these were characterised by elemental analysis and by NMR (^1H and ^{13}C), FTIR and visible spectroscopy. The visible spectra showed a peak with λ_{max} at 415 and ϵ falling between 17000 (**2a**) and 33145 (**2g**). All products were stable on storage in the air and at room temperature. The *tert*-butyl ester of the dipeptide phenylalanyl-valine was also acylated under identical conditions to yield 78% of the expected stable derivative (**2i**) with λ_{max} 419 and ϵ 27648. With the aim to test the possibility to recover the initial amino acid esters by removal of the chromophore and considering that formyl, acetyl and benzoyl groups can be cleaved by *N,N*-diethylaminoethyl-amine (DEAEA) from the amide bond of Boc-acylamides under very mild conditions,¹¹ compounds **2 b-h** were converted into the corresponding equally coloured *tert*-butoxycarbonyl derivatives. For this purpose they were reacted at room temperature with di-*tert*-butyl pyrocarbonate in dry acetonitrile and in the presence of catalytic amount of 4-(*N,N*-dimethylamino)-pyridine (DMAP). After purification by dry chromatography, the coloured reaction products **3 b-h** were obtained in yields ranging from 56 to 99% (Table 1) and characterised as above. Their visible

spectra showed λ_{max} falling within 418 nm (**3b**) and 470 nm (**3e**), with ϵ values varying between 7723 (**3d**) and 24589 (**3g**).

Deacylation of the coloured Boc-acylamides (**3 b-h**) was then carried out by aminolysis with DEAEA in dry acetonitrile at room temperature; the expected Boc-amino acids (**4 b-h**) were isolated as colourless and usually non-crystalline materials in yields within the range 40-78% (Scheme 1, Table 2). TLC showed that a coloured by-product was also formed in all cleavages of compounds **3 b-h** with DEAEA; in a few cases (**3b**, **3f**, **3g** and **3h**) it was isolated in yields within the range 74-100% and characterised. Suspecting that this was the transamination product resulting from transfer of the dye moiety to DEAEA, a genuine sample of this compound was prepared by direct acylation of DEAEA with the dye. The product (**5**) was characterised by IR and NMR (^1H and ^{13}C) spectroscopy and by high resolution mass spectrometry and compared well with the by-product referred above. As formation of this by-product consumed DEAEA, at least two equivalents of this reagent had to be used in the cleavage reactions.

In order to investigate the electrochemical behaviour of the dye (**1**) with the aim to test deacylation by electrolysis, a cyclic voltammogram of its methyl ester (**14**) in dimethylformamide (DMF) containing tetrabutylammonium tetrafluoroborate (TBAB) as the supporting electrolyte was obtained, showing

peaks at -1.16 and -2.22 V. The compound was then electrolysed¹² at a constant potential 50 mV more negative than that of the second peak, as no reaction was observed when electrolysis was attempted at the first peak; the material was completely consumed after a few hours, which we assigned to reduction of the azo group.¹³⁻¹⁵ Compound **3h** showed to behave similarly to the dye, showing reduction peaks at -1.19 and -2.03 V and no reaction at the first peak. Thus, it was electrolysed at a constant potential 50 mV more negative than that of the second peak. The reaction was monitored by HPLC, showing that 88% of the initial material had been consumed 6 hours after it had been started; work up of the reaction mixture at this stage yielded 32% of the expected cleavage product (**4h**) together with a material that showed to correspond to 32% of *N*-(3-aminobenzoyl)-*N*-*tert*-butyloxycarbonyl-phenylalanine ethyl ester (**6h**) (Scheme 2, Table 2). The latter must have resulted from reductive cleavage of the azo group and its cyclic voltammogram showed a single peak at -2.31 V, which falls within the region of potentials that would be expected for cleavage of the 3-aminobenzoyl group. When **6h** was electrolysed at a potential 50 mV more negative than that corresponding to this peak, the expected cleavage product (**4h**) was obtained in a yield of 30%. These results suggest that reductive cleavage at the azo group and that at the amide nitrogen atom occur at similar potentials, somehow below -2 V, compound **6h**

being possibly an intermediate. Following this result, **3h** was electrolysed at – 2.5 V to give **4h** in a yield of 48% of pure product.

Colourless products were also obtained by reductive cleavage of the azo bond of compounds **3c** and **3h** by zinc powder¹⁶ in the presence of formic acid to give the corresponding *N*-(3-aminobenzoyl)-amino acid esters **6c** and **6h** in yields of 39 and 45% as pure materials (Scheme 2, Table 2). As it would be expected, for this reduction to occur there would be no need for the aid of a Boc group, which was confirmed when **2h** was reacted under the same conditions as above to give phenylalanine ethyl ester (**7h**) in a yield of 30%. The low yields in pure products obtained in these reactions with zinc were assigned to difficulties met during purification. In fact, the reactions reached completion within only ten minutes and no sign of side products was found, which supports our believe that the initial coloured materials were completely converted into the expected colourless derivatives.

In addition to labelling amino acids or peptides at their N-terminus, the alternative acylation at a lysine ω -amine group was also investigated. Thus, the methyl ester of *N*-benzyloxycarbonyl lysine was reacted with **1** under the conditions reported above, to give the expected coloured derivative (**8**) in a yield of 66%. The product was then reacted with di-*tert*-butyl pyrocarbonate; when only a slight excess of the reagent was used (2.4 equivalents), a mixture

of the required diacylation product (**9**, 45%) was obtained together with the mono acylated one (**10**, 35%). Thus, the reaction was repeated with a larger excess of pyrocarbonate (6 equivalents), which yielded 90% of **9**. This was treated with DEAEA as above, giving the product of cleavage (**11**) of both the ω -acyl group and the initial N-protecting group (Z) in a yield of 92%. The product (**12**) of saponification of **8** was obtained with a yield of 98% and then coupled with alanine methyl ester (**13c**) and phenylalanine ethyl ester (**13h**) in high yields (95 and 97%, respectively).

In conclusion, acylation with the dye can be performed at the N-terminus of either amino acid esters or peptide esters to give coloured products in high yields. As the compounds explored in our work are acceptable models for larger peptides or even proteins, our results suggest that the dye is suitable for marking materials of biological interest. Alternatively, marking can be performed at a lysine residue when such is appropriate either before or after the peptide is made. Moreover, if required, the colour can be eliminated *in situ* either by removing the chromophore with base or by electrolysis, the efficiency of the latter being not as satisfactory as that of the former. However, our best approach seems to lie on breaking the chromophore by reducing the azo group with zinc.

Experimental

All melting points are uncorrected and they were measured on a Gallenkamp melting point apparatus. TLC analyses were carried out on 0.25 mm thick precoated silica plates (Merck Fertigplatten Kieselgel 60F₂₅₄) and spots were visualised under UV or by exposure to vaporised iodine. Dry and column chromatography were carried out on Merck Kieselgel (230-240 mesh). Light petroleum refers to the fraction boiling within the range (40-60) °C. IR spectra were determined on a Perkin Elmer FTIR-1600 and UV spectra were determined on a Hitachi U-2000 spectrophotometer. ¹H NMR spectra were recorded on a Varian 300 spectrometer in 5% CDCl₃ solution at 25 °C. All chemical shifts are given in δ ppm using $\delta_{\text{H}} \text{Me}_4\text{Si}=0$ as reference and *J* values are given in Hz. Assignments were made by comparison of chemical shifts, peak multiplicities and *J* values. ¹³C NMR spectra were run in the same instrument but at 75.4 MHz using the solvent peak as internal reference. Spectrometric analysis were performed at the "Unidad de Espectrometria de Masas" of the University of Vigo, Spain. Elemental analyses were carried out on a Leco CHNS 932 instrument. For controlled potential electrolysis experiments a Hi-Tek potentiostat DT 2101, and a Hi-Tek wave generator PP RI, connected to a Philips recorder PM 8043 were used. The electrolysis cell was a conventional two-compartment, three-electrode, home-built batch cell of

the type illustrated elsewhere.¹⁷ HPLC experiments were run on a Shimadzu instrument, type 6A, connected to a Merck pre-packed column, type Hibar RT 250-4, filled with LiChrospher 100 CH-18/2 (5 μ m) and the eluent was a mixture of acetonitrile and water. The peaks were measured with a Shimadzu integrator, type C-R6A Chromatopack. Phenylalanine methyl ester and *N*-benzyloxycarbonyl-lysine were a commercial products. Other amino acid methyl esters were prepared with thionyl chloride by the usual procedure and compound **1** was synthesised according to the procedure described elsewhere.¹⁰

General method for acylation with the dye

3-[(*N,N*-Dimethylaminophenyl)-4'-diazanyl]-benzoic acid (**1**) was reacted in a 1.12-mmolar scale with an amino acid methyl or ethyl ester hydrochloride (or peptide *tert*-butyl ester) in DMF by a standard DCC/HOBt coupling. After dry chromatography on silica and recrystallisation from ethyl acetate/hexane, the required product was obtained as an orange solid.

N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazanyl]-benzoyl}-proline methyl ester **2a*

The product of reaction of **1** with proline methyl ester hydrochloride (185 mg, 1.12 mmol) was chromatographed using chloroform/methanol 6.5:0.5 as the

eluent to give the ester **2a** (237 mg, 56%), mp 119.4-121.3 °C, R_f 0.94 (chloroform/methanol 6.5:0.5) (Found: C, 66.38; H, 6.45; N, 14.70. C₂₁H₂₄N₄O₃ requires C, 66.30; H, 6.36; N, 14.73%).

N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazanyl]-benzoyl}-glycine methyl ester **2b*

The product of reaction of **1** with glycine methyl ester hydrochloride (141 mg, 1.12 mmol) was chromatographed using diethyl ether/hexane (mixtures of increasing polarity) as the eluent to give the ester **2b** (375 mg, 99%), mp 122.2-124.6 °C, R_f 0.20 (ethyl ether/hexane 9:1) (Found: C, 63.72; H, 6.15; N, 16.26. C₁₈H₂₀N₄O₃ requires C, 63.51; H, 5.92; N, 16.46%).

N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazanyl]-benzoyl}-alanine methyl ester **2c*

The product of reaction of **1** with alanine methyl ester hydrochloride (156 mg, 1.12 mmol) was chromatographed using diethyl ether/hexane 9:1 as the eluent to give the ester **2c** (360 mg, 91%), mp 132.4-134.1 °C, R_f 0.56 (diethyl ether/hexane 9:1) (Found: C, 64.65; H, 6.21; N, 15.68. C₁₉H₂₂N₄O₃ requires C, 64.39; H, 6.26; N, 15.81%).

N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazenyl]-benzoyl}-valine methyl ester **2d*

The product of reaction of **1** with valine methyl ester hydrochloride (188 mg, 1.12 mmol) was chromatographed using diethyl ether/hexane 9:1 as the eluent to give the ester **2d** (346 mg, 81%), mp 114.3-115.9 °C, R_f 0.70 (diethyl ether/hexane 9:1) (Found: C, 65.74; H, 6.88; N, 14.74. $C_{21}H_{26}N_4O_3$ requires C, 65.95; H, 6.85; N, 14.65%).

N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazenyl]-benzoyl}-isoleucine methyl ester **2e*

The product of reaction of **1** with isoleucine methyl ester hydrochloride (203 mg, 1.12 mmol) was chromatographed using diethyl ether/hexane 9:1 as the eluent to give the ester **2e** (351 mg, 79%), mp 117.3-118.1 °C, R_f 0.78 (diethyl ether/hexane 9:1) (Found: C, 66.68; H, 7.16; N, 14.21. $C_{22}H_{28}N_4O_3$ requires C, 66.64; H, 7.12; N, 14.13%).

N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazenyl]-benzoyl}-leucine methyl ester **2f*

The product of reaction of **1** with leucine methyl ester hydrochloride (162 mg, 1.12 mmol) was chromatographed using diethyl ether/light petroleum 9.5: 0.5

as the eluent to give the ester **2f** (260 mg, 59%), mp 136.4-137.5 °C, R_f 0.71 (diethyl ether/light petroleum 9.5:0.5) (Found: C, 66.69; H, 7.11; N, 14.10. C₂₂H₂₈N₄O₃ requires C, 66.64; H, 7.12; N, 14.13%).

***N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazenyl]-benzoyl}-methionine**

methyl ester 2g

The product of reaction of **1** with methionine methyl ester hydrochloride (223 mg, 1.12 mmol) was chromatographed using diethyl ether/light petroleum 9.5:0.5 as the eluent to give the ester **2g** (323 mg, 70%), mp 120.5-121.3 °C, R_f 0.68 (diethyl ether/light petroleum) (Found: C, 60.81; H, 6.32; N, 13.56; S, 7.53. C₂₁H₂₆N₄O₃S requires C, 60.84; H, 6.32; N, 13.52; S, 7.74%).

***N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazenyl]-benzoyl}-phenylalanine**

ethyl ester 2h

The product of reaction of **1** with phenylalanine ethyl ester hydrochloride (257 mg, 1.12 mmol) was chromatographed using chloroform/methanol 6:1 as the eluent to give the ester **2h** (385 mg, 77%), mp 152.7-153.8 °C, R_f 0.70 (diethyl ether/ light petroleum 9.5:0.5) (Found: C, 70.28; H, 6.37; N, 12.68. C₂₆H₂₈N₄O₃ requires C, 70.25; H, 6.35; N, 12.61%).

***N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazenyl]-benzoyl}-phenylalanyl-
valine *tert*-butyl ester **2i****

The product of reaction of **1** with phenylalanyl-valine *tert*-butyl ester (161 mg, 0.60 mmol) was chromatographed using chloroform/methanol (mixtures of increasing polarity) as the eluent to give ester **2i** (267 mg, 78%), mp 164.8-166.9 °C, R_f 0.71 (ethyl acetate/hexane 6:4) (Found C, 69.02; H, 7.00; N, 12.30. C₃₃H₄₁N₅O₄ requires C, 69.33; H, 7.23; N 12.25%).

General method for preparation of Boc-acylamides

To a solution of the required substrate in dry acetonitrile (47 mmol dm⁻³) 0.3 eq. of DMAP was added followed by 3.6 eq. of di-*tert*-butyl pyrocarbonate under rapid stirring over night at room temperature, the reaction being monitored by TLC. Evaporation under reduced pressure followed by dry chromatography on silica gel and recrystallisation gave the required Boc-acylamide as an orange residue.

***N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazenyl]-benzoyl}-*N-tert*-
butyloxycarbonyl-glycine methyl ester **3b****

The product of reaction of **2b** (247 mg, 0.73 mmol) was chromatographed with diethyl ether/hexane 6:4 as the eluent; the solid material thus obtained was

recrystallised from ethyl acetate/hexane to give ester **3b** (288 mg, 90%), mp 114.8-116.0 °C, R_f 0.40 (diethyl ether/hexane 6:4) (Found: C, 62.61; H, 6.38; N, 12.55. C₂₃H₂₈N₄O₅ requires C, 62.71; H, 6.41; N, 12.72%).

N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazenyl]-benzoyl}-*N*-*tert*-butyloxycarbonyl-alanine methyl ester **3c*

The product of reaction of **2c** (50 mg, 0.14 mmol) was chromatographed with diethyl ether/light petroleum 9:1 as the eluent to give ester **3c** (63 mg, 99%), mp 101.8-103.4 °C, R_f 0.74 (diethyl ether/hexane 8:2) (Found: C, 63.52; H, 6.55; N, 12.21. C₂₄H₃₀N₄O₅ requires C, 63.42; H, 6.65; N, 12.33%).

N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazenyl]-benzoyl}-*N*-*tert*-butyloxycarbonyl-valine methyl ester **3d*

The product of reaction of **2d** (189 mg, 0.50 mmol) was chromatographed using diethyl ether/hexane 2:8 as the eluent to give ester **3d** (203 mg, 85%), mp 147.6-149.0 °C, R_f 0.62 (diethyl ether/hexane 6:4) (Found: C, 64.94; H, 6.82 N, 11.63. C₂₆H₃₄N₆O₅ requires C, 64.71; H, 7.10; N, 11.61%).

***N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazenyl]-benzoyl}-*N*-*tert*-
butyloxycarbonyl-isoleucine methyl ester **3e****

The product of reaction of **2e** (220 mg, 0.56 mmol) was chromatographed with diethyl ether/hexane 4:6 as the eluent to give ester **3e** (154 mg, 56%), mp 89.2-91.0 °C, R_f 0.56 (diethyl ether/hexane 4:6) (Found: C, 65.20; H, 7.39; N, 11.04. C₂₇H₃₆N₄O₅ requires C, 65.30; H, 7.31; N, 11.28%).

***N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazenyl]-benzoyl}-*N*-*tert*-
butyloxycarbonyl-leucine methyl ester **3f****

The product of reaction of **2f** (130 mg, 0.33 mmol) was chromatographed with ethyl acetate/hexane 6:4 as the eluent to give ester **3f** (161 mg, 99%), R_f 0.59 (diethyl ether/hexane 6:4); found: M⁺, 496.268479. C₂₇H₃₆N₄O₅ requires *m/z*, 496.268571.

***N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazenyl]-benzoyl}-*N*-*tert*-
butyloxycarbonyl-methionine methyl ester **3g****

The product of reaction of **2g** (210 mg, 0.51 mmol) was chromatographed with diethyl ether/hexane 3:7 as the eluent to give ester **3g** (155 mg, 60%), R_f 0.47 (diethyl ether/hexane 6:4); *m/z* (EI) 514.225675 (M⁺. C₂₆H₃₄N₄O₅S requires 514.224992).

N*-{3-[(*N,N*-Dimethylaminophenyl)-4'-diazenyl]-benzoyl}-*N-tert*-butyloxycarbonyl-phenylalanine ethyl ester **3h*

The product of reaction of **2h** (62.6 mg, 0.14 mmol) was chromatographed with diethyl ether/hexane 1:1 as the eluent to give ester **3h** (76 mg, 99%), R_f 0.40 (diethyl ether/hexane 1:1) (Found: C, 68.48; H, 6.75; N, 9.88. $C_{31}H_{36}N_4O_5$ requires C, 68.36; H, 6.66; N, 10.29%).

General method of aminolysis of the coloured amino acid esters

The coloured substrates **3 b-h** were treated with a given amount of DEAEA for one or more days according to the procedure of Grehn¹¹ *et al.* The products were purified by flash chromatography to give the corresponding Boc-amino acid esters **4 b-h** together with the transamination product **5**.

N-tert*-Butyloxycarbonyl-glycine methyl ester **4b** by aminolysis of **3b*

The product of a 1-day reaction of **3b** (100 mg, 0.23 mmol) with DEAEA (0.16 cm³, 0.11 mmol) was chromatographed with diethyl ether/hexane 4:6 as the eluent to give ester **4b** (24 mg, 56%), R_f 0.54 (diethyl ether/hexane 6:1) (Found: C, 50.57; H, 7.82, N, 7.09 requires $C_8H_{15}NO_4$: C, 50.78; H, 7.99; N, 7.40%).

N*-tert-Butyloxycarbonyl-alanine methyl ester **4c** by aminolysis of **3c*

The product of a 2-day reaction of **3c** (75 mg, 0.17 mmol) with DEAEA, ($47 \times 10^{-3} \text{ cm}^3$, 0.33 mmol) was chromatographed with diethyl ether/hexane 8:26 as the eluent to give ester **4c** (21 mg, 63%), R_f 0.67 (diethyl ether/hexane 6:4) (Found: C, 53.47; H, 8.28; N, 6.59. $\text{C}_9\text{H}_{17}\text{NO}_4$ requires C, 53.18; H, 8.43; N, 6.89%).

N*-tert-Butyloxycarbonyl-valine methyl ester **4d** by aminolysis **3d*

The product of a 2-day reaction of **3d** (79.5 mg, 0.17 mmol) with DEAEA, ($93 \times 10^{-3} \text{ cm}^3$, 0.66 mmol) was chromatographed with diethyl ether/hexane 4:6 to give ester **4d** (15 mg, 40%), R_f 0.79 (diethyl ether/hexane; 6:4) (Found: C, 57.36; H, 8.92; N, 5.65. $\text{C}_{11}\text{H}_{21}\text{NO}_4$ requires C, 57.12; H, 9.15; N, 6.06%).

N*-tert-Butyloxycarbonyl-isoleucine methyl ester **4e** by aminolysis of **3e*

The product of a 1-day reaction of **3e** (81.8 mg, 0.17 mmol) with DEAEA ($93 \times 10^{-3} \text{ cm}^3$, 0.66 mmol). was chromatographed with diethyl ether/hexane 2:8 to give ester **4e** (29 mg, 72%), R_f 0.78 (diethyl ether/hexane 6:4) (Found: C, 58.89; H, 9.43; N, 5.32. $\text{C}_{12}\text{H}_{23}\text{NO}_4$ requires C, 58.75; H, 9.45; N, 5.71%).

N*-tert-Butyloxycarbonyl-leucine methyl ester **4f** by aminolysis of **3f*

The product of a 1-day reaction of **3f** (79 mg, 0.16 mmol) with DEAEA ($93 \times 10^{-3} \text{ cm}^3$, 0.66 mmol) was chromatographed with diethyl ether/light petroleum 2:8 to give ester **4f** (19 mg, 49%), R_f 0.76 (diethyl ether/hexane 6:4) (Found: C, 58.96; H, 9.55; N, 5.43. $\text{C}_{12}\text{H}_{23}\text{NO}_4$ requires C, 58.75; H, 9.45; N, 5.71%).

N*-tert-Butyloxycarbonyl-methionine methyl ester **4g** by aminolysis of **3g*

The product of a 1-day reaction of **3g** (83 mg, 0.16 mmol) with DEAEA ($93 \times 10^{-3} \text{ cm}^3$, 0.66 mmol) was chromatographed with diethyl ether/light petroleum 2:8 to give ester **4g** (31 mg, 78%), R_f 0.66 (diethyl ether/ light petroleum 6:4) (Found: C, 50.13; H, 7.80; N, 5.28; S 12.59. $\text{C}_{11}\text{H}_{21}\text{NO}_4\text{S}$ requires C, 50.17; H, 8.04; N, 5.32; S, 12.18%).

N*-tert-Butyloxycarbonyl-phenylalanine ethyl ester **4h** by aminolysis of **3h*

The product of a 1-day reaction of **3h** (60 mg, 0.11 mmol) with DEAEA ($31 \times 10^{-3} \text{ cm}^3$, 0.22 mmol) was chromatographed with diethyl ether/hexane 2:8 to give ester **4h** (23 mg, 71%), R_f 0.90 (chloroform/methanol 5.8:0.2) (Found: C, 65.38; H, 7.77; N, 4.86. $\text{C}_{16}\text{H}_{23}\text{NO}_4$ requires C, 65.51; H, 7.90; N, 4.78; %).

General method of controlled potential electrolysis of the coloured amino acid esters

Both compartments of a two-compartment cell for controlled-potential electrolysis were filled with acetonitrile containing Et_4NHCl (0.1 mol dm^{-3}) as supporting electrolyte and Et_3NHCl (0.12 mol dm^{-3}) as a proton donor.¹² At this stage the substrate (0.31 mmol) was added to the cathodic compartment and a cyclic voltammogram recorded at a sweep rate of 100 mV s^{-1} . Then, the potential was adjusted to a value 50 mV more negative than that corresponding to the peak chosen for electrolysis and the apparatus switched on. When the intensity of the current was almost zero, the reaction mixture (catholyte) was transferred to a round-bottomed flask and the solvent evaporated under reduced pressure. The residue was dissolved in water and extracted with ethyl acetate, dried over MgSO_4 and after concentration of the organic layer under reduced pressure the residue was chromatographed on silica gel (diethyl ether/hexane, mixtures of increasing polarity).

N*-tert-Butyloxycarbonyl-phenylalanine ethyl ester **4h** by electrolysis of **3h*

Electrolysis of **3h** (54 mg 0.10 mmol) at a potential of -2.50 V gave ester **4h** (14 mg, 48%). When **3h** (167 mg, 0.31 mmol) was electrolysed at a potential of -2.03 V , **4h** was obtained (18 mg, 32%) together with the corresponding

aminobenzoyl derivative **6h** (25 mg, 32%); m/z (EI) 412.198227 (M^+ . $C_{23}H_{28}N_2O_5$ requires 412.199822).

N*-tert-Butyloxycarbonyl-phenylalanine ethyl ester **4h** by electrolysis of **6h*

Electrolysis of **6h** (90 mg, 0.22 mmol) at a potential of -2.31 V gave ester **4h** (19 mg, 30%).

2-(*N,N*-Diethylamino)-*N'*-{3-[(*N,N*-dimethylaminophenyl)-4'-diazenyl]-benzoyl}-ethylamine **5**

To a dispersion of **1** (52.9 mg, 0.22 mmol) in dry acetonitrile (1.1 cm^3) DEAEA ($62 \times 10^{-3}\text{ cm}^3$, 0.44 mmol) was added and the mixture left at room temperature for 6.5 hours under stirring, the reaction being monitored by TLC (chloroform/methanol 6:4). This mixture was filtered and the solid residue recrystallised from ethyl acetate/hexane to give compound **5** as an orange solid (53 mg, 66%), mp $86.3\text{-}88.3\text{ }^\circ\text{C}$, R_f 0.66 (chloroform/methanol 6:4); m/z (EI) 367.238227 (M^+ . $C_{21}H_{29}N_5O$ requires 367.237211).

General method of reductive cleavage of the coloured amino acid esters with zinc dust

Reductive cleavage by zinc dust in methanol in the presence of formic acid was carried out according to the procedure described by Gowda¹⁶ *et al.* The required product was isolated by flash chromatography (silica: ethyl acetate/hexane, mixtures of increasing polarity) and then characterised.

N*-tert-Butyloxycarbonyl-*N*-(3-aminobenzoyl)alanine methyl ester **6c** by chemical reduction of **3c*

Reduction of **3c** with zinc (357 mg, 0.79 mmol) gave the corresponding aminobenzoyl derivative **6c** (99 mg, 39%) as an oil; *m/z* 322.152869. (M⁺. C₁₆H₂₂N₂O₅ requires 322.152872).

N*-tert-Butyloxycarbonyl-*N*-(3-aminobenzoyl)phenylalanine ethyl ester **6h** by chemical reduction of **3h*

Reduction of **3h** (280 mg, 0.52 mmol) with zinc gave the corresponding aminobenzoyl derivative **6h** (96 mg, 45%), which compared well with a sample obtained by electrolysis.

N*-(3-Aminobenzoyl)-phenylalanine ethyl ester **7h** by chemical reduction of **2h*

Reduction of **2h** (198 mg, 0.45 mmol) with zinc gave the amino acid ester **7h** (41 mg, 30%), mp 121.0-123.4 °C, R_f 0.55 (ethyl acetate/hexane 8:2); *m/z* (EI) 312.147630 (M⁺. C₁₈H₂₀N₂O₅ requires 321.147393).

N*-Benzyloxycarbonyl- ω {3-[(*N,N*-dimethylaminophenyl)-4'-diazenyl]-benzoyl}-lysine methyl ester **8*

The product of reaction of **1** with *N*-Benzyloxycarbonyl-lysine methyl ester hydrochloride (329 mg, 1.12 mmol) carried out according to the general method described above for acylation with the dye was chromatographed using chloroform/methanol 5.8:0.2 as the eluent to give ester **8** (401 mg, 66%), mp 114.0-115.9 °C, R_f 0.75 (chloroform/methanol 5:1) (Found: C, 65.76; H, 6.56; N, 12.49. C₃₀H₃₅N₅O₅ requires C, 66.04; H, 6.47; N, 12.84%).

N*-Benzyloxycarbonyl-*N*, ω -bis(*tert*-butyloxycarbonyl)- ω {3-[(*N,N*-dimethylaminophenyl)-4'-diazenyl]-benzoyl}-lysine methyl ester **9*

The product of a 2-day reaction of **8** (100 mg, 0.18 mmol) with di-*tert*-butyl pyrocarbonate (240 mg, 1.10 mmol) carried out according to the general method described above for preparation of Boc-acylamides was

chromatographed with ethyl acetate/hexane 2:8 as the eluent to give ester **9** (127 mg, 90%), R_f 0.88 (diethyl ether/hexane 6:4); m/z (EI) 745.369343 (M^+ . $C_{40}H_{51}N_5O_9$ requires 745.368679).

N*-Benzyloxycarbonyl-*N*-*tert*-butyloxycarbonyl- ω -{3-[(*N,N*-dimethylaminophenyl)-4'-diazenyl]-benzoyl}-lysine methyl ester **10*

The product of a 3-day reaction of **9** (234 mg, 0.43 mmol) with di-*tert*-butyl pyrocarbonate (224 mg, 1.03 mmol) was chromatographed with ethyl acetate/hexane 2:8 to give ester **10** (100 mg, 35%), R_f 0.77 (diethyl ether/hexane 6:4); m/z (EI) 645.315487 (M^+ . $C_{35}H_{43}N_5O_7$ requires 645.316249) together with **9** (148 mg, 45%).

N*, ω -Bis(*tert*-butyloxycarbonyl-lysine methyl ester **11** by aminolysis of **9*

The fully acylated ester **9** (100 mg, 0.13 mmol) was reacted for 3 days with DEAEA (0.148 cm³, 1.04 mmol) according to the general method described above for aminolysis of the coloured amino acid esters and the product chromatographed with diethyl ether/hexane (mixtures of increasing polarity) to give ester **11** (43 mg, 92%) (Found: C, 56.88; H, 8.93; N, 7.43. $C_{17}H_{32}N_2O_6$ requires C, 56.64; H, 8.95; N, 7.76%).

N*-Benzyloxycarbonyl- ω -{3-[(*N,N*-dimethylaminophenyl)-4'-diazenyl]-benzoyl}-lysine **12*

To the fully protected amino acid **8** (151 mg, 0.28 mmol) in 1,4- dioxane (1.39 cm³) 1M NaOH (0.28 cm³, 0.28 mmol) was added. The solution was stirred at room temperature for 4h and acidified to pH 2-3 with 1M KHSO₄. The red precipitate thus formed was filtered off to give the amino acid derivative **12** (144 mg, 98.0%); mp 112.5-114.5 °C.

General method for coupling the coloured amino acid

One equivalent of compound **12** was reacted with an amino acid methyl or ethyl ester hydrochloride in DMF by a standard DCC/HOBt coupling. After dry chromatography on silica and recrystallisation from ethyl acetate/hexane, the required product was obtained as an orange solid.

N*-Benzyloxycarbonyl- ω -{3-[(*N,N*-dimethylaminophenyl)-4'-diazenyl]-benzoyl}-lysyl-alanine methyl ester **13c*

The product of reaction of **12** (144 mg, 0.27 mmol) with alanine methyl ester (38 mg, 0.27 mmol) was chromatographed with chloroform/methanol 5.8: 0.2 as the eluent to give ester **13c** (159 mg, 95%), mp 179.9-181.7 °C, R_f 0.5

(chloroform/methanol 5.5: 0.5) (Found: C, 63.99; H, 6.58; N, 13.58. $C_{33}H_{40}N_6O_6$ requires C, 64.27; H, 6.54; N, 13.63%).

N*-Benzyloxycarbonyl- ω {3-[(*N,N*-dimethylaminophenyl)-4'-diazenyl]-benzoyl}-lysyl-phenylalanine ethyl ester **13h*

The product of reaction of **12** (163 mg, 0.31 mmol) with phenylalanine methyl ester (71 mg, 0.31 mmol) was chromatographed with chloroform/methanol 5.8: 0.2 was used as the eluent to give ester **13h** (266 mg, 97%), mp 146.9-148.9 °C, R_f 0.68 (ethyl acetate/hexane 8:2) (Found: C, 67.94; H, 6.45; N, 11.87. $C_{40}H_{46}N_6O_6$ requires C, 68.00; H, 6.56; N, 11.89%).

Methyl 3-[(*N,N*-dimethylaminophenyl)-4'-diazenyl]-benzoate **14**

Compound **1** (269 mg, 1.0 mmol) was reacted with thionyl chloride (73×10^{-3} cm³, 2.0 mmol) by the usual procedure. Recrystallisation of the product from acetone/hexane gave the required ester **14** (283 mg, 100%) as a dark red solid, mp 82.8-84.7 °C; R_f 0.77 (chloroform).


References

- 1 J.C.V.P. Moura, in *Colorants for Non-Textile Applications*, eds. H.S. Freeman and A.T. Peters, Elsevier Science B.V., New York, 2000, p. 189
- 2 T. Imai, *Med. Technol.* (Tokyo), 1985, **13**, 1403
- 3 J. Kapuscinski., *J.Histochem. Cytochem.*, 1990, **38**, 1323
- 4 W. Beisker and W.G. Eisert, *J.Histochem. Cytochem.*, 1989, **37**, 1699
- 5 Y.Kozaki, S.Kozai and H. Matsumoto, *Tokushima Bunri Daigaku Kenkyu Kiyō*, 1995, **50**, 135
- 6 R. Schmidt, Ger. Offen. DE 4 117 619, 1992
- 7 J. Backhaus, D. Mangold and W.R. Knappe, Gr. Offen. DE 3 813 503, 1989
- 8 R. Behrendt, M.Schenk, H. Musiol and L. Moroder, *J. Peptide Sci.*, 1999, **5**, 519
- 9 F. Sebestyén, G. Szendrei, M. Mák, M. Dóda, E. Illyés, G. Szókán, K. Kindla, W. Rapp, P. Szegó, E. Câmpian and À. Furka, *J. Peptide Sci*, 1998, **4**, 294
- 10 J.I.N.R. Gomes, J. Griffiths, H.L.S. Maia, J.C.V.P. Moura and A.M.F. Oliveira-Campos, *Dyes and Pigments*, 1991, **17**, 269


- 11 L. Grehn, K. Gunnarsson and U. Ragnarsson, *Acta Chem. Scand., Ser. B*, 1987, **B41**, 18
- 12 L. Grehn, H.L.S. Maia, L.S. Monteiro, M.I. Montenegro and U. Ragnarsson, *J. Chem. Res.*, 1991, (S), 144; (M), 1501
- 13 L. Horner at H. Lung, in *Organic Electrochemistry*, eds. M.M. Baizer and H. Lung, Marcel Dekker, New York, 1991
- 14 T.M. Florence, D.A. Johnson and G.E. Batley, *Electroanalytical Chemistry and Interfacial Electrochemistry*, 1974, **50**, 113
- 15 T.M. Florence, *Aust. J. Chem.*, 1965, **18**, 609
- 16 Gowda, K. Abiraj and D. C. Gowda, *Tetrahedron Lett.*, 2002, **43**, 1329
- 17 L. Grehn, K. Gunnarsson, H.L.S. Maia, M.I. Montenegro, L. Pedro and U. Ragnarsson, *J. Chem. Res.*, 1988, (S) 399; (M), 3081

Acknowledgements

We thank the Foundation for Science and Technology (Portugal) for financial support to the Institute of Biotechnology and Fine Chemistry (University of Minho).




Scheme 1



c $\text{R} = \text{CH}_3, \text{R}^1 = \text{Me}$
h $\text{R} = \text{CH}_2\text{Ph}, \text{R}^1 = \text{Et}$

Scheme 2



Scheme 3

Table 1 - Synthesis of coloured compounds **2** and **3**

Product	Compound no.	Yield/%	
		2 (R = H)	3 (R = Boc)
Dpa-Pro-OMe (a)		56	–
Dpa-Gly(<i>N</i> -R)-OMe (b)		99	90
Dpa-Ala(<i>N</i> -R)-OMe (c)		91	99
Dpa-Val(<i>N</i> -R)-OMe (d)		81	85
Dpa-Ile(<i>N</i> -R)-OMe (e)		79	56
Dpa-Leu(<i>N</i> -R)-OMe (f)		59	99
Dpa-Met(<i>N</i> -R)-OMe (g)		70	60
Dpa-Phe(<i>N</i> -R)-OEt (h)		77	99
Dpa-Phe(<i>N</i> -R)-Val- <i>O</i> <i>t</i> Bu (i)		78	–

Table 2 - Selective cleavage of the chromophores

Starting material	Deprotection method	Product	Yield/%
3b	DEAEA	4b	56
3c	DEAEA	4c	63
3d	DEAEA	4d	40
3e	DEAEA	4e	72
3f	DEAEA	4f	49
3g	DEAEA	4g	78
3h	DEAEA	4h	71
9	DEAEA	11	84
3h	Electrolysis at –2.03 V	4h + 6h	32 + 32
3h	Electrolysis at –2.5 V	4h	48
6h	Electrolysis at –2.31 V	4h	30
2h	Zn / HCO ₂ H	7h	30
3c	Zn / HCO ₂ H	6c	39
3h	Zn / HCO ₂ H	6h	45

Table 3 - Results obtained in the synthesis of compounds **8-12**

Product (compound no.)	Yield/%
Z-Lys(ω -Dpa)-OMe (8)	66
Z-Lys(<i>N</i> -Boc, ω -Dpa, ω -Boc)-OMe (9)	90
Z-Lys(ω -Dpa)-OH (12)	98
Z-Lys(ω -Dpa)-Ala-OMe (13c)	95
Z-Lys(ω -Dpa)-Phe-OEt (13h)	97