

## New perspectives of *Juglans regia* L. phytochemicals against *Candida* species

Natália Martins,<sup>1,2</sup> Lillian Barros,<sup>1</sup> Ana Maria Carvalho<sup>1</sup>, Sónia Silva,<sup>2</sup> Mariana Henriques,<sup>2</sup> Isabel C.F.R. Ferreira<sup>1</sup>

<sup>1</sup>Mountain Research Centre (CIMO), ESA, Polytechnic Institute of Bragança, Portugal.

<sup>2</sup>IBB - Institute for Biotechnology and Bioengineering, Centre of Biological Engineering, University of Minho, Braga, Portugal.

Opportunistic fungal infections have deserved special relevance in the last decades, presenting itself, a serious problem in terms of public health. Despite *Candida albicans* was considered the main agent responsible for those infections, other non-*albicans* *Candida* species have also been described in the last years [1-3]. Most of the species are susceptible to antimicrobial drugs, but recently it has been observed a growing number of microorganisms with drug resistance. Therefore, the discover/use of alternative therapies is crucial [4].

*Juglans regia* L. (walnut) leaves are commonly used in traditional medicine as antiseptic, antimicrobial and anti-inflammatory [5]; those benefits could be related with its richness in phenolic compounds [6]. In the present work, the antifungal potential of the hydroalcoholic extract prepared from walnut leaves was evaluated against a total of nineteen *Candida* strains (from the species: *C. albicans*, *C. glabrata*, *C. parapsilosis* and *C. tropicalis*), using the disc diffusion halo assay.

All the tested strains were sensible to the plant extract. The obtained values of the inhibitory zones ranged between 0.9-1.4 cm, being the halo maintained after 48h. The observed antifungal activity is certainly related to the phenolic compounds previously determined in the extract [6]: five phenolic acid derivatives-caffeoylquinic and p-coumaroylquinic acid derivatives, two dimers and one trimer of procyanidins, twelve flavonols- quercetin, myricetin and kaempferol derivatives, and five taxifolin O-pentoside isomers; 3-O-caffeoylquinic acids and quercetin O-pentoside were the main phenolic compounds. Further studies are necessary in order to elucidate the most active compounds and the specific role of each one.

Acknowledgments: The authors are grateful to FCT (Portugal) for financial support to CIMO (PEst-OE/AGUI0690/2011), N. Martins (SFRH/BD/87658/2012), L. Barros (\*Compromisso para a Ciência 2008) and S. Silva (NORTE-07-0124-FEDER-000027; QREN) contracts.

### References:

- [1] Kim J, Sudbery P. (2011) *J Microbiol.* 49, 171-177.
- [2] Tsai P-W, Chen Y-T, Hsu P-C, Lan C-Y. (2013) *BioMedicine.* 3, 51-64.
- [3] Vázquez-González D, Perusquia-Ortiz AM, Hündeiker M, Bonifaz A. (2013) *J Ger Soc Dermatol.* 11, 381-94.
- [4] Kanafani ZA, Perfect JR. (2008) *Clin Infect Dis - Antimicrob Resist.* 46, 120-128.
- [5] Murray MT. (2004) 2<sup>nd</sup> ed. New York, NY. Random House.
- [6] Santos A, Barros L, Calheta RC, Dueñas M, Carvalho AM, Santos-Buelga C, Ferreira ICFR. (2013) *Ind Crops Prod.* 51, 430-436.