

# Characterization of New 5-Aminoimidazole Derivatives

Soraia Fernandes, Eva Pereira, M. Fernanda Proença and Alice M. Dias


Centro de Química, Universidade do Minho, 4710-057 Braga, Portugal

[ad@quimica.uminho.pt](mailto:ad@quimica.uminho.pt)

The imidazole nucleus is present in a significant number of biomolecules and the inclusion of this moiety in organic scaffolds is considered an important synthetic strategy in drug discovery.<sup>1</sup>

5-Aminoimidazoles are attractive building blocks in chemical synthesis because they are key components in many bioactive compounds. Few reports on the chemistry of these compounds appear in the literature due to their known instability.<sup>2</sup>

In our research group, imidazoles **1** are easily accessible from an efficient method previously developed. Studies on the reactivity of the 4-cyanoformimidoyl group with primary amines showed that competitive nucleophilic displacement of NH<sub>3</sub> or HCN occur leading to two alternative products.<sup>3</sup> The experimental conditions are determinant to achieve the selective synthesis of each product. As an extension of this work, condensation of imidazoles **1** and hydrazines have been studied. Here, we report the characterization of the products obtained from condensation of imidazoles **1** with 1-aminopiperidine.



Attempts to react imidazoles **1** with 1-aminopiperidine led us to isolate two different compounds. From the <sup>1</sup>H NMR data we confirmed that both products have in common the 5-aminoimidazole nucleus and the piperidine moiety. The presence of an extra amino group in the <sup>1</sup>H NMR spectrum of one of the products was a very good evidence for the formation of compound **3**. The presence of a cyano band in the IR spectrum of the second product supported the formation of product **2**. Finally, <sup>13</sup>C and 2D NMR spectra were determinant to the assignment of structure **2** and **3**. The <sup>1</sup>H, <sup>13</sup>C and 2D NMR and IR data will be presented and discussed.

## Acknowledgements:

The authors gratefully acknowledge the University of Minho and the Foundation for the Science and Technology (FCT, Portugal) for financial support to the NMR portuguese network (PTNMR, Bruker Avance III 400-Univ. Minho). FCT and FEDER (European Fund for Regional Development)-COMPETE-QREN-EU for financial support to the Research Centre, CQ/UM [PEst-C/QUI/UI0686/2013 (FCOMP-01-0124-FEDER-037302)].

## References:

- [1] Helmick, R. A.; Fletcher, A. E.; Gardner, A. M.; Gessner, C.R.; Hvitved, A. N.; Gustin, M.C.; Gardner, P.R. *Antimicrob. Agents Chemother.* (2005), **49**,1837.
- [2] Soh, C. H. *et al.*, *J. Comb. Chem.* (2006), **8**, 464-468 and references there in.
- [3] Dias, A. M.; Vila-Chã, A. M.; Costa, A. L.; Cunha, D; Senhorães, N; Proença, M. F.; *Synlett*, (2011), 2675.