

Short Communication

Extended-spectrum β -lactamase and carbapenemase-producing *Aeromonas* species in wild animals from Portugal

C. Dias, C. R. Serra, L. C. Simões, M. Simões, A. Martinez-Murcia, M. J. Saavedra

AEROMONAS are Gram-negative, facultative-anaerobic, non-spore-forming, glucose-fermenting, oxidase- and catalase-positive rods (Martin-Carnahan and Joseph 2005). Apart from fish, which are widely reported hosts for *Aeromonas*, insects, crustaceans, reptiles, birds and mammals were also found to harbour *Aeromonas* species, both in healthy and disease state (Pearson and others 2000, Turutoglu and others 2005, Evangelista-Barreto and others 2006, Ceylan and others 2009). An increase in resistance levels of the genus, particularly to β -lactam antimicrobial agents, has been observed not only in clinical isolates, but also in environmental strains (Saavedra and others 2004, 2007).

The most common mechanism of antibacterial resistance is the production of three chromosomally encoded β -lactamases, which have been described and identified in different *Aeromonas* (Janda and Abbott 2010). They may or may not concomitantly occur in the same strain, and their coordinated expression is induced by the presence of β -lactam antibiotics (Walsh and others 1997, Avison and others 2004). These enzymes comprise cephalosporinases (Ambler's class C), penicillinases/oxacillinases (class D) such as OXA-type enzymes and metallo- β -lactamases (class B) (Hayes and others 1994). The

most common metallo- β -lactamases produced by this genus are of the 'CphA'-type, whose sequences appear to be widely distributed in *Aeromonas hydrophila* and *Aeromonas veronii* strains (Walsh and others 1997). Recently, two other metallo- β -lactamases, VIM and IMP, have




FIG 1: Unrooted phylogenetic tree based on *gyrB* gene sequences of strains isolated in this study, and the representative strains of the genus, *Aeromonas*, of the MDC collection. The isolated included in the study are present in bold-face

Veterinary Record (2014)

doi: 10.1136/vr.101803

C. Dias, MSc,
C. R. Serra, PhD,
M. J. Saavedra, PhD,
Veterinary and Animal Science
Research Centre, Carla Dias,
Centre for the Research and Technology
for Agro-Environment and Biological
Sciences, University of Trás-os-Montes
e Alto Douro, Vila Real 5000-801,
Portugal

L. C. Simões, PhD,
IBB – Institute for Biotechnology and
Bioengineering, Centre of Biological
Engineering, University of Minho,
Braga 4710-057, Portugal

L. C. Simões, PhD,
M. Simões, PhD,
LEPAE – Department of Chemical
Engineering, Faculty of Engineering,
University of Porto, Porto 4200-465,
Portugal

A. Martinez-Murcia, PhD,
Area de Microbiología, EPSO,
Universidad Miguel Hernández,
Orihuela E-03300, Alicante, Spain

M. J. Saavedra, PhD,
Department of Veterinary Sciences,
School of Agriculture and Veterinary
Science, University of Trás-os-Montes
e Alto Douro, Vila Real 5000-801,
Portugal;

E-mail for correspondence:
saavedra@utad.pt

CD and CRS contributed equally

Provenance: Not commissioned;
externally peer reviewed

Accepted March 20, 2014

Short Communication

TABLE 1: Characterisation of strains isolated as concerns resistance phenotypes and *bla* genotypes

Animal species name	Strains	Identification (<i>gyrB</i> gene sequencing)	Resistance phenotype	β -lactamase genes content				
				<i>bla</i> _{OXA-aer}	<i>bla</i> _{CTX-M}	<i>bla</i> _{FOX}	<i>bla</i> _{MOX}	<i>bla</i> _{CphA}
<i>Cervus elaphus</i>	MDC 180	<i>Aeromonas salmonicida</i>	AML, AMC, TIC, TIM, KF, S, E, TE	-	-	-	-	-
<i>C elaphus</i>	MDC 182	<i>A salmonicida</i>	KF, S, E	-	-	-	-	-
<i>C elaphus</i>	MDC 190	<i>A salmonicida</i>	AML, AMC, TIC, TIM, KF, S, E, TE	-	-	-	-	+
<i>C elaphus</i>	MDC 193	<i>A salmonicida</i>	AML, AMC, TIC, TIM, KF, S, E, TE	-	-	+	-	+
<i>Strix aluco</i>	MDC 2447	<i>A salmonicida</i>	AML, AMC, TIM, PRL, TZP, KF, E, TE	-	-	-	+	+
<i>C elaphus</i>	MDC 2492	<i>A salmonicida</i>	AML, KF, E	+	-	-	-	+
<i>C elaphus</i>	MDC 2493	<i>A salmonicida</i>	AML, TIC, KF, S	+	-	+	-	-
<i>C elaphus</i>	MDC 2494	<i>A salmonicida</i>	AML, TIC, KF, S	+	-	+	-	-
<i>C elaphus</i>	MDC 2495	<i>A salmonicida</i>	AML, KF	+	-	+	-	-
<i>Sciurus vulgaris</i>	MDC 2485	<i>Aeromonas bestiarum</i>	AML, E	+	-	-	-	+
<i>S vulgaris</i>	MDC 2486	<i>A bestiarum</i>	AML, E	+	-	-	-	+
Colubridae	MDC 2487	<i>Aeromonas eucrenophila</i>	AML, KF, E	+	-	+	-	-
Colubridae	MDC 2488	<i>A eucrenophila</i>	AML, KF, E	+	-	+	-	-
Colubridae	MDC 2489	<i>A eucrenophila</i>	AML, KF, E	+	-	+	-	-
Colubridae	MDC 2490	<i>A eucrenophila</i>	AML, KF, E	+	-	+	-	-
<i>Circaetus gallicus</i>	MDC 2491	<i>Aeromonas veronii</i>	AML, TIC, TIM, NA, S, E, TE	+	+	+	+	+

been identified in *A hydrophila* encoded on an integron and a plasmid, respectively (Neuwirth and others 2007, Libisch and others 2008).

This study reports the identification of *Aeromonas* species from wild animals, the antibiotic resistance found in the strains isolated and the association of resistance with the presence of *bla*_{CphA}, *bla*_{IMP}, *bla*_{VIM}, *bla*_{OXA}, *bla*_{TEM}, *bla*_{CTX-M}, *bla*_{SHV}, *bla*_{MOX} and *bla*_{FOX} genes in the strains of *Aeromonas* species.

A total of 140 fecal samples from different wild animal species (birds, reptiles and mammals), were collected aseptically, directly from the animal rectum or from freshly voided faecal material. Collection was done as soon as the animal entered the Centre for Treatment of Wild Animals (University of Trás-os-Montes and Alto Douro). Samples were enriched in 5 ml of Brain Heart Infusion Broth (Oxoid), for 24 hours at 30°C. Direct streaking of faecal swabs on glutamate starch phenol-red agar medium (Merck) was used to select, isolate and purify bacterial isolates, by incubating plates aerobically at 30°C for 24 hours. All colonies that were morphologically suspected as *Aeromonas* species (yellow, smooth and round) were then selected to establish pure cultures.

Total genomic DNA was extracted from overnight pure colonies as previously described (Soler and others 2004). The presence of genes encoding TEM, SHV, CTX-M, MOX, FOX, CphA, VIM, OXA-Aer, OXA-B, OXA-C and IMP β -lactamases was analysed by PCR. The oligonucleotides were designed by Fontes (2009) within her PhD thesis (in publication).

Sequencing of 16S rRNA and *gyrB* genes was performed as previously described (Martínez-Murcia and others 1999, Soler and others 2004).

Susceptibility of *Aeromonas* isolates was determined by the disk method (Bauer and others 1966) according to the Clinical and Laboratory Standards Institute (2006). The following antibiotic-containing discs were obtained from Oxoid: amoxicillin/clavulanic acid, ticarcillin, ticarcillin/clavulanic acid, piperacillin, piperacillin/tazobactam, cephalothin, cefoxitin, ceftriaxone, cefoperazone, ceftazidime, cefotaxime, cefepime, imipenem, aztreonam, streptomycin, kanamycin, amikacin, gentamicin, tobramycin, nalidixic acid, ciprofloxacin, erythromycin, tetracycline, trimethoprim-sulfamethoxazole, chloramphenicol, fosfomycin.

A total of 325 different Gram-negative isolates were obtained from the 140 faecal samples, with 8 samples positive for the presence of *Aeromonas* species (4 from *Cervus elaphus* (red deer), 1 of a *Strix aluco* (tawny owl), 1 of a *Sciurus vulgaris* (red squirrel), 1 of a Colubridae (snake) and 1 of a *Circaetus gallicus* (short-toed snake eagle)).

Ten different *gyrB* gene partial sequences were obtained that grouped *Aeromonas* into four different species: *Aeromonas salmonicida* (n=9), *Aeromonas eucrenophila* (n=4), *Aeromonas bestiarum* (n=2) and *A veronii*

(n=1). The unrooted phylogenetic tree was constructed by using the *gyrB* gene partial sequence of each isolate and previously published reference sequences (Martínez-Murcia and others 1992, Yáñez and others 2003, Saavedra and others 2006) from the Molecular Diagnostics Center (Biomolecular Technologies S.L.U., Spain) culture collection, including type strains (Fig 1).

Resistance to at least two antibiotic was recorded for all isolates, and almost half (43.8 per cent) were found to be multiresistant (Table 1). Our results showing 93.7 per cent of isolates resistant to amoxicillin, 81.25 per cent to cephalothin and 37.5 per cent to ticarcillin are in agreement with this report. The combination of an aminopenicillin and a carboxipenicillin with a β -lactamase inhibitor (clavulanic acid), was effective in reducing resistance, as shown by the decrease in the proportion of resistant strains. This reduction was more pronounced with amoxicillin (93.7 per cent v 25 per cent) than with ticarcillin (37.5 per cent v 31 per cent). No resistance was found to the following antimicrobials: aztreonam, imipenem and cephalosporins.

The β -lactamases genes were found in different combinations, and in one case, five β -lactamase genes were detected in the same isolate, *A veronii* MDC2491. The most prevalent genes were *bla*_{OXA-aer} present in 11 isolates (69 per cent) followed by *bla*_{FOX} in nine isolates (56 per cent) and the metallo- β -lactamase encoding gene *bla*_{CphA} in seven isolates (44 per cent). The gene *bla*_{MOX} was detected in two isolates (*A salmonicida*, *A veronii*) and *bla*_{CTX-M} in one *A salmonicida* isolate. No PCR specific for *bla*_{TEM}, *bla*_{SHV}, *bla*_{OXA-B}, *bla*_{OXA-C}, *bla*_{IMP} and *bla*_{VIM} encoding sequences were detected, and yielded no evidence for the presence of these genes in any isolate.

The housekeeping gene *gyrB* has been demonstrated to be an excellent molecular chronometer for phylogenetic inference in the genus *Aeromonas* than the commonly used 16S rRNA gene sequencing (Janda and Abbott 2007), as shown by our results.

Although the majority of isolates were susceptible to the fluoroquinolones, confirming the usefulness of such antibiotics in the treatment of *Aeromonas* infections, one isolate from *C gallicus*, MDC2491, identified as *A veronii* was resistant to nalidixic acid. Isolates were susceptible to all other antimicrobials.

The complete genome sequencing of *A hydrophila* ATCC7699 and *A salmonicida* A449 revealed that both carry an array of β -lactamases genes to counteract antibacterial factors present in the environment, including several antibiotics used for human and animal clinical treatment (Seshadri and others 2006, Reith and others 2008). Inducible chromosomal β -lactamases is the resistance mechanism against β -lactam antibiotics for most *Aeromonas*, with examples described and identified in different species (Henriques and others 2006). Expression of metallo- β -lactamases active against carbapenems is also

a concern with regards to *Aeromonas* infections (Parker and Shaw 2011). This later, the 'CphA-type', is considered the most common metallo- β -lactamase produced by *Aeromonas* species (Janda and Abbott 2010), corroborating our results.

Acknowledgments

C Dias was the recipient of grant SFRH/BD/90120/2012 from the Fundação para a Ciência e Tecnologia (FCT) under OREN – POPH- Potencial Humano, Tipologia 4.1 – Formação avançada da União Europeia, and strategic research project PEst-OE/AGR/UI0772/2014.

Competing interests None.

References

- AVISON, M. B., NIUMSUP, P., NURMAHOMED, K., WALSH, T. R. & BENNETT, P. M. (2004) Role of the 'cre/blr-tag' DNA sequence in regulation of gene expression by the *Aeromonas hydrophila* β -lactamase regulator, *BlrA*. *The Journal of Antimicrobial Chemotherapy* **53**, 197–202
- BAUER, A. W., KIRBY, W. M., SHERRIS, J. C. & TURCK, M. (1966) Antibiotic susceptibility testing by a standardized single disk method. *American Journal of Clinical Pathology* **45**, 493–496
- CEYLAN, E., BERKTAS, M. & AĞAOĞLU, Z. (2009) The occurrence and antibiotic resistance of motile *Aeromonas* in livestock. *Tropical Animal Health and Production* **41**, 199–204
- CLSI (2006) Methods for antimicrobial dilution and disk susceptibility testing of infrequently isolated or fastidious bacteria. *Approved guideline M45-A*
- EVANGELISTA-BARRETO, S. N., VIEIRA, R. H., CARVALHO, F., TORRES, R., SANT'ANNA, E. S., RODRIGUES, D. P. & REIS, C. M. (2006) *Aeromonas* spp. isolated from oysters (*Crassostrea rhizophorea*) from a natural oyster bed, Ceará, Brazil. *Revista do Instituto de Medicina Tropical de Sao Paulo* **48**, 129–133
- FONTES, M. C. (2009) Diversidade filogenética e resistência a antibióticos em *Aeromonas* spp. Isoladas de suínos abatidos para consumo e de alheiras. PhD Thesis in Veterinary Sciences. University of Trás-os-Montes e Alto Douro, Vila Real, Portugal
- HAYES, M. V., THOMSON, C. J. & AMYES, S. G. (1994) Three beta-lactamases isolated from *Aeromonas salmonicida*, including a carbapenemase not detectable by conventional methods. *European Journal of Clinical Microbiology & Infectious Diseases: Official Publication of the European Society of Clinical Microbiology* **13**, 805–811
- HENRIQUES, I. S., FONSECA, F., ALVES, A., SAAVEDRA, M. J., & CORREIA, A. (2006) Occurrence and diversity of integrons and β -lactamase genes among ampicillin-resistant isolates from estuarine waters. *Research in Microbiology* **157**, 938–947
- JANDA, J. M. & ABBOTT, S. L. (2007) 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory: pluses, perils, and pit-falls. *Journal of Clinical Microbiology* **45**, 2761–2764
- JANDA, J. M. & ABBOTT, S. L. (2010) The genus *Aeromonas*: taxonomy, pathogenicity, and infection. *Clinical Microbiology Reviews* **23**, 35–73
- LIBISCH, B., GISKE, C. G., KOVÁCS, B., TÓTH, T. G. & FÜZI, M. (2008) Identification of the first VIM metallo- β -lactamase-producing multidrug-resistant *Aeromonas hydrophila* strain. *Journal of Clinical Microbiology* **46**, 1878–1880
- MARTIN-CARNAHAN, A. & JOSEPH, S. W. (2005) *Aeromonadaceae*. In *Bergey's manual of systematic bacteriology*. Eds. G. M. GARRITY, D. J. BRENNER, N. R. KRIEG, & J. T. STALEY. New York: Springer. pp. 556–578
- MARTÍNEZ-MURCIA, A. J., ANTÓN, A. I. & RODRIGUEZ-VALERA, F. (1999) Patterns of sequence variation in two regions of the 16S rRNA multigene family of *Escherichia coli*. *International Journal of Systematic and Evolutionary Microbiology* **49**, 601–610
- MARTÍNEZ-MURCIA, A. J., BENLLOCH, S. & COLLINS, M. D. (1992) Phylogenetic interrelationships of members of the genera *Aeromonas* and *Plesiomonas* as determined by 16S ribosomal DNA sequencing: lack of congruence with results of DNA–DNA hybridizations. *International Journal of Systematic and Evolutionary Microbiology* **42**, 412–421
- NEUWIRTH, C., SIEBOR, E., ROBIN, F. & BONNET, R. (2007) First occurrence of an IMP metallo- β -lactamase in *Aeromonas caviae*; IMP-19 in an isolate from France. *Antimicrobial Agents and Chemotherapy* **51**, 4486–4488
- PARKER, J. L. & SHAW, J. G. (2011) *Aeromonas* spp. clinical microbiology and disease. *Journal of Infection* **62**, 109–118
- PEARSON, M. D., HIRONO, I., AOKI, T., MIRANDA, R. & INGLIS, V. (2000) Virulence properties of motile aeromonads isolated from farmed frogs *Rana tigrina* and *R. rugulosa*. *Diseases of Aquatic Organisms* **40**, 185–193
- REITH, M. E., SINGH, R. K., CURTIS, B., BOYD, J. M., BOUEVITCH, A., KIMBALL, J., MUNHOLLAND, J., MURPHY, C., SARTY, D., WILLIAMS, J., NASH, J. H., JOHNSON, S. C. & BROWN, L. L. (2008) The genome of *Aeromonas salmonicida* subsp. *salmonicida* A449: insights into the evolution of a fish pathogen. *BMC Genomics* **9**, 427
- SAAVEDRA, M. J., FIGUERAS, M. J. & MARTÍNEZ-MURCIA, A. J. (2006) Update phylogeny of the genus *Aeromonas*. *International Journal of Systematic and Evolutionary Microbiology* **56**, 2481–2487
- SAAVEDRA, M. J., GUEDES-NOVAIS, S., ALVES, A., REMA, P., TACÃO, M., CORREIA, A. & MARTÍNEZ-MURCIA, A. (2004) Resistance to β -lactam antibiotics in *Aeromonas hydrophila* isolated from rainbow trout (*Oncorhynchus mykiss*). *International Microbiology* **7**, 207–211.
- SAAVEDRA, M. J., PEREA, V., FONTES, M. C., MARTINS, C. & MARTÍNEZ-MURCIA, A. (2007) Phylogenetic identification of *Aeromonas* strains isolated from carcasses of pig as new members of the species *Aeromonas allosaccharophila*. *Antonie Van Leeuwenhoek* **91**, 159–167
- SESHADRI, R., JOSEPH, S. W., CHOPRA, A. K., SHA, J., SHAW, J., GRAF, J., HAFT, D., WU, M., REN, Q., ROSEVITZ, M. J., MADUPU, R., TALLON, L., KIM, M., JIN, S., VUONG, H., STINE, O. C., ALI, A., HORNEMAN, A. A. J. & HEIDELBERG, J. F. (2006) Genome sequence of *Aeromonas hydrophila* ATCC 7966T: jack of all trades. *Journal of Bacteriology* **188**, 8272–8282
- SOLER, L., YAÑEZ, A., CHACÓN, M. R., AGUILERA-ARREOLA, M. G., CATALÁN, V., FIGUERAS, M. J. & MARTÍNEZ-MURCIA, A. J. (2004) Phylogenetic analysis of the genus *Aeromonas* based on two housekeeping genes. *International Journal of Systematic and Evolutionary Microbiology* **54**, 1511–1519
- TURUTOGLU, H., ERCELIK, S. & CORLU, M. (2005) *Aeromonas hydrophila*-associated skin lesions and septicemia in a Nile crocodile (*Crocodylus niloticus*). *Journal of the South African Veterinary Association* **76**, 40–42
- WALSH, T. R., STUNT, R. A., NABI, J. A., MACGOWAN, A. P. & BENNETT, P. M. (1997) Distribution and expression of β -lactamase genes among *Aeromonas* spp. *The Journal of Antimicrobial Chemotherapy* **40**, 171–178
- YAÑEZ, M. A., CATALÁN, V., APRÁIZ, D., FIGUERAS, M. J. & MARTÍNEZ-MURCIA, A. J. (2003) Phylogenetic analysis of members of the genus *Aeromonas* based on *GyrB* gene sequences. *International Journal of Systematic and Evolutionary Microbiology* **53**, 875–883



CrossMark