Activity of phenolic compounds from plant origin against *Candida* species

Natália Martins^{a,b}, Lillian Barros^a, Mariana Henriques^b, Sónia Silva^b,
Isabel C.F.R. Ferreira^{a,*}

^a Mountain Research Centre (CIMO), ESA, Polytechnic Institute of Bragança, Campus de Santa Apolónia, Apartado 1172, 5301-855 Bragança, Portugal

^b CEB, Centre of Biological Engineering, LIBRO – Laboratório de Investigação em Biofilmes Rosário Oliveira, University of Minho, 4710-057 Braga, Portugal

ARTICLE INFO

Article history:

Received 29 December 2014

Received in revised form 19 April 2015

Accepted 27 May 2015

Keywords:

Phytochemicals

Plant extracts

Phenolic compounds

Antifungal activity

Candida species

ABSTRACT

Candida albicans and other *Candida* species have been highly associated with several opportunistic fungal infections. Their ability to develop host infections is incited by different determinants, being virulence factors the most highlighted. Molecular targets of the antifungal drugs are crucial components for determination of yeast survival. Ergosterol, nucleic acids and glucan are the most studied molecular targets to destroy *Candida* species, being considered the basis of the development of new antifungal drugs. However, increasing levels of resistant *Candida* species to the current antifungal drugs have been observed, making ineffective those agents. Thus, other therapies more effective and safer than the current ones, are being studied, namely the use plant of extracts enriched in phenolic compounds. In this sense, this manuscript provides an historical perspective of the opportunistic fungal infections, molecular targets of the current anti-*Candida* drugs, as well as a general description of the active principles present in plants, focused on the antifungal potential of whole plant extracts and isolated phenolic compounds, against *Candida* species.

© 2015 Elsevier B.V. All rights reserved.

Contents

1. Introduction	649
2. Opportunistic fungal infections	649
2.1. Historical perspective	649
2.2. Anti- <i>Candida</i> targets	650
2.3. Current anti- <i>Candida</i> drugs	650
3. Active principles of plants	651
3.1. Plants as a rich source of bioactive compounds	651
3.2. Primary metabolites	651
3.3. Secondary metabolites	652
4. Activity of phenolic extracts/compounds against <i>Candida</i> species	653
4.1. The use of phenolic extracts against <i>Candida</i> species	653
4.2. The use of phenolic compounds against <i>Candida</i> species	662
5. Conclusions	666
Acknowledgements	666
References	666

* Corresponding author. Tel.: +351 273303219; fax: +351 273325405.

E-mail address: iferreira@ipb.pt (I.C.F.R. Ferreira).

1. Introduction

Fungi and other microbial species are widespread in the environment, including the soil, plants, trees, and even in the skin and other parts of the body. Most of them are not dangerous, even conferring some benefits to the host, being considered as commensal flora. Notwithstanding, some species in face to an abnormal overgrowth, might become harmful to the host and, in last instance, could compromise its life (Blanco and Garcia, 2008; Brunke and Hube, 2013).

In the last two decades, it has been observed a considerable increase in the incidence of deep fungal infections, not only in hospital environments, due to the increasing magnitude of organ transplantations, the rise of AIDS, the use of invasive devices (catheters, artificial joints and valves) and in immunocompromised patients, but also in the rest of population (Abi-Said et al., 1997; Eggimann et al., 2003a; Fanello et al., 2001; Kim and Sudbery, 2011; Li et al., 2006; Mayer et al., 2013; Raman et al., 2013; Silva et al., 2011a; Tsai et al., 2013; Wächter et al., 2012). In parallel with this, and despite the advances on medical and chemical industries increased the life expectancy, due to the indiscriminate use of some chemical substances, it has been observed an increasing resistance of pathogenic microorganisms to conventional drugs, leading to development of other complications (Agarwal et al., 2010; Alves-Silva et al., 2013; Asgarpanah and Kazemivash, 2012; Asl and Hosseinzadeh, 2008; Bakkali et al., 2008; Kanafani and Perfect, 2008; Rana et al., 2011; Sher, 2009; Shojaii and Fard, 2012; Silva et al., 2011b; Singh et al., 2010).

Currently, fungal infections are the fourth leading causes of hematogenous infections and the most common involved fungi are *Candida* species, commensal microorganisms present in any healthy people (Pierce, 2005; Tsai et al., 2013). Symptoms and signals of fungal infections are not always visible, and an evolution to a systemic infection thrives very slowly, mostly of times, without any mistrust by the individual infected. There are several causal factors that contribute to those opportunistic fungal infections, some of them difficult, or even impossible, to avert. Pollution, smoking, certain drugs, poor diet, sedentary lifestyle and stress-inducing agents are some examples of aggressors (Devasagayam et al., 2004). However, not only external factors increase the organisms vulnerability to disorders, but also genetic changes in the cells, degeneration, reduction of the cells functionality (characteristic of aging process); damage in white blood cells and attacks in other cells of the organic defense system, weak the immune system, leading to a decrease in the ability to respond to invading organisms (Valko et al., 2007).

Natural defense systems are present in living organisms, but it is necessary to consider that in some conditions/situations, an external strengthening and complementation diets are essential (Carocho and Ferreira, 2013a; Devasagayam et al., 2004). It is an established fact that any unbalance affects the well-being of body and potentiates the growing and colonization of certain invaders, among other organic disorders. Thus, a balanced diet, containing all macro and micronutrients in right proportions is crucial to an optimal health, which was already stated by Hippocrates, the father of modern medicine, nearly 2.500 years ago: “Let food be the medicine and medicine be the food”. There are some nutrients present in foods that can help to protect and strength the body, at greater or lesser degree (Devasagayam et al., 2004; Murray and Pizzorno, 1998; Singh et al., 2004; Valko et al., 2007).

Since the pre-historic era, plants, and other living organisms are used by primitive societies due to healing properties, being sought through botanical preparations. In fact, natural matrices, in particular, plants are extremely rich sources of natural biomolecules. Despite, with the passage of time, their use, as prevention or treatment of various conditions, became secondary and recently, plants have deserved a great relevance for scientific researchers that have

been studying their bioactive properties; many of those properties have been related to their richness in phenolic compounds (Carocho and Ferreira, 2013a; Murray and Pizzorno, 1998; Wojdylo et al., 2007). Therefore, in the present manuscript, the activity of phenolic extracts and compounds against *Candida* species was revised and highlighted as an alternative to current antifungal drugs, which failed in treatment of several opportunistic fungal infections related with *Candida* species.

2. Opportunistic fungal infections

2.1. Historical perspective

Microorganisms are ubiquitous in the world, with a wide and fascinating variety and diversity. In the absence of microorganisms, the life on earth is not possible, with a close dependency on them. Otherwise, they are responsible for the onset of several diseases and disorders in the host (Amara and Shibl, 2013; Isolauri et al., 2002; Kaur et al., 2009).

In particular, yeasts, which belong to a category of fungi are part of the commensal flora of the healthy population. Colonization by some yeast species is beneficial to the host, because it not only limits the growth of other opportunistic pathogenic fungi, but also stimulates the functioning of the immune system. However, in face to an abnormal overgrowth, they are able to cause a wide variety of dysfunctions/disorders to the host. *Candida* species are a good example; although they are deemed a commensal microorganism, living smoothly in the inner warm creases and crevices of the gastrointestinal (GI) tract (and vaginal tract), they are able to cause problems, mainly vaginal infections (Asmundsdóttir et al., 2009; Brunke and Hube, 2013; Eggimann et al., 2003a; McCullough et al., 1996; Tsai et al., 2013).

During some days after birth, primarily, the mucosa of GI tract and upper respiratory passages are colonized by those species, as well as mouth, pharynx and larynx. Apart of the observed biochemical variations in the last trimester of pregnancy, namely progesterone, estradiol and glycogen levels, associated with an increase in vaginal pH, which favors the emergence of these infections (Carrara et al., 2010; Vázquez-González et al., 2013), other clinical status increase the vulnerability to yeast infections (diabetes, patient's submitted to broad-spectrum of antibiotic therapy, the use of oral contraceptives) (Epstein and Polsky, 1998; Geiger et al., 1995; Liu et al., 2009; Lott et al., 2005; Sobel, 2007; Tarry et al., 2005). Furthermore, and not least important, the increasing magnitude of organ transplantations, the rise of AIDS and the use of invasive devices (catheters, artificial joints and valves) are also major factors that contribute to higher patient's susceptibility to infections in hospitals (Epstein and Polsky, 1998; Kim and Sudbery, 2011; Pierce, 2005; Vázquez-González et al., 2013).

Seven *Candida* species are classified as having major medical importance, namely *Candida albicans*, *Candida tropicalis*, *Candida glabrata*, *Candida parapsilosis*, *Candida stellatoidea*, *Candida kru-sei* and *Candida kyfer*, being *Candida albicans* the most important (Greenberg and Glick, 2003; McCullough et al., 1996; Silva et al., 2011a; Sullivan et al., 2004; Westwater et al., 2007).

Candidiasis, the most common opportunistic yeast infection in the world has been related in majority with *Candida albicans*. This microorganism is a causative agent of mouth and mucocutaneous infections, among others more complicated, such as septicemia, endocarditis, meningitis, and peritonitis, especially in patients with reduced immune function or taking antibiotic therapies (Brunke and Hube, 2013; Greenberg and Glick, 2003; McCullough et al., 1996). Nevertheless, in the last two decades, in face to the rapid increase in the incidence of opportunistic fungal infections, numerous studies had revealed that also other species of

fungi are involved in some hospital infections, like *Aspergillus* spp., *Zygomycetes* spp., *Fusarium* spp., *Scedosporium* spp., *Cryptococcus* spp., *Trichosporon* spp., *Geotrichum* spp., and *Rhodotorula* spp. (Kwamin et al., 2013; Martins et al., 2014; Tsai et al., 2013; Vázquez-González et al., 2013). But, *C. albicans* still counting nearly 50–90% of isolates from fungal infections, being able to cause serious hematogenous infections that, directly, affect the welfare of individuals (Vázquez-González et al., 2013).

2.2. Anti-Candida targets

Due to the high incidence of opportunistic fungal infections, namely triggered by *Candida* species, a wide variety of experimental studies have been carried out towards the discovery of effective treatments against these opportunistic and pathogenic microorganisms. Cellular morphology, physiology, and metabolism are crucial aspects in fungal cells survival.

All living organisms communicate with the surrounding environment and, in the case of *Candida* cells, the communication is established through the cellular membrane. Quorum-sensing is considered the mechanism of microbial communication, and for *C. albicans*, the main quorum sensing molecules are farnesol, tyrosol and dodecanol. This mechanism is not only incited by yeast growth but also by the hyphal formation (Lu et al., 2006; Mayer et al., 2013; Wyk et al., 2009). One of the main functions of cellular membrane is to maintain cells homeostasis to ensure the protection of internal organelles, being the first barrier in *Candida* cells. In the membrane, there are a wide variety of components present, playing different functions. The phospholipid bilayer is a common membrane to all eukaryotic cells, as also proteins, carbohydrates, other lipids (i.e. sterols, fatty acids, glycerol and esters) and even association of those substances. In the case of fungal cells, such as *Candida* species, ergosterol is the specific and most abundant sterol. Those molecules play important roles in the maintenance of integrity and fluidity of the membrane, acting as secondary messenger in the signal development in order to warrant the proper cell function (i.e. growth, division and reproduction). Thus, ergosterol is considered the main target of antifungal drugs (Ghannoum and Rice, 1999; Kanafani and Perfect, 2008; White et al., 1998).

Chemical substances could act by three main different ways in *Candida* cells: in the ergosterol biosynthesis (i.e. azoles) (Ghannoum and Rice, 1999; Kanafani and Perfect, 2008; Lamb et al., 1999; Lupetti et al., 2002; Sanglard, 2002); interacting with ergosterol and intercalating between membrane (i.e. polyenes) (Ghannoum and Rice, 1999; Kanafani and Perfect, 2008; Kontoyiannis and Lewis, 2002); or even causing ergosterol depletion (i.e. allylamines) (Kanafani and Perfect, 2008; Kontoyiannis and Lewis, 2002; White et al., 1998).

Unfortunately, in the last years, it has been observed an increasing drug resistance to those classes of antifungal agents, being ineffective even in higher doses. Some virulence mechanisms have been studied, not only to explain the increasing rates of antifungal drugs resistance, but also to discover/create new and promising antifungal strategies. So far, the most frequent virulence mechanism associated with *Candida* species infections is the ability of adaptation to a wide variety of habitats, commonly known as morphological transition. Notwithstanding, other factors, as the ability for biofilm formation, the secretion of hydrolytic enzymes (such as proteases and phospholipases), and the expression of adhesins and invasins linked with the dimorphic phases and phenotypic switching, have been associated with *Candida* species infections (Mayer et al., 2013; Tsai et al., 2013).

In this sense, different molecular targets needs to be studied as also more effective and selective compounds against the fungal cell wall; compounds to inhibit nucleic acids, to affect the plasma membrane, ergosterol, phospholipid, sphingolipid and pro-

tein biosynthesis; intermediates of the metabolism of nucleic acids and amino acids; compounds that affect efflux pumps, proton ATPases and other enzymes. The study of the fungal virulence factors and involved genes on antifungal resistance, among other important biological components with a direct interference on drug resistance, is crucial (Franz et al., 1998; Kanafani and Perfect, 2008; Lupetti et al., 2003; Mayer et al., 2013; Noël, 2012; Perlin, 2014; Sangamwar et al., 2008; Sanglard and Odds, 2002; White et al., 2002, 1998). Mayer et al. (2013) quoted polymorphism, secretion of hydrolases; pH sensing and regulation; metabolic adaptation; environmental stress response; heat shock proteins and small heat shock proteins, and metal acquisition as the most important virulence factors and fitness attributes, while Lewis et al. (2012) proposed that the inhibition of macrophage cell division or arrest of the host macrophages to confine the pathogens is another virulence attribute of *Candida* species, in particular *C. albicans*.

2.3. Current anti-Candida drugs

Currently, there are five main classes of antifungal agents widely used at a medical level against the major fungal pathogens, including *Candida* species: polyenes, azoles, allylamines/thiocarbamates, fluoropyrimidines and echinocandins (Maurya et al., 2013).

Azoles, polyenes and allylamine/thiocarbamates act, direct or indirectly, in the ergosterol present in all *Candida* species (Kontoyiannis and Lewis, 2002; Lupetti et al., 2003).

Polyenes comprise a class of antifungal agents that affect cellular membranes containing ergosterol. These agents present amphipathic properties which allow their intercalation between membranes containing ergosterol, forming channels that favour the ionic unbalance and destroy the proton gradient (Ghannoum and Rice, 1999). Nystatin and amphotericin-B are the most common used polyenes (Eggimann et al., 2003b). Nevertheless, not only primary but also secondary resistance of *Candida* species, namely *C. albicans*, *C. lusitanae*, *C. guilliermondii* and *C. glabrata*, have been observed (Ghannoum and Rice, 1999; Kontoyiannis and Lewis, 2002; White et al., 1998).

Azoles are another class of antifungal agents in which the target is over again ergosterol, but in this case affects its biosynthesis. Firstly introduced during the 1980s and used by oral administration, to treat oropharyngeal candidiasis in patients with VIH⁺, they act as inhibitors of sterol 14 α -demethylation. The remotion of the C32-methyl group at a 14 α - from precursor sterols, catalyzed by a microsomal cytochrome P450 monooxygenase, affect the main sterol present on fungal cells membranes: Erg11p, a lanosterol 14-alpha-demethylase. Miconazole and econazole were the original used azole compounds, followed by ketoconazole, fluconazole and itraconazole (Eggimann et al., 2003b; Ghannoum and Rice, 1999; Sardi et al., 2013). However, and despite during several years those drugs presented a great efficacy as antifungal agents, resulting in a widespread use, currently it is being observed an increasing emergence of *Candida* species-resistance (namely *C. albicans*, *C. glabrata*, *C. parapsilosis*, *C. krusei* and *C. lusitanae*) (Develoux and Bretagne, 2005; Noël, 2012; Vázquez-González et al., 2013).

Allylamines (i.e. naftifine and terbinafine) and thiocarbamates (i.e. tolnaftate and tolciclate) are the third class of antifungal agents that interact with ergosterol. But, in this case, they act as inhibitors of squalene epoxidase, blocking the conversion of squalene into 2,3-oxidosqualene, that result in ergosterol depletion and accumulation of toxic sterols. It was shown that the increasing of the squalene levels is toxic, leading to cell membrane permeability and then disrupting the cellular organization (Ghannoum and Rice, 1999; White et al., 1998). However, this class of antifungal drugs presents a weak activity against the most frequent *Candida* species, being only active against dermatophytes (Kontoyiannis and Lewis, 2002; Sanglard, 2002).

On the other hand, fluoropyrimidines and echinocandins act differently from the previous drugs; the first act as inhibitor of nucleic acids, while the second as inhibitor of the glucan synthesis and, therefore, are effective against *Candida* biofilm cells (Kontoyiannis and Lewis, 2002; Pereira Gonzales and Maisch, 2012). Fluoropyrimidines, namely 5-fluorocytosine (5FC) is a fluorinated pyrimidine that impairs nucleic acid biosynthesis due to the formation of toxic fluorinated pyrimidine antimetabolites. A permease enzyme aids 5FC to enter on fungal cells, where it is converted into 5-fluorouracil (5FU), catalyzed by the cytosine deaminase. After this, 5FU is converted into 5-fluorouridylic acid (FUMP) and, then, phosphorylated and incorporated into RNA, leading to a disruption of protein synthesis. Another mechanism of action of 5FU, results of its conversion into 5-fluorodeoxyuridine monophosphate, a potent inhibitor of the enzyme involved in DNA synthesis and nuclear division (Waldorf and Polak, 1983). There are some studies that refer an increasing resistance to 5FC (Ghannoum and Rice, 1999; Gubbins and Anaissie, 2009; Liu et al., 2009; Souza et al., 2010), and despite several mechanisms of action responsible for 5FC resistance are still not established, they could be related with losses of permease activity or decrease of enzymatic activity responsible for the conversion of 5FC into FMUP. This process was observed in *C. glabrata*, while mutational losses of one of the pyrimidine salvage enzymes and decrease in UPRTase in a gene dose-dependent manner were observed in *C. albicans* strains, resistant to 5FC (Ghannoum and Rice, 1999).

Lastly, echinocandins (i.e. caspofungin, anidulofungin, and micafungin) are the latest and, at the same time, the first new class of antifungal drugs. The main target of this drug is the fungal cell wall, acting as inhibitors of the α -1,3-D-glucan biosynthesis by blocking of β -1,3-D-glucan synthase (Perlin, 2014; Sanglard and Odds, 2002). Despite a higher level of efficiency associated to this class of antifungal drugs, exerting rapid fungicidal effects against *Candida* species, including azole-resistance species (Kontoyiannis and Lewis, 2002), in the last years, it has been observed an increasing level of resistance, namely for *C. albicans*, *C. glabrata*, *C. krusei* and *C. parapsilosis* (Kanafani and Perfect, 2008; Perlin, 2009). The involved resistance mechanisms were not completely elucidated and other are still unknown; however, drug tolerance conferred by adaptive cellular physiology to the environmental stress, as well as genetic mutations and amino acids substitutions are the main supposed contributors associated with *Candida* species echinocandin-resistant (Perlin, 2009).

Nevertheless, new perspectives of development of new orally effective drugs are open and different experiments have been made (Kanafani and Perfect, 2008; Perlin, 2014, 2009; Sanglard, 2002). So, considering all the advances on antifungal drugs research, it is necessary to take into account the safety, tolerability, efficacy, toxicity, and even side effects of these substances/chemical molecules. In this sense, other alternatives safer than the current antifungal drugs are necessary.

3. Active principles of plants

3.1. Plants as a rich source of bioactive compounds

Plants have been used for several generations in traditional herbal medicine, as first choice treatment. The first civilizations established the therapeutic potential of plants taking into account the more or less prominent effects observed in the body. They also observed that certain plants evidenced toxic effects according to the dose. However, they did not know the reasons that provided those effects, as well as, the nature of plant constituents. Thus, much of the knowledge was empirical, passing from generation to generation, but lacking of a solid foundation of the use and pre-

scription, mostly due to the low literacy and written information (Halberstein, 2005; Murray and Pizzorno, 1998; Petrovska, 2012).

Currently, it is well recognized that hundreds of biological active chemical compounds are present, in each plant, working in synergism, and conferring a broad variety of bioactivities, being subject of an increasing scientific interest. More and more, people worldwide use medicinal plants as first priority to maintain a good health, to prevent or even treat some health conditions. Linked to this, some studies have been conducted, using a wide variety of plants, towards the identification of bioactive molecules, their properties and mechanisms of action (Agarwal et al., 2010; Alves-Silva et al., 2013; Asgarpanah and Kazemivash, 2012; Asl and Hosseinzadeh, 2008; Bakkali et al., 2008; Kanafani and Perfect, 2008; Rana et al., 2011; Sher, 2009; Shojaii and Fard, 2012; Silva et al., 2011b; Singh et al., 2010). In fact, natural matrices, in particular, plants are extremely rich sources of biomolecules that have sparked an increasing interest for scientific researchers, in different areas of knowledge. Until now, several chemical compounds have been identified, but only a few properties have been clarified, with many remaining undefined. Nevertheless, many compounds are still unknown, as also the synergic interactions among them (Carocho and Ferreira, 2013a; Choudhary and Atta-ur-Rahman, 1999; Devasagayam et al., 2004; Rubió et al., 2013).

It is truly believed that the biologically active chemical compounds present in plants act synergistically and confer, both direct and indirect effects, to the human body. The direct impact is commonly due to their phytopharmacological action, while the indirect impact is linked to the simultaneous interaction with other plants or drugs. Furthermore, it should be highlighted that the fact that one plant from a particular genus contains a specific chemical composition and a significant medicinal value does not mean that all the other plants from the same genus had the same value. Even some studies report that the same species, but with different origins might have considerable differences on chemical composition and, therefore, a different efficacy and potency in the bioactivity (Farhat et al., 2009, 2013; Ghasemi Pirbalouti et al., 2013; Papageorgiou et al., 2008; Politeo et al., 2006).

In the modern medical classification, a plant acquire the “medicinal” status when it presents several pharmacologically active compounds, making possible its use at a therapeutic level, conferring not only benefits in the treatment of some conditions, but also in prevention/prophylaxis (Balch, 2006; Murray, 2004; Murray and Pizzorno, 1998). Notwithstanding, other chemical compounds present in trace amounts could influence the action of the main molecules, which explains the higher potential of using the whole natural matrix instead of an isolate chemical compound (Ettfagh et al., 2011; Junio et al., 2011).

Thus, it is very important to provide a correct and complete characterization of the bioactive compounds present in medicinal plants, in order to complement and unify the prescriptions, and at the same time, to establish the amount/dosage of active principles needed to treat/avert various health problems. Currently, due to the advances of the analytical methods, it was possible to elucidate the structure of thousands of phytochemicals, either primary or secondary metabolites (Fig. 1).

3.2. Primary metabolites

Primary metabolites comprise the organic compounds derived from the primary metabolism of plants; they are often concentrated in seeds and/or vegetative organs and play an important role in the physiological development and basic cell metabolism. In general, these compounds are not limited to the plant kingdom, being even used in several biosynthetic pathways (Nelson and Cox, 2000).

Higher plants are able to synthesize organic compounds and produce oxygen from the light, water and carbon dioxide, which

Fig. 1. Relationship between the primary and secondary metabolism in plants.

are the main energetic and food sources for animals. Furthermore, these same organic compounds are also an energetic source for biosynthetic organisms. This process, in which autotrophic organisms produce organic compounds according to their needs is commonly known as photosynthesis: light phase and dark phase. Organic compounds are synthesized in the dark phase, through some biosynthetic pathways (Nelson and Cox, 2000). Proteins, lipids, carbohydrates and chlorophyll are the main compounds derived from the primary metabolism (Fig. 1).

Besides many other physiological functions, proteins and even some free amino acids are considered crucial transport elements of a wide variety of molecules; integrate the genetic material binding to nucleic acids; include enzymes play an important reserve role and confer high protection against invaders (Nelson and Cox, 2000; Proença da Cunha et al., 2010, 2012). On the other hand, lipids, commonly defined as water insoluble compounds but soluble in organic solvents play important organic benefits as structural components of cellular membranes; energetic reserve; protective function; hormonal regulators and synthesizers of inflammatory immunomodulators (eicosanoids: prostaglandins, thromboxanes and leukotrienes), among other crucial functions in living organisms (Campos, 2008; Nelson and Cox, 2000).

Carbohydrates, or glycids, represent the basis of dietary food of a majority of living organisms. According to the length and the type of glycosidic linkages, these molecules play different biological properties and have innumerable applications. One of the most common is their energetic cellular reservoir function, being accumulated in the main reserve organs of plants – seeds, grains and tubers, as well as leaves. The latest have a limited storage capacity (Campos, 2008; Nelson and Cox, 2000; Proença da Cunha et al., 2010).

Lastly, chlorophyll, a green pigment mainly present in the higher plant protoplasts, but also in red algae and purple bacteria, essentially exerts functions of photosynthetic pigments. Despite the existence of two types of chlorophyll, *a* and *b*, both of them are involved in organic compounds synthesis and, therefore, in the primary function of plants: nutrition (Campos, 2008; Nelson and Cox, 2000).

In fact, primary metabolites are much more than merely macronutrients. Some of the bioactive properties reported for plants, such as antioxidant, antimicrobial, anti-inflammatory, and antitumor activities have been related with primary metabolites, mainly lipids (polyunsaturated fatty acids, sterols), proteins, carotenoids, vitamins and carbohydrates (polysaccharides and glycosides) (Ksouri et al., 2012).

3.3. Secondary metabolites

Secondary metabolites are organic compounds that derived from the primary metabolites of plants. It is well established that these metabolites play specialized functions in a wide variety of living organisms, being even considered as biologically active compounds. They are renowned as possess higher value than the primary metabolites, in relation to the life expectancy, wellbeing

and even to prevent/avert some diseases/disorders (Nelson and Cox, 2000; Rubió et al., 2013).

Terpenes are a class of secondary metabolites formed by several isoprene units, i.e. several combinations of 5-carbon-base (C5). Different combinations of the isoprene units originate structurally and functionally different classes of terpenes (Rubió et al., 2013). Thus, they could occur as monoterpenes (C10), diterpenes (C20), triterpenes (C30), tetraterpenes (C40), but also hemiterpenes (C5) and sesquiterpenes (C15). Monoterpenes and diterpenes are the main classes of terpenes present on herbs and spices. Furthermore, these compounds have important biological activities, including antimicrobial properties against a wide variety of pathogens such as *Candida* species (Alves et al., 2013; Rubió et al., 2013; Saleem et al., 2010).

Steroids are secondary metabolites derived from the tetracyclic triterpenes (C30), composed by isoprenoid C5 units derived from isopentenyl (3-methylbut-3-en-1-yl) pyrophosphate, and presenting a characteristic branched chain structure. Stigmasterol has been reported as one of the most abundant steroids in plant oils. These compounds are commonly highly degraded into other products, namely steroid hormones that play important organic functions (Carocho and Ferreira, 2013b; Nelson and Cox, 2000). Moreover, the antimicrobial activity of steroids has also been described, including the antifungal properties against some pathogens such as *Candida* species (Alves et al., 2013; Saleem et al., 2010; Sanglard, 2002). One of the proposed mechanisms of action is their ability to disrupt membrane integrity (Saleem et al., 2010), but others remain unclear and even unknown.

Alkaloids comprise a diverse group of heterocyclic nitrogen substances, being commonly considered high toxic molecules that play an important defensive role, but reflecting a general biological principle of poisonous plants (Inbaraj et al., 2001, 2006; Sertel et al., 2011). Nevertheless, in some situations, their toxicity is beneficial; for example, Sertel et al. (2011) reported that vinblastine, vincristine (from *Catharanthus roseus*, formerly: *Vinca rosea*) and semisynthetic vindesine and vinorelbine, presented high cytotoxicity against cancer cells, highlighting the potential of these dimeric indole alkaloids for cancer, leukemia and lymphomas treatments. The efficacy of alkaloids against bacteria, parasites, dermatophytes and fungi, including *Candida* species has also been described (Ali-Shtayeh and Ghdeib, 1999; Arif et al., 2011, 2009; Carocho and Ferreira, 2013b; Dzoyem et al., 2014; Rose, 1999; Sher, 2009; Silva and Fernandes Júnior, 2010; Wyk et al., 2009; Xie et al., 2012). The intercalation into cell wall and/or DNA is considered one of the main described mechanisms of bioactive action of alkaloids (Silva and Fernandes Júnior, 2010).

Phenolic compounds constitute a major class of plant secondary metabolites being broadly distributed, with more than 8000 phenolic structures currently identified. These compounds can contain one or more aromatic rings with one or more hydroxyl groups (Dai and Mumper, 2010). In fact, the term “phenolic” or “polyphenol” can be chemically defined as a substance which possesses an aromatic ring bearing at least one (phenol) or more (polyphenol)

Fig. 2. Main classes and subclasses of phenolic compounds.

nol) hydroxyl substituent, including functional derivatives (esters, methyl ethers, glycosides, etc.), ranging from simple molecules, such as phenolic acids, to high complex structures, such as tannins (Dai and Mumper, 2010; Santos-Buelga et al., 2012). In the modern classification, the broad category of phenolics is divided into several groups, according to the number of phenol subunits present: polyphenols and simple phenols. Thus, polyphenols possessing at least two phenol subunits are classified as flavonoids, while those compounds which possess three or more phenol subunits are included in tannins (hydrolysable and non-hydrolysable) (Giada, 2013; Robbins, 2003). Phenolic acids, flavonoids and tannins are the most common, being stilbenes and lignans less common (Dai and Mumper, 2010). Curiously, these classes of phenolic compounds are currently the main subjects of scientific researches. But, beyond these, other phenolic compounds, like chalcones, coumarins, xanthenes and lignins are also present in natural matrices (Fig. 2).

Several researchers have demonstrated the antioxidant potential of phenolic compounds (Çekiç et al., 2013; Rice-Evans et al., 1996; Walch et al., 2011), but currently other bioactive properties have also been studied. Curiously, some of these biological properties are directly linked to the antioxidant activity. It is a fact, that the chemical structure of phenolic compounds determine their properties as well as mechanism of action, being conferred by different ways (Giada, 2013). Nevertheless, and not less important is the fact that despite reactive oxygen and nitrogen species (ROS/RNS) at physiological concentrations may be required for normal cell functioning and to play an important role in plant and animal evolutions, these species are able to attack biological macromolecules (e.g. cellular DNA), causing a wide variety of cellular damages, disorders and even diseases; to disestablish of the defense systems increasing the vulnerability to infections by pathogens, parasites and other invaders (Carocho and Ferreira, 2013a; Çekiç et al., 2013; Chaturvedi and Beal, 2013; Devasagayam et al., 2004; Valko et al., 2007).

4. Activity of phenolic extracts/compounds against *Candida* species

4.1. The use of phenolic extracts against *Candida* species

Several studies have been performed in order to evaluate the antifungal potential of polar extracts from plant origin, which are enriched in phenolic compounds, such as acetone (Table 1), aqueous (Table 2), ethanolic (Table 3), methanol–dichloromethane (Table 4), methanolic (Table 5), hydroalcoholic (methanol:or ethanol:water mixtures) (Table 6), DMSO, diethyl ether and ethyl acetate extracts, and tinctures (Table 7). The aqueous extracts were the most studied preparations, followed by methanolic and ethanolic extracts.

Different techniques/assays have been used for that purpose. Agar disc diffusion and broth dilution (microdilution and macrodilution) methods are the most commonly used, being, normally, performed according to the Clinical and Laboratory Standards Institute (CLSI), with some modifications. Both of them are considered antimicrobial susceptibility tests, expressing the results as halo inhibition zones or minimal inhibitory (MIC) and minimal fungicidal (MFC) concentrations, respectively. The most tested *Candida* species were *C. albicans*, *C. tropicalis*, *C. krusei*, *C. glabrata*, *C. parapsilosis*, *C. lusitanae* and *C. guilliermondii*, being considered the main species associated to opportunistic fungal infections (Vázquez-González et al., 2013).

In general, the plant extracts obtained using methanol and water, as solvent extraction, and mixtures of the previous (hydroalcoholic extracts) were more effective against *Candida* species than the other extracts prepared with a unique solvent. Thus, the methanol (Table 5), aqueous (Table 2) and hydroalcoholic extracts (ethanol:water and methanol:water mixtures) (Table 6) presented lower MICs than the extracts obtained with acetone (Table 1), methanol–dichloromethane (Table 4) ethanol (Table 3) or other

Table 1
Activity of acetone extracts against *Candida* species.

Plant source	<i>Candida</i> species	MIC	References
<i>Combretum albopunctatum</i>	<i>C. albicans</i>	0.64 mg/mL	Masoko et al. (2010)
<i>Combretum imberbe</i>	<i>C. albicans</i>	2.5 mg/mL	Masoko et al. (2010)
<i>Combretum nelsonii</i>	<i>C. albicans</i>	0.04 mg/mL	Masoko et al. (2010)
<i>Origanum vulgare</i>	<i>C. albicans</i>	5 mg/mL	Ličina et al. (2013)
<i>Paullinia cupana</i>	<i>C. albicans</i>	250 µm/mL	Basile et al. (2013)
<i>Sclerocarya birrea</i>	<i>C. parapsilosis</i>	0.32 mg/mL	Masoko et al. (2008)
<i>Terminalia sericea</i>	<i>C. albicans</i>	0.64 mg/mL	Masoko et al. (2010)

MIC – minimum inhibitory concentration.

Table 2
Activity of aqueous extracts against *Candida* species.

Plant source	<i>Candida</i> species	MIC	References
<i>Acorus calamus</i>	<i>C. albicans</i>	12.5 mg/mL	Webster et al. (2008)
	<i>C. krusei</i>	12.5 mg/mL	
	<i>C. lusitaniae</i>	12.5 mg/mL	
<i>Adiantum capillus veneris</i> <i>Allium sativum</i>	<i>C. parapsilosis</i>	12.5 mg/mL	Ishaq et al. (2014) Shamim et al. (2004) Fei et al. (2008) Pathak (2012) Pathak (2012) Shamim et al. (2004) Pathak (2012) Fei et al. (2008) Pathak (2012) Shamim et al. (2004) Fei et al. (2008) Pathak (2012) Shamim et al. (2004) Fei et al. (2008) Pathak (2012)
	<i>C. albicans</i>	1 mg/mL ^a	
	<i>C. albicans</i>	0.2 mg/mL	
		100 mg/mL ^a	
		1 mg/mL ^a	
	<i>C. dublinensis</i>	1 mg/mL ^a	
	<i>C. grabrata</i>	0.2 mg/mL	
		1 mg/mL ^a	
	<i>C. krusei</i>	100 mg/mL ^a	
		1 mg/mL ^a	
	<i>C. tropicalis</i>	0.2 mg/mL	
		100 mg/mL ^a	
		1 mg/mL ^a	
		12.5 mg/mL	
		25 mg/mL	
<i>Alnus viridis</i>	<i>C. glabrata</i>	25 mg/mL	Webster et al. (2008)
	<i>C. krusei</i>	0.2 mg/mL	
	<i>C. lusitaniae</i>	6 mg/mL	
	<i>C. parapsilosis</i>	25 mg/mL	
	<i>C. tropicalis</i>	3 mg/mL	
<i>Aloe barbadensis</i>	<i>C. albicans</i>	1 mg/mL	Shamim et al. (2004)
	<i>C. grabrata</i>	1 mg/mL	
	<i>C. tropicalis</i>	0.2 mg/mL	
<i>Aloe excelsa</i>	<i>C. albicans</i>	1 mg/mL	Coopoosamy and Magwa (2007)
	<i>C. tropicalis</i>	0.2 mg/mL	
		10 mg/mL ^a	
<i>Arbutus unedo</i> <i>Arctium minus</i>	<i>C. albicans</i>	12.5 mg/mL	Dib et al. (2010) Lubian et al. (2010)
	<i>C. dubliniensis</i>	12.5 mg/mL	
	<i>C. glabrata</i>	12.5 mg/mL	
	<i>C. krusei</i>	12.5 mg/mL	
	<i>C. stellatoidea</i>	12.5 mg/mL	
	<i>C. tropicalis</i>	25 mg/mL	
		12.5 mg/mL	
	<i>C. glabrata</i>	25 mg/mL	
	<i>C. krusei</i>	50 mg/mL	
<i>Artemisia campestris</i>	<i>C. lusitaniae</i>	25 mg/mL	Webster et al. (2008)
	<i>C. parapsilosis</i>	50 mg/mL	
	<i>C. tropicalis</i>	25 mg/mL	
	<i>C. glabrata</i>	0.4 mg/mL	
	<i>C. krusei</i>	6 mg/mL	
<i>Artemisia frigida</i>	<i>C. lusitaniae</i>	12.5 mg/mL	Webster et al. (2008)
	<i>C. parapsilosis</i>	50 mg/mL	
	<i>C. tropicalis</i>	6 mg/mL	
	<i>C. albicans</i>	1 mg/mL ^a	
	<i>C. dubliniensis</i>	1 mg/mL ^a	
<i>Azadirachta indica</i>	<i>C. glabrata</i>	1 mg/mL ^a	Pathak (2012)
	<i>C. krusei</i>	1 mg/mL ^a	
	<i>C. tropicalis</i>	1 mg/mL ^a	
	<i>C. albicans</i>	0.8 mg/mL	
	<i>C. glabrata</i>	0.05 mg/mL	
<i>Betula alleghaniensis</i>	<i>C. krusei</i>	0.4 mg/mL	Webster et al. (2008)
	<i>C. lusitaniae</i>	0.4 mg/mL	
	<i>C. parapsilosis</i>	3 mg/mL	
		0.05 mg/mL	
	<i>C. tropicalis</i>	6 mg/mL	
<i>Brucea javanica</i>	<i>C. albicans</i>	50 mg/mL	Nordin et al. (2013)
	<i>C. dubliniensis</i>	3.13 mg/mL	
	<i>C. glabrata</i>	25 mg/mL	
	<i>C. krusei</i>	50 mg/mL	
	<i>C. lusitaniae</i>	25 mg/mL	
	<i>C. parapsilosis</i>	25 mg/mL	
	<i>C. tropicalis</i>	50 mg/mL	
	<i>C. albicans</i>	26.90 mg	
	<i>C. albicans</i>	10 mg/mL ^a	
<i>Cassia alata</i> <i>Camellia sinensis</i> <i>Cinnamomum</i> sp.	<i>C. albicans</i>	200 mg/mL ^a	Timothy et al. (2012) Doddanna et al. (2013) Qader et al. (2013)
	<i>C. dubliniensis</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
	<i>C. tropicalis</i>	200 mg/mL ^a	
<i>Emblica officinalis</i>	<i>C. albicans</i>	200 mg/mL ^a	Saeed and Tariq (2007)
		200 mg/mL ^a	

Table 2 (Continued)

Plant source	Candida species	MIC	References
<i>Epilobium angustifolium</i>	<i>C. albicans</i>	0.2 mg/mL 0.4 mg/mL 0.4 mg/mL 0.4 mg/mL	Webster et al. (2008)
	<i>C. glabrata</i>	0.025 mg/mL	
	<i>C. krusei</i>	0.2 mg/mL	
	<i>C. lusitaniae</i>	0.05 mg/mL	
	<i>C. parapsilosis</i>	0.4 mg/mL 0.1 mg/mL	
	<i>C. tropicalis</i>	1.6 mg/mL	
<i>Eucalyptus sideroxylon</i>	<i>C. albicans</i>	400 mg/mL ^a	Ashour (2008)
<i>Eucalyptus torquata</i>	<i>C. albicans</i>	400 mg/mL ^a	Ashour (2008)
<i>Fragaria virginiana</i>	<i>C. albicans</i>	0.8 mg/mL	Webster et al. (2008)
	<i>C. glabrata</i>	0.05 mg/mL	
	<i>C. krusei</i>	0.05 mg/mL	
	<i>C. lusitaniae</i>	0.1 mg/mL	
	<i>C. parapsilosis</i>	0.2 mg/mL	
	<i>C. tropicalis</i>	0.8 mg/mL	
<i>Funtumia elastica</i>	<i>C. albicans</i>	100 mg/mL ^a	Adekunle and Ikumapayi (2006)
<i>Hyssopus officinalis</i> subsp. <i>pilifer</i>	<i>C. albicans</i>	7.00 mg/mL	Džamić et al. (2013)
<i>Lonicera japonica</i>	<i>C. albicans</i>	Viability (%): 81.48	Wong and Tsang (2009)
<i>Mallotus oppositifolius</i>	<i>C. albicans</i>	100 mg/mL ^a	Adekunle and Ikumapayi (2006)
<i>Morinda citrifolia</i>	<i>C. albicans</i>	40 mg/mL	Jaikittivong et al. (2009)
<i>Morinda morindoides</i>	<i>C. albicans</i>	62.50 mg/mL	Touré et al. (2011)
<i>Murraya koenigii</i>	<i>C. albicans</i>	1 mg/mL ^a	Pathak (2012)
	<i>C. dublinensis</i>	1 mg/mL ^a	
	<i>C. glabrata</i>	1 mg/mL ^a	
	<i>C. krusei</i>	1 mg/mL ^a	
	<i>C. tropicalis</i>	1 mg/mL ^a	
<i>Ocimum sanctum</i>	<i>C. glabrata</i>	1 mg/mL ^a	Pathak (2012)
<i>Origanum vulgare</i>	<i>C. albicans</i>	200 mg/mL ^a 20 mg/mL	Abdul et al. (2012) Ličina et al. (2013)
<i>Paeonia suffruticosa</i>	<i>C. albicans</i>	Viability (%): 100.53	Wong and Tsang (2009)
<i>Plantago media</i>	<i>C. albicans</i>	200 mg/mL ^a	Qader et al. (2013)
	<i>C. dublicans</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
	<i>C. tropicalis</i>	200 mg/mL ^a	
<i>Populus tremuloides</i>	<i>C. albicans</i>	1.6 mg/mL	Webster et al. (2008)
	<i>C. glabrata</i>	0.025 mg/mL	
	<i>C. krusei</i>	0.2 mg/mL	
	<i>C. lusitaniae</i>	0.2 mg/mL	
	<i>C. parapsilosis</i>	0.4 mg/mL	
	<i>C. tropicalis</i>	1.6 mg/mL	
<i>Punica granatum</i>	<i>C. albicans</i>	200 mg/mL ^a	Qader et al. (2013)
	<i>C. dublicans</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
<i>Quercus infectoria</i>	<i>C. albicans</i>	200 mg/mL ^a	Qader et al. (2013)
	<i>C. dublicans</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
	<i>C. tropicalis</i>	200 mg/mL ^a	
<i>Rhus angustifolia</i>	<i>C. albicans</i>	200 mg/mL ^a	Qader et al. (2013)
	<i>C. dublicans</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
	<i>C. tropicalis</i>	200 mg/mL ^a	
<i>Ribes nigrum</i>	<i>C. glabrata</i>	3.85 mg/mL	Krisch et al. (2008)
	<i>C. inconspicua</i>	2.82 mg/mL	
	<i>C. lipolytica</i>	5.15 mg/mL	
	<i>C. norvegica</i>	3.62 mg/mL	
	<i>C. parapsilosis</i>	2.76 mg/mL	
	<i>C. tropicalis</i>	3.75 mg/mL	
	<i>C. zeylanoides</i>	1.83 mg/mL	
<i>Rosmarinus officinalis</i>	<i>C. albicans</i>	300 mg/mL ^a	Abdul et al. (2012)
<i>Salvia officinalis</i>	<i>C. albicans</i>	15 mg/mL	Jasim and Al-khaliq (2011)
	<i>C. parapsilosis</i>	2.5 mg/mL	Martins et al. (2015)
	<i>C. tropicalis</i>	1.25 mg/mL	(Martins et al., 2015)
<i>Sapindus saponaria</i>	<i>C. parapsilosis</i>	0.16 mg/mL	Tsuzuki et al. (2007)
<i>Scutellaria baicalensis</i>	<i>C. albicans</i>	2.5 mg/mL 5 mg/mL	Wong and Tsang (2009)
<i>Solanum nigrum</i>	<i>C. albicans</i>	0.2 mg/mL	Shamim et al. (2004)
	<i>C. glabrata</i>	1 mg/mL	
	<i>C. tropicalis</i>	0.2 mg/mL	

Table 2 (Continued)

Plant source	<i>Candida</i> species	MIC	References
<i>Solidago gigantea</i>	<i>C. albicans</i>	0.8 mg/mL	Webster et al. (2008)
	<i>C. glabrata</i>	0.1 mg/mL	
	<i>C. krusei</i>	0.8 mg/mL	
	<i>C. lusitaniae</i>	1.6 mg/mL	
	<i>C. tropicalis</i>	12.5 mg/mL	
<i>Sonneratia alba</i>	<i>C. albicans</i>	0.25 mg/mL	Kaewpiboon et al. (2012) Kaewpiboon et al. (2012) (Kim and Lee (2012)
<i>Sonneratia caseolaris</i>	<i>C. albicans</i>	0.125 mg/mL	
<i>Syzygium aromaticum</i>	<i>C. albicans</i>	0.782 mg/mL	
<i>Thymus kotschyana</i>	<i>C. glabrata</i>	0.025 mg/mL	Qader et al. (2013)
	<i>C. krusei</i>	0.098 mg/mL	
	<i>C. tropicalis</i>	0.003125 mg/mL	
	<i>C. albicans</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
	<i>C. tropicalis</i>	200 mg/mL ^a	
<i>Thymus vulgaris</i>	<i>C. albicans</i>	300 mg/mL ^a	Abdul et al. (2012)
<i>Zingiber officinalis</i>	<i>C. albicans</i>	200 mg/mL ^a	Qader et al. (2013)
	<i>C. dublicans</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
	<i>C. tropicalis</i>	200 mg/mL ^a	

MIC – minimum inhibitory concentration.

^a Concentration used to evaluate antifungal activity by disc diffusion.

extracts (Table 7). For example, hydroalcoholic extracts of *Cassia fistula* (MIC = 0.025 mg/mL), *Embllica officinalis* (MIC = 0.1024 mg/mL), *Glycyrrhiza glabra* (MIC = 0.512 mg/mL) and *Punica granatum* (MIC = 0.0039 mg/mL) presented a higher antifungal effect, against *C. albicans*, than the extracts of the same plant species prepared with a single solvent. The only exceptions were the aqueous extracts of *Sapindus saponaria* (MIC = 0.16 mg/mL), against *C. parapsilosis*, and *Syzygium aromaticum* (MIC = 0.003125 mg/mL), against *C. tropicalis*, that gave a higher effect than its hydroalcoholic extracts, and the methanolic extract of *Salvia officinalis* (MIC = 0.16 mg/mL) that evidenced a higher antifungal potential, against *C. tropicalis*, than the hydroalcoholic (MIC = 1.25 mg/mL) and aqueous (MIC = 1.25 mg/mL) extracts. Notwithstanding, the number of studies that report antifungal properties of hydroalcoholic extracts is lower than the studies using methanol, ethanol or water.

Comparing the antifungal potential of the extracts prepared with a unique solvent, the methanolic extracts (Table 5) showed the lowest MICs for the majority of the *Candida* species, followed by aqueous (Table 2) and ethanolic (Table 3) extracts. In particular, *Adiantum capillus veneris*, *Artemisia herba-alba*, *Caparis spinosa*, *Eucalyptus sideroxylon*, *E. torquata*, *Glycyrrhiza glabra*, *Laurus nobilis*, *Pimpinella anisum* and *Salvia officinalis* methanolic extracts showed higher antifungal potential than their aqueous and ethanolic extracts. On the other hand, *Acorus calamus*, *Funtumia elastica*, *Hyssopus officinalis* subsp. *pilifer*, *Murraya koenigii*, *Ribes nigrum* and *Zingiber officinalis* aqueous extracts evidenced most pronounced antifungal effects than their ethanolic and methanolic extracts. *Allium sativum*, *Aloe barbadensis*, *Aloe excelsa*, *Cinnamomum* spp. and *Solanum nigrum* aqueous and ethanolic extracts evidenced similar potential against *Candida* species. Lastly, *Camellia sinensis*, *Cassia alata*, *Mallotus oppositifolius*, *Morinda morindoides*, *Origanum vulgare*, *Plantago media*, *Punica granatum*, *Quercus infectoria*, *Rhus angustifolia*, *Sclerocarya birrea* and *Thymus kotschyana* ethanolic extracts evidenced higher antifungal potential than aqueous and methanolic extracts.

In specific plant species, the tendency described was modified. For example, the methanol-dichloromethane extract of *Andrographis paniculata* showed a higher antifungal potential than their ethanolic and methanolic extracts; ethanol, acetone, diethyl ether and ethyl acetate extracts of *Origanum vulgare*, as well as aqueous and ethyl acetate extracts of *Hyssopus officinalis* subsp.

Pilifer and aqueous and hydroalcoholic extracts of *Salvia officinalis* showed similar antifungal effects. Furthermore, *Crossandra infundibuliformis* ethyl acetate and methanol extracts showed similar effects, whereas *Gonzalagunia rosea* and *Montanoa* methanolic extracts showed higher antifungal potential than the corresponding methanol:dichloromethane extracts; the acetone extract of *Paullinia cupana* seems to be more effective than the ethanolic extract against *C. albicans*, but due to the differences in the units used to express the results, it is difficult to compare both values.

Analysing the antifungal activity against specific *Candida* species, hydroalcoholic extracts showed the most pronounced properties against *C. albicans*, followed by methanolic and methanol–dichloromethane extracts. *Punica granatum* hydroalcoholic (ethanol:water, 50:50 v/v) extract revealed the highest antifungal activity (MIC = 0.0039 mg/mL). In general, methanolic extracts followed by aqueous and hydroalcoholic extracts, showed a higher antifungal potential against *C. glabrata* than the ethanolic and methanol–dichloromethane extracts. *Dorstenia turbinata* (0.039 mg/mL) and *Thymbra spicata* (0.01 mg/mL) methanolic extracts evidenced the most pronounced effects. For *C. guilliermondii*, the hydroalcoholic extract of *Syzygium aromaticum* (MICs = 0.05 mg/mL) was the most effective preparations, but *Punica granatum* ethanolic extract (MIC = 0.125 mg/mL) also evidenced a lower MIC value. For *C. krusei*, the methanolic extracts were the most efficient plant extract preparations, followed by ethanolic and aqueous extracts. *Thymbra spicata* (MIC = 0.01 mg/mL) and *Laurus nobilis* (MIC = 0.07 mg/mL) comprises the methanolic extracts that evidenced highest antifungal activity against *C. krusei*. Concerning *C. lusitaniae*, aqueous extracts presented the highest antifungal potential, followed by ethanolic, methanol–dichloromethane and ethyl acetate extracts. *Epilobium augustifolium* aqueous extract (MIC = 0.05 mg/mL) revealed to be the best against *C. lusitaniae*. On the other hand, to the *C. parapsilosis*, the hydroalcoholic extracts were the most effective preparations, followed by methanolic, aqueous and ethanolic extracts. *Punica granatum* hydroalcoholic extract (MIC = 0.0039 mg/mL) presented the most pronounced antifungal potential against *C. parapsilosis*. Lastly, aqueous extracts were the most efficient plant extract preparations against *C. tropicalis*, followed by methanolic, methanol–dichloromethane and hydroalcoholic extracts. *Syzygium aromaticum* (MIC = 0.003125 mg/mL) aqueous extract revealing the highest potential.

Table 3
Activity of ethanolic extracts against *Candida* species.

Plant source	<i>Candida</i> species	MIC	References
<i>Achillea biebersteinii</i>	<i>C. albicans</i>	100 mg/mL	Hassawi and Kharma (2006)
<i>Achillea fragrantissima</i>	<i>C. albicans</i>	150 mg/mL	Hassawi and Kharma (2006)
<i>Acorus calamus</i>	<i>C. albicans</i>	28.80 mg/mL	Thirach et al. (2003)
<i>Adiantum capillus veneris</i>	<i>C. albicans</i>	1 mg/mL ^a	Ishaq et al. (2014)
<i>Allium cepa</i>	<i>C. albicans</i>	10 mg/mL ^a	Doddanna et al. (2013)
<i>Allium sativum</i>	<i>C. albicans</i>	1 mg/mL ^a	Joe et al. (2009)
<i>Allium sativum</i>	<i>C. albicans</i>	0.2 mg/mL	Shamim et al. (2004)
	<i>C. grabrata</i>	0.2 mg/mL	
	<i>C. tropicalis</i>	0.2 mg/mL	
<i>Aloe barbadensis</i>	<i>C. albicans</i>	1 mg/mL	Shamim et al. (2004)
	<i>C. grabrata</i>	1 mg/mL	
	<i>C. tropicalis</i>	0.2 mg/mL	
<i>Aloe excelsa</i>	<i>C. albicans</i>	1 mg/mL	Coopoosamy and Magwa (2007)
	<i>C. tropicalis</i>	0.2 mg/mL	Doddanna et al. (2013)
<i>Aloe vera</i>	<i>C. albicans</i>	10 mg/mL ^a	Wong and Tsang (2009)
<i>Andrographis paniculata</i>	<i>C. albicans</i>	Viability: 103.85%	Hassawi and Kharma (2006)
<i>Anthemis pseudocotula</i>	<i>C. albicans</i>	150 mg/mL	Hassawi and Kharma (2006)
<i>Artemisia herba-alba</i>	<i>C. albicans</i>	100 mg/mL	Hashem (2011)
<i>Ballota undulate</i>	<i>C. albicans</i>	25 mg/mL	Doddanna et al. (2013)
<i>Camellia sinensis</i>	<i>C. albicans</i>	10 mg/mL ^a	Mandeel and Taha (2005)
<i>Capparis spinosa</i>	<i>C. albicans</i>	100 mg/mL ^a	Ertürk (2006)
<i>Capsicum annum</i>	<i>C. albicans</i>	17.5 mg/mL	Timothy et al. (2012)
<i>Cassia alata</i>	<i>C. albicans</i>	5.60 mg	Prabhakar et al. (2008)
<i>Cassia siamea</i>	<i>C. albicans</i>	100 mg/mL	
	<i>C. glabrata</i>	100 mg/mL	
	<i>C. guilliermondii</i>	100 mg/mL	
	<i>C. krusei</i>	100 mg/mL	
	<i>C. parapsilosis</i>	100 mg/mL	
	<i>C. tropicalis</i>	100 mg/mL	
<i>Caulerpa scalpelliformis</i>	<i>C. albicans</i>	100 mg/mL	Prabhakar et al. (2008)
	<i>C. glabrata</i>	100 mg/mL	
	<i>C. guilliermondii</i>	100 mg/mL	
	<i>C. krusei</i>	100 mg/mL	
	<i>C. parapsilosis</i>	100 mg/mL	
	<i>C. tropicalis</i>	100 mg/mL	
<i>Cinnamomum</i> sp.	<i>C. albicans</i>	200 mg/mL ^a	Qader et al. (2013)
	<i>C. dubicans</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
	<i>C. tropicalis</i>	200 mg/mL ^a	
<i>Cleome amplyocarpa</i>	<i>C. albicans</i>	75 mg/mL	Hashem (2011)
<i>Colutea arborescens</i>	<i>C. albicans</i>	12.5 mg/mL	Ertürk (2006)
<i>Convolvulus arvensis</i>	<i>C. albicans</i>	100 mg/mL ^a	Mandeel and Taha (2005)
<i>Convolvulus arvensis</i>	<i>C. albicans</i>	150 mg/mL	Hassawi and Kharma (2006)
<i>Coscinium fenestratum</i>	<i>C. albicans</i>	0.5 mg/mL	Kaewpiroon et al. (2012)
<i>Crataegus oxyacantha</i>	<i>C. albicans</i>	10 mg/mL ^a	Kostić et al. (2012)
<i>Cressa cretica</i>	<i>C. albicans</i>	100 mg/mL ^a	Mandeel and Taha (2005)
<i>Cuminum cyminum</i>	<i>C. albicans</i>	15 mg/mL	Ertürk (2006)
<i>Cynara scolymus</i>	<i>C. albicans</i>	12.5 mg/mL	Zhu et al. (2005)
	<i>C. lusitaniae</i>	12.5 mg/mL	
<i>Dianthus coryophyllum</i>	<i>C. albicans</i>	20 mg/mL	Ertürk (2006)
<i>Emex spinosa</i>	<i>C. albicans</i>	100 mg/mL ^a	Mandeel and Taha (2005)
<i>Erica arborea</i>	<i>C. albicans</i>	2.5 mg/mL	Ertürk (2006)
<i>Eugenia caryophyllus</i>	<i>C. albicans</i>	17.41 mg/mL	Thirach et al. (2003)
<i>Eugenia uniflora</i>	<i>C. krusei</i>	0.25 mg/mL	Correa-Royero et al. (2010)
<i>Fagonia indica</i>	<i>C. albicans</i>	100 mg/mL ^a	Mandeel and Taha (2005)
<i>Fraxinus rhynchophylla</i>	<i>C. albicans</i>	Viability: 77.56%	Wong and Tsang (2009)
<i>Funtumia elastica</i>	<i>C. albicans</i>	100 mg/mL ^a	Adekunle and Ikumapayi (2006)
<i>Glycyrrhiza glabra</i>	<i>C. albicans</i>	1 mg/mL	Fatima et al. (2009)
	<i>C. krusei</i>	2 mg/mL	
	<i>C. nerformans</i>	0.5 mg/mL	
	<i>C. pseudotropicalis</i>	1 mg/mL	
<i>Heliotropium curassavicum</i>	<i>C. albicans</i>	100 mg/mL ^a	Mandeel and Taha (2005)
<i>Illicium verum</i>	<i>C. albicans</i>	16 mg/mL	Yazdani et al. (2009)
<i>Isatis indigotica</i>	<i>C. albicans</i>	Viability: 94.05%	Wong and Tsang (2009)
<i>Juniperus oxycedrus</i>	<i>C. albicans</i>	5 mg/mL	Ertürk (2006)
<i>Lantana fucata</i>	<i>C. krusei</i>	0.5 mg/mL	Correa-Royero et al. (2010)
<i>Laurus nobilis</i>	<i>C. albicans</i>	5 mg/mL	Ertürk (2006)
<i>Mahonia aquifolium</i>	<i>C. tropicalis</i>	1000 µg/mL	Volleková et al. (2003)
<i>Mallotus oppositifolius</i>	<i>C. albicans</i>	100 mg/mL ^a	Adekunle and Ikumapayi (2006)
<i>Melissa officinalis</i>	<i>C. albicans</i>	25 mg/mL	Ertürk (2006)
<i>Mentha piperita</i>	<i>C. albicans</i>	5 mg/mL	Ertürk (2006)
<i>Mentha piperita</i>	<i>C. albicans</i>	10 mg/mL ^a	Doddanna et al. (2013)
<i>Morinda morindoides</i>	<i>C. albicans</i>	31.25 mg/mL	Touré et al. (2011)
<i>Morinda royoc</i>	<i>C. krusei</i>	0.25 mg/mL	Correa-Royero et al. (2010)

Table 3 (Continued)

Plant source	Candida species	MIC	References
<i>Murraya koenigii</i>	<i>C. albicans</i>	10 mg/mL ^a	Doddanna et al. (2013)
<i>Myrcia cucullata</i>	<i>C. krusei</i>	0.03125 mg/mL	Correa-Royero et al. (2010)
<i>Odina woder</i>	<i>C. albicans</i>	500 mg/mL	Prabhakar et al. (2008)
	<i>C. glabrata</i>	500 mg/mL	
	<i>C. guilliermondii</i>	500 mg/mL	
	<i>C. krusei</i>	500 mg/mL	
	<i>C. parapsilosis</i>	500 mg/mL	
	<i>C. tropicalis</i>	500 mg/mL	
<i>Origanum vulgare</i>	<i>C. albicans</i>	5 mg/mL	Ličina et al. (2013)
<i>Paullinia cupana</i>	<i>C. albicans</i>	0.5 mg/mL	Basile et al. (2013)
<i>Pimpinella anisum</i>	<i>C. albicans</i>	19%, v/v	Kosalec et al. (2005)
	<i>C. krusei</i>	20%, v/v	
	<i>C. parapsilosis</i>	18%, v/v	
	<i>C. pseudotropicalis</i>	17%, v/v	
	<i>C. tropicalis</i>	18%, v/v	
<i>Piper nigrum</i>	<i>C. albicans</i>	12.5 mg/mL	Ertürk (2006)
		1.5 mg/mL ^a	Joe et al. (2009)
<i>Piper regnellii</i>	<i>C. albicans</i>	0.125 mg/mL	Pessini et al. (2005)
	<i>C. krusei</i>	0.5 mg/mL	
	<i>C. parapsilosis</i>	0.5 mg/mL	
<i>Plantago lanceolata</i> (France)	<i>C. albicans</i>	200 mg/mL	Hassawi and Kharma (2006)
<i>Plantago media</i>	<i>C. albicans</i>	200 mg/mL ^a	Qader et al. (2013)
	<i>C. dublicans</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
	<i>C. tropicalis</i>	200 mg/mL ^a	
<i>Pluchea ovalis</i>	<i>C. albicans</i>	100 mg/mL ^a	Mandeel and Taha (2005)
<i>Punica granatum</i>	<i>C. albicans</i>	0.25 mg/mL	Anibal et al. (2013)
	<i>C. dubliniensis</i>	0.25 mg/mL	
	<i>C. glabrata</i>	0.25 mg/mL	
	<i>C. guilliermondii</i>	0.125 mg/mL	
	<i>C. krusei</i>	0.125 mg/mL	
	<i>C. lusitaniae</i>	0.125 mg/mL	
	<i>C. parapsilosis</i>	0.125 mg/mL	
	<i>C. rugosa</i>	0.125 mg/mL	
	<i>C. tropicalis</i>	0.25 mg/mL	
	<i>C. utilis</i>	0.25 mg/mL	
<i>Punica granatum</i>	<i>C. albicans</i>	200 mg/mL ^a	Qader et al. (2013)
	<i>C. dublicans</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
	<i>C. tropicalis</i>	200 mg/mL ^a	
<i>Quercus infectoria</i>	<i>C. albicans</i>	200 mg/mL ^a	Qader et al. (2013)
	<i>C. dublicans</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
	<i>C. tropicalis</i>	200 mg/mL ^a	
<i>Rhus angustifolia</i>	<i>C. albicans</i>	200 mg/mL ^a	Qader et al. (2013)
	<i>C. dublicans</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
	<i>C. tropicalis</i>	200 mg/mL ^a	
<i>Rhus coriaria</i>	<i>C. albicans</i>	15 mg/mL	Ertürk (2006)
<i>Sargassum wightii</i>	<i>C. albicans</i>	100 mg/mL	Prabhakar et al. (2008)
	<i>C. glabrata</i>	100 mg/mL	
	<i>C. guilliermondii</i>	100 mg/mL	
	<i>C. krusei</i>	100 mg/mL	
	<i>C. parapsilosis</i>	100 mg/mL	
	<i>C. tropicalis</i>	100 mg/mL	
<i>Sclerocarya birrea</i>	<i>C. parapsilosis</i>	0.21 mg/mL	Masoko et al. (2008)
<i>Solanum nigrum</i>	<i>C. albicans</i>	0.2 mg/mL	Shamim et al. (2004)
	<i>C. glabrata</i>	1 mg/mL	
	<i>C. tropicalis</i>	0.2 mg/mL	
<i>Syzygium jambolanum</i>	<i>C. albicans</i>	100 mg/mL	Prabhakar et al. (2008)
	<i>C. glabrata</i>	100 mg/mL	
	<i>C. guilliermondii</i>	100 mg/mL	
	<i>C. krusei</i>	100 mg/mL	
	<i>C. parapsilosis</i>	100 mg/mL	
	<i>C. tropicalis</i>	100 mg/mL	
<i>Tamarix arabica</i>	<i>C. albicans</i>	100 mg/mL ^a	(Mandeel and Taha, 2005)
<i>Thymus kotschyana</i>	<i>C. albicans</i>	200 mg/mL ^a	(Qader et al., 2013)
	<i>C. dublicans</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
	<i>C. tropicalis</i>	200 mg/mL ^a	
<i>Zingiber officinale</i>	<i>C. albicans</i>	1 mg/mL ^a	(Joe et al., 2009)

Table 3 (Continued)

Plant source	Candida species	MIC	References
<i>Zingiber officinalis</i>	<i>C. albicans</i>	200 mg/mL ^a	(Qader et al., 2013)
	<i>C. dublicans</i>	200 mg/mL ^a	
	<i>C. glabrata</i>	200 mg/mL ^a	
	<i>C. krusei</i>	200 mg/mL ^a	
	<i>C. tropicalis</i>	200 mg/mL ^a	

MIC – minimum inhibitory concentration.

^a Concentration used to evaluate antifungal activity by disc diffusion.**Table 4**Activity of methanol–dichloromethane extracts against *Candida* species.

Plant source	Species	MIC	References
<i>Aframomum citratum</i>	<i>C. albicans</i>	3.12 mg/mL	Dzoyem et al. (2014)
	<i>C. glabrata</i>	4.68 mg/mL	
	<i>C. guilliermondii</i>	3.12 mg/mL	
	<i>C. krusei</i>	0.39 mg/mL	
	<i>C. lusitaniae</i>	0.78 mg/mL	
	<i>C. parapsilosis</i>	6.25 mg/mL	
<i>Aframomum melegueta</i>	<i>C. tropicalis</i>	6.25 mg/mL	Dzoyem et al. (2014)
	<i>C. albicans</i>	6.25 mg/mL	
	<i>C. guilliermondii</i>	1.56 mg/mL	
	<i>C. krusei</i>	6.25 mg/mL	
	<i>C. lusitaniae</i>	3.12 mg/mL	
	<i>C. parapsilosis</i>	6.25 mg/mL	
<i>Andrographis paniculata</i> ^a	<i>C. tropicalis</i>	1.56 mg/mL	Sule et al. (2012)
	<i>C. albicans</i>	0.1 mg/mL	
	<i>C. krusei</i>	0.25 mg/mL	
	<i>C. tropicalis</i>	0.1 mg/mL	
<i>Bielschmiedia cinnamomea</i>	<i>C. albicans</i>	3.12 mg/mL	Dzoyem et al. (2014)
	<i>C. glabrata</i>	6.25 mg/mL	
	<i>C. guilliermondii</i>	6.25 mg/mL	
	<i>C. lusitaniae</i>	3.12 mg/mL	
	<i>C. tropicalis</i>	3.12 mg/mL	
<i>Cinnamomum zeylanicum</i>	<i>C. albicans</i>	3.12 mg/mL	Dzoyem et al. (2014)
	<i>C. glabrata</i>	3.12 mg/mL	
	<i>C. guilliermondii</i>	0.78 mg/mL	
	<i>C. krusei</i>	0.78 mg/mL	
	<i>C. lusitaniae</i>	0.78 mg/mL	
	<i>C. parapsilosis</i>	3.12 mg/mL	
<i>Dichrostachys glomerata</i>	<i>C. tropicalis</i>	0.097 mg/mL	Dzoyem et al. (2014)
	<i>C. albicans</i>	0.39 mg/mL	
	<i>C. glabrata</i>	3.12 mg/mL	
	<i>C. guilliermondii</i>	3.12 mg/mL	
	<i>C. krusei</i>	3.12 mg/mL	
	<i>C. lusitaniae</i>	0.39 mg/mL	
<i>Diospyros crassiflora</i>	<i>C. parapsilosis</i>	6.25 mg/mL	Dzoyem et al. (2007)
	<i>C. tropicalis</i>	3.12 mg/mL	
	<i>C. albicans</i>	12.5 mg/mL	
	<i>C. glabrata</i>	25 mg/mL	
	<i>C. krusei</i>	12.5 mg/mL	
	<i>C. tropicalis</i>	12.5 mg/mL	
<i>Dorstenia psilurus</i>	<i>C. albicans</i>	0.39 mg/mL	Dzoyem et al. (2014)
	<i>C. glabrata</i>	3.12 mg/mL	
	<i>C. guilliermondii</i>	1.56 mg/mL	
	<i>C. krusei</i>	3.12 mg/mL	
	<i>C. lusitaniae</i>	3.12 mg/mL	
	<i>C. parapsilosis</i>	3.12 mg/mL	
<i>Echinops giganteus</i>	<i>C. tropicalis</i>	0.39 mg/mL	Dzoyem et al. (2014)
	<i>C. albicans</i>	6.25 mg/mL	
	<i>C. glabrata</i>	1.56 mg/mL	
	<i>C. guilliermondii</i>	3.12 mg/mL	
	<i>C. lusitaniae</i>	6.25 mg/mL	
	<i>C. parapsilosis</i>	6.25 mg/mL	
<i>Fagara lepreurii</i>	<i>C. tropicalis</i>	6.25 mg/mL	Dzoyem et al. (2014)
	<i>C. albicans</i>	3.12 mg/mL	
	<i>C. glabrata</i>	3.12 mg/mL	
	<i>C. krusei</i>	6.25 mg/mL	
	<i>C. parapsilosis</i>	6.25 mg/mL	
	<i>C. tropicalis</i>	6.25 mg/mL	
<i>Fagara macrophylla</i>	<i>C. albicans</i>	3.12 mg/mL	Dzoyem et al. (2014)
	<i>C. glabrata</i>	3.12 mg/mL	
	<i>C. guilliermondii</i>	1.56 mg/mL	
	<i>C. krusei</i>	6.25 mg/mL	
	<i>C. lusitaniae</i>	1.56 mg/mL	
	<i>C. tropicalis</i>	0.39 mg/mL	
<i>Gonzalagunia rosea</i> ^a	<i>C. albicans</i>	2.50 mg/mL	Niño et al. (2007)

Table 4 (Continued)

Plant source	Species	MIC	References
<i>Imperata cylindrical</i>	<i>C. albicans</i>	6.25 mg/mL	Dzoyem et al. (2014)
	<i>C. glabrata</i>	3.12 mg/mL	
	<i>C. guilliermondii</i>	3.12 mg/mL	
	<i>C. krusei</i>	1.56 mg/mL	
	<i>C. lusitaniae</i>	3.12 mg/mL	
	<i>C. parapsilosis</i>	6.25 mg/mL	
	<i>C. tropicalis</i>	3.12 mg/mL	
<i>Liabum asclepiadeum</i> ^a	<i>C. albicans</i>	0.62 mg/mL	Niño et al. (2007)
<i>Mondia whitei</i>	<i>C. albicans</i>	3.12 mg/mL	Dzoyem et al. (2014)
	<i>C. guilliermondii</i>	6.25 mg/mL	
	<i>C. lusitaniae</i>	3.12 mg/mL	
<i>Monodora myristica</i>	<i>C. tropicalis</i>	3.12 mg/mL	Dzoyem et al. (2014)
	<i>C. albicans</i>	6.25 mg/mL	
	<i>C. guilliermondii</i>	3.12 mg/mL	
	<i>C. krusei</i>	6.25 mg/mL	
	<i>C. lusitaniae</i>	3.12 mg/mL	
<i>Montanoa</i> ^a	<i>C. tropicalis</i>	3.12 mg/mL	Dzoyem et al. (2014)
	<i>C. albicans</i>	1.25 mg/mL	
	<i>C. albicans</i>	0.097 mg/mL	
	<i>C. glabrata</i>	1.56 mg/mL	
	<i>C. guilliermondii</i>	0.78 mg/mL	
<i>Olax subscorpioidea</i>	<i>C. krusei</i>	1.56 mg/mL	Dzoyem et al. (2014)
	<i>C. lusitaniae</i>	0.19 mg/mL	
	<i>C. parapsilosis</i>	0.39 mg/mL	
	<i>C. tropicalis</i>	0.048 mg/mL	
	<i>C. albicans</i>	3.12 mg/mL	
	<i>C. glabrata</i>	6.25 mg/mL	
	<i>C. guilliermondii</i>	6.25 mg/mL	
<i>Pentadiplandra brazzeana</i>	<i>C. lusitaniae</i>	6.25 mg/mL	Dzoyem et al. (2014)
	<i>C. tropicalis</i>	1.56 mg/mL	
	<i>C. albicans</i>	3.12 mg/mL	
	<i>C. guilliermondii</i>	6.25 mg/mL	
	<i>C. krusei</i>	6.25 mg/mL	
<i>Piper capense</i>	<i>C. tropicalis</i>	3.12 mg/mL	Dzoyem et al. (2014)
	<i>C. albicans</i>	3.12 mg/mL	
	<i>C. guilliermondii</i>	3.12 mg/mL	
	<i>C. krusei</i>	3.12 mg/mL	
	<i>C. lusitaniae</i>	1.56 mg/mL	
	<i>C. parapsilosis</i>	3.12 mg/mL	
	<i>C. tropicalis</i>	0.19 mg/mL	
<i>Piper guineense</i>	<i>C. albicans</i>	3.12 mg/mL	Dzoyem et al. (2014)
	<i>C. glabrata</i>	3.12 mg/mL	
	<i>C. lusitaniae</i>	1.56 mg/mL	
	<i>C. parapsilosis</i>	6.25 mg/mL	
	<i>C. tropicalis</i>	3.12 mg/mL	
	<i>C. albicans</i>	2.5 mg/mL	
	<i>C. albicans</i>	2.34 mg/mL	
<i>Schizogygia caffaeoides</i> ^a	<i>C. glabrata</i>	4.68 mg/mL	Dzoyem et al. (2014)
	<i>C. guilliermondii</i>	6.25 mg/mL	
	<i>C. krusei</i>	3.90 mg/mL	
	<i>C. lusitaniae</i>	2.34 mg/mL	
	<i>C. parapsilosis</i>	6.25 mg/mL	
	<i>C. tropicalis</i>	4.68 mg/mL	
	<i>C. guilliermondii</i>	6.25 mg/mL	
<i>Scorodophloeus zenkeri</i>	<i>C. tropicalis</i>	3.12 mg/mL	Dzoyem et al. (2014)
	<i>C. albicans</i>	3.12 mg/mL	
<i>Solanum melongena</i>	<i>C. glabrata</i>	6.25 mg/mL	Dzoyem et al. (2014)
	<i>C. guilliermondii</i>	3.12 mg/mL	
	<i>C. tropicalis</i>	3.12 mg/mL	
<i>Tetrapleura tetraptera</i>	<i>C. albicans</i>	3.12 mg/mL	Dzoyem et al. (2014)
	<i>C. glabrata</i>	6.25 mg/mL	
	<i>C. guilliermondii</i>	3.12 mg/mL	
	<i>C. krusei</i>	6.25 mg/mL	
	<i>C. lusitaniae</i>	3.12 mg/mL	
	<i>C. parapsilosis</i>	3.12 mg/mL	
	<i>C. tropicalis</i>	6.25 mg/mL	
<i>Xylopi aethiopica</i>	<i>C. albicans</i>	3.12 mg/mL	Dzoyem et al. (2014)
	<i>C. krusei</i>	3.12 mg/mL	
	<i>C. parapsilosis</i>	3.12 mg/mL	
	<i>C. tropicalis</i>	3.12 mg/mL	
<i>Xylopi parviflora</i>	<i>C. albicans</i>	6.25 mg/mL	Dzoyem et al. (2014)
	<i>C. glabrata</i>	6.25 mg/mL	
	<i>C. guilliermondii</i>	3.12 mg/mL	
	<i>C. krusei</i>	3.12 mg/mL	
	<i>C. lusitaniae</i>	1.56 mg/mL	
	<i>C. parapsilosis</i>	3.12 mg/mL	
	<i>C. tropicalis</i>	3.12 mg/mL	

MIC – minimum inhibitory concentration.

^a Dichloromethane extracts.

It can be concluded that the type of solvent used in the extraction of the phytochemicals (including phenolic compounds) has influence in the final composition of the extract, and consequently,

in the bioactivity. Nevertheless, the *Candida* species, as also the strains used had also influence in the antifungal results. Several studies have highlighted some aspects involved in the mecha-

Table 5
Activity of methanolic extracts against *Candida* species.

Plant sources	<i>Candida</i> species	MIC	References
<i>Adiantum capillus veneris</i>	<i>C. albicans</i>	1 mg/mL ^a	Ishaq et al. (2014)
<i>Anagyris foetida</i>	<i>C. albicans</i>	12.5 mg/mL	Darwish and Aburjai (2011)
	<i>C. glabrata</i>	25 mg/mL	
	<i>C. krusei</i>	12.5 mg/mL	
<i>Andrographis paniculata</i>	<i>C. albicans</i>	0.2 mg/mL	Sule et al. (2012)
	<i>C. krusei</i>	0.2 mg/mL	
	<i>C. tropicalis</i>	0.150 mg/mL	
<i>Arrabidaea chica</i>	<i>C. albicans</i>	0.5 mg/mL	Höfling et al. (2011)
<i>Artemisia herba-alba</i>	<i>C. albicans</i>	6.3 mg/mL	Darwish and Aburjai (2011)
	<i>C. glabrata</i>	25 mg/mL	
	<i>C. krusei</i>	12.5 mg/mL	
<i>Caesalpinia pulcherrima</i>	<i>C. albicans</i>	0.125 mg/disc ^a	Parekh and Chanda (2008)
<i>Capparis spinosa</i>	<i>C. albicans</i>	12.5 mg/mL	Darwish and Aburjai (2011)
	<i>C. glabrata</i>	12.5 mg/mL	
	<i>C. krusei</i>	12.5 mg/mL	
<i>Cassia fistula</i>	<i>C. albicans</i>	0.150 mg/mL	Irshad et al. (2011)
	<i>C. glabrata</i>	0.1 mg/mL	
	<i>C. tropicalis</i>	0.250 mg/mL	
<i>Centaurium erythraea</i>	<i>C. albicans</i>	0.1 mg/mL	Siler et al. (2014)
<i>Centaurium littorale</i> spp. <i>uliginosum</i>	<i>C. albicans</i>	0.2 mg/mL	Siler et al. (2014)
<i>Centaurium pulchellum</i>	<i>C. albicans</i>	0.4 mg/mL	Siler et al. (2014)
<i>Centaurium spicatum</i>	<i>C. albicans</i>	0.2 mg/mL	Siler et al. (2014)
<i>Centaurium tenuiflorum</i>	<i>C. albicans</i>	0.2 mg/mL	Siler et al. (2014)
<i>Crossandra infundibuliformis</i>	<i>C. krusei</i>	125 mg/mL	Madhumitha and Saral (2011)
<i>Diospyros canaliculata</i>	<i>C. albicans</i>	0.0125 mg/mL	Dzoyem et al. (2011)
	<i>C. kefyr</i>	0.025 mg/mL	
	<i>C. parapsilosis</i>	0.0125 mg/mL	
<i>Dorstenia turbinata</i>	<i>C. albicans</i>	0.039 mg/mL	Ngameni et al. (2009)
	<i>C. glabrata</i>	0.039 mg/mL	
<i>Ephedra pachyclada</i>	<i>C. albicans</i>	0.0005 mg/mL ^a	Parsaeimehr et al. (2010)
<i>Ephedra procera</i>	<i>C. albicans</i>	0.0005 mg/mL ^a	Parsaeimehr et al. (2010)
<i>Ephedra strobilacea</i>	<i>C. albicans</i>	0.001 mg/mL ^a	Parsaeimehr et al. (2010)
<i>Equinops polyceras</i>	<i>C. albicans</i>	6.3 mg/mL	Darwish and Aburjai (2011)
	<i>C. glabrata</i>	12.5 mg/mL	
	<i>C. krusei</i>	3.1 mg/mL	
<i>Eucalyptus intertexta</i>	<i>C. albicans</i>	0.0625 mg/mL	Safaei-Ghomi and And (2010)
<i>Eucalyptus largiflorens</i>	<i>C. albicans</i>	0.0312 mg/mL	Safaei-Ghomi and Ahd (2010)
<i>Eucalyptus sideroxylon</i>	<i>C. albicans</i>	400 mg/mL [*]	Ashour (2008)
<i>Eucalyptus torquata</i>	<i>C. albicans</i>	400 mg/mL [*]	Ashour (2008)
<i>Euphorbia hirta</i>	<i>C. albicans</i>	3.13 mg/mL	Rajeh et al. (2010)
<i>Euphorbia macroclada</i>	<i>C. albicans</i>	12.5 mg/mL	Darwish and Aburjai (2011)
		25 mg/mL	
	<i>C. glabrata</i>	12.5 mg/mL	
		25 mg/mL	
	<i>C. krusei</i>	12.5 mg/mL	
		25 mg/mL	
<i>Euphorbia tirucalli</i>	<i>C. albicans</i>	0.125 mg/disc ^a	Parekh and Chanda (2008)
<i>Glycyrrhiza glabra</i>	<i>C. albicans</i>	0.0005 mg/mL ^a	Karomi et al. (2012)
<i>Gonzalagunia rosea</i>	<i>C. albicans</i>	1.25 mg/mL	Niño et al. (2007)
<i>Holarrhena antidysenterica</i>	<i>C. tropicalis</i>	0.125 mg/disc ^a	Parekh and Chanda (2008)
<i>Hybiscus sabdariffa</i>	<i>C. krusei</i>	25 mg/mL	Darwish and Aburjai (2011)
<i>Hypericum</i>	<i>C. albicans</i>	12.5 mg/mL	Darwish and Aburjai (2011)
<i>trequetriofolium</i>	<i>C. glabrata</i>	25 mg/mL	
	<i>C. krusei</i>	12.5 mg/mL	
<i>Hyssopus officinalis</i> subsp. <i>pilifer</i>	<i>C. albicans</i>	10 mg/mL	Džamić et al. (2013)
<i>Laurus nobilis</i>	<i>C. albicans</i>	0.13 mg/mL	Unver et al. (2008)
	<i>C. glabrata</i>	0.13 mg/mL	
	<i>C. holmii</i>	0.16 mg/mL	
	<i>C. insane</i>	0.07 mg/mL	
	<i>C. krusei</i>	0.07 mg/mL	
	<i>C. tropicalis</i>	0.04 mg/mL	
<i>Leiothrix spiralis</i>	<i>C. albicans</i>	0.5 mg/mL	Araújo et al. (2012)
	<i>C. krusei</i>	0.5 mg/mL	
	<i>C. parapsilosis</i>	0.25 mg/mL	
	<i>C. tropicalis</i>	1 mg/mL	
<i>Lepidium sativum</i>	<i>C. albicans</i>	25 mg/mL	Darwish and Aburjai (2011)
<i>Mangifera indica</i>	<i>C. albicans</i>	0.125 mg/disc ^a	Parekh and Chanda (2008)
<i>Mentha longifolia</i>	<i>C. albicans</i>	6.3 mg/mL	Darwish and Aburjai (2011)
	<i>C. glabrata</i>	12.5 mg/mL	
	<i>C. krusei</i>	12.5 mg/mL	
<i>Mesua ferra</i>	<i>C. albicans</i>	0.125 mg/disc ^a	Parekh and Chanda (2008)
<i>Mindium laevigatum</i>	<i>C. albicans</i>	0.1 mg/mL	Modaressi et al. (2013)
<i>Montanoa</i>	<i>C. albicans</i>	0.62 mg/mL	Niño et al. (2007)
<i>Munnozia senecionidis</i>	<i>C. albicans</i>	2.5 mg/mL	Niño et al. (2007)

Table 5 (Continued)

Plant sources	<i>Candida</i> species	MIC	References
<i>Origanum syriacum</i>	<i>C. albicans</i>	6.3 mg/mL	Darwish and Aburjai (2011)
	<i>C. glabrata</i>	12.5 mg/mL	
	<i>C. krusei</i>	12.5 mg/mL	
<i>Paeonia rockii</i>	<i>C. albicans</i>	0.025 mg/mL	Picerno et al. (2011)
<i>Phlomis brachydon</i>	<i>C. albicans</i>	6.3 mg/mL	Darwish and Aburjai (2011)
	<i>C. glabrata</i>	12.5 mg/mL	
	<i>C. krusei</i>	3.1 mg/mL	
<i>Pimpinella anisum</i>	<i>C. albicans</i>	12.5 mg/mL	Darwish and Aburjai (2011)
	<i>C. glabrata</i>	12.5 mg/mL	
	<i>C. krusei</i>	25 mg/mL	
<i>Pimpinella anisum</i>	<i>C. albicans</i>	16 mg/mL	Yazdani et al. (2009)
<i>Psidium sartorianum</i>	<i>C. albicans</i>	16 mg/mL	Camacho-Hernández et al. (2004)
<i>Ribes x nidigrolaria</i>	<i>C. glabrata</i>	2.91 mg/mL	Krisch et al. (2008)
	<i>C. inconspicua</i>	3.33 mg/mL	
	<i>C. lipolytica</i>	4.67 mg/mL	
	<i>C. norvegica</i>	10.98 mg/mL	
	<i>C. parapsilosis</i>	3.56 mg/mL	
	<i>C. tropicalis</i>	4.23 mg/mL	
	<i>C. zeylanoides</i>	5.68 mg/mL	
	<i>C. guilliermondii</i>	6.13 mg/mL	
	<i>C. inconspicua</i>	4.22 mg/mL	
<i>Ribes nigrum</i>	<i>C. parapsilosis</i>	4.41 mg/mL	Krisch et al. (2008)
	<i>C. tropicalis</i>	7.16 mg/mL	
	<i>C. glabrata</i>	4.60 mg/mL	
	<i>C. lipolytica</i>	4.63 mg/mL	
<i>Ribes uva-crispa</i>	<i>C. glabrata</i>	4.60 mg/mL	Krisch et al. (2008)
	<i>C. lipolytica</i>	4.63 mg/mL	
<i>Salvia officinalis</i>	<i>C. clus</i>	0.16 mg/mL	Unver et al. (2008)
	<i>C. tropicalis</i>	0.16 mg/mL	
<i>Saussurea lappa</i>	<i>C. albicans</i>	0.125 mg/disc ^a	Parekh and Chanda (2008)
<i>Schistocarpha sinforosi</i>	<i>C. albicans</i>	2.5 mg/mL	Niño et al. (2007)
<i>Schizozygia coffaeoides</i>	<i>C. albicans</i>	10 mg/mL	Kariba et al. (2001)
<i>Sclerocarya birrea</i>	<i>C. parapsilosis</i>	0.32 mg/mL	Masoko et al. (2008)
<i>Thea sinensis</i>	<i>C. albicans</i>	6.3 mg/mL	Darwish and Aburjai (2011)
	<i>C. glabrata</i>	12.5 mg/mL	
	<i>C. krusei</i>	12.5 mg/mL	
<i>Thecacoris annobonae</i>	<i>C. albicans</i>	0.3125 mg/mL	Kuete et al. (2010)
<i>Thymbra spicata</i>	<i>C. albicans</i>	0.04 mg/mL	Unver et al. (2008)
	<i>C. clus</i>	0.01 mg/mL	
	<i>C. glabrata</i>	0.01 mg/mL	
	<i>C. krusei</i>	0.01 mg/mL	
	<i>C. parapsilosis</i>	0.01 mg/mL	
	<i>C. tropicalis</i>	0.01 mg/mL	
	<i>C. tropicalis</i>	0.125 mg/disc ^a	
	<i>C. albicans</i>	12.5 mg/mL	
	<i>C. krusei</i>	12.5 mg/mL	
<i>Trapa natans</i>	<i>C. tropicalis</i>	0.125 mg/disc ^a	Parekh and Chanda (2008)
<i>Trigonella</i>	<i>C. albicans</i>	12.5 mg/mL	Darwish and Aburjai (2011)
<i>foenum-graecum</i>	<i>C. krusei</i>	12.5 mg/mL	(2011)
<i>Varthemia iphionoides</i>	<i>C. albicans</i>	6.3 mg/mL	Darwish and Aburjai (2011)
<i>Verbesina nudipes</i>	<i>C. albicans</i>	2.5 mg/mL	Niño et al. (2007)
<i>Vitex negundo</i>	<i>C. albicans</i>	0.125 mg/disc ^a	Parekh and Chanda (2008)

MIC – minimum inhibitory concentration.

^a Concentration used to evaluate antifungal activity by disc diffusion.

nisms of action such as interaction with the lipid bilayer of the cell membrane, which consequently affect the membrane permeability, respiratory chain and cellular energy production (Tintino et al., 2014), as well as the inhibition of the H⁺-ATPase (Kuete et al., 2010). Beyond that it is possible to distinguish the antifungal potential, between fungistatic and/or fungicidal. The majority of the studies reported fungistatic effects of plant extracts (Anibal et al., 2013; Barros et al., 2013a; Basile et al., 2013; Correa-Royero et al., 2010; Dzoyem et al., 2014; Kariba et al., 2001; Kim and Lee, 2012; Kurdelas et al., 2010; Lubian et al., 2010; Nordin et al., 2013; Pathak, 2012; Thirach et al., 2003; Tomczykowa et al., 2008; Tsuzuki et al., 2007), but there are also some reports evidencing fungicidal effects, most of them in a dose-dependent manner (Askun et al., 2008; Barros et al., 2013a; Dzoyem et al., 2014; Korukluoglu et al., 2008; Lubian et al., 2010; Nordin et al., 2013; Rajeh et al., 2010; Siler et al., 2014; Sule et al., 2012; Yazdani et al., 2009).

It is crucial to characterize the phytochemical composition of the plant extracts with antifungal activity, in order to identify the individual compounds responsible for this bioactivity, and to elucidate the main molecular targets in *Candida* species. Some studies have been performed in individual phenolic compounds (one of the

most abundant molecules in polar extracts), as it will be discussed in the next section but, unfortunately, the majority of the studies report anti-*Candida* activity of crude extracts without any chemical characterization.

4.2. The use of phenolic compounds against *Candida* species

Phenolic acids, flavonoids and tannins are the most studied groups of phenolic compounds, comprising molecules with evidenced antifungal properties. These molecules have been isolated from different plant sources, as also commercially obtained as pure compounds.

A commercial sample of gallic acid presented higher antifungal potential against *C. albicans* than the same compound isolated from *Lythrum salicaria*, *Paeonia rockii* or *Pelargonium reniforme* subsp. *reniforme* (Table 8). The same behaviour was observed for commercial quercetin, in comparison to the one obtained from *Buddleja salviifolia* and *Halimodendron halodendron* (Table 9). On the other hand, commercial sample of salicylic acid presented lower antifungal potential than the same compound isolated from *Halimodendron halodendron* (Table 8). It would be expected a similar

Table 6
Activity of hydroalcoholic extracts against *Candida* species.

	Plant source	<i>Candida</i> species	MIC	References
Ethanol: water (90:10 v/v)	<i>Sapindus saponaria</i>	<i>C. parapsilosis</i>	0.4 mg/mL	Tsuzuki et al. (2007)
Ethanol: water (70:30 v/v)	<i>Embllica officinalis</i>	<i>C. albicans</i>	0.1024 mg/mL	Tharkar et al. (2010)
Ethanol: water (70:30 v/v)	<i>Glycyrrhiza glabra</i>	<i>C. albicans</i>	0.512 mg/mL	Tharkar et al. (2010)
Ethanol: water (70:30 v/v)	<i>Momordica charantia</i>	<i>C. guilhermondii</i>	0.1 mg/mL ^a	Silva (2012)
Ethanol: water (70:30 v/v)	<i>Punica granatum</i>	<i>C. albicans</i>	0.1 mg/mL ^a	Silva (2012)
		<i>C. guilhermondii</i>	0.1 mg/mL ^a	
		<i>C. krusei</i>	0.1 mg/mL ^a	
		<i>C. parapsilosis</i>	0.1 mg/mL ^a	
		<i>C. tropicalis</i>	0.1 mg/mL ^a	
Ethanol: water (70:30 v/v)	<i>Syzygium aromaticum</i>	<i>C. guilhermondii</i>	0.05 mg/mL	Silva (2012)
		<i>C. parapsilosis</i>	0.1 mg/mL ^a	
		<i>C. tropicalis</i>	0.1 mg/mL ^a	
Ethanol: water (50:50 v/v)	<i>Achyranthes aspera</i>	<i>C. albicans</i>	0.3 mg/mL	Mathur et al. (2011)
Ethanol: water (50:50 v/v)	<i>Punica granatum</i>	<i>C. albicans</i>	0.0039 mg/mL	Endo et al. (2010)
		<i>C. parapsilosis</i>	0.0039 mg/mL	
Methanol: water (90:10 v/v)	<i>Cassia fistula</i>	<i>C. albicans</i>	0.025 mg/mL	Bhalodia et al. (2012)
Methanol: water (80:20 v/v)	<i>Castanea sativa</i>	<i>C. albicans</i>	0.625 mg/mL	Barros et al. (2013a)
		<i>C. tropicalis</i>	0.625 mg/mL	
Methanol: water (80:20 v/v)	<i>Cistus lanadifer</i>	<i>C. tropicalis</i>	0.625 mg/mL	Barros et al. (2013b)
Methanol: water (80:20 v/v)	<i>Filipendula ulmaria</i>	<i>C. albicans</i>	0.625 mg/mL	Barros et al. (2013a)
		<i>C. tropicalis</i>	0.155 mg/mL	
Methanol: water (80:20 v/v)	<i>Rosa micrantha</i>	<i>C. albicans</i>	0.05 mg/mL	Barros et al. (2013a)
		<i>C. tropicalis</i>	0.155 mg/mL	
Methanol: water (80:20 v/v)	<i>Salvia officinalis</i>	<i>C. parapsilosis</i>	2.5 mg/mL	Martins et al. (2015)
		<i>C. tropicalis</i>	1.25 mg/mL	

MIC – minimum inhibitory concentration.

^a Concentration used to evaluate antifungal activity by disc diffusion.

bioactivity for the same compound, independently of its source. Therefore, the observed differences are probably related with the purity grade of the tested compounds, as well as with the vehicles/solvents used to test antifungal activities.

Table 8 shows the activity of phenolic acids (hydroxycinnamic acid derivatives, hydroxybenzoic acid derivatives and other related compounds) against *Candida* species. In general, hydroxycinnamic acid derivatives presented lower MICs against *Candida* species, than the hydroxybenzoic acid derivatives. The only exception was reported by Faria et al. (2011); however, in this case, it is necessary to take into account that the results referred to tested concentrations and not MIC values. Analyzing the results by each *Candida* species, in general, commercial compounds, mainly albicanil caffeate (MIC=4 µg/mL) presented the most pronounced antifungal effects against *C. albicans*. Two phenolic acid related compounds, dihydro-*N*-caffeoyltyramine and *trans-N*-caffeoyltyramine (MICs = 5 µg/mL) obtained from *Lycium chinense* also evidenced a significant potential, followed by gallic acid

isolated from *Paeonia rockii* (30 µg/mL) and gentisic, syringic, vanillic and 4-*O*-β-D-(6-*O*-gentisoylglucosyl) vanillic acids from *Stenoloma chusanum* (50 µg/mL). Regarding other *Candida* species, only Latte and Kolodziej (2000) report the antifungal activity of gallic acid from *Pelargonium reniforme* subsp. *reniforme* against *C. glabrata* (MIC = 62 µg/mL) and *C. krusei* (MIC = 125 µg/mL). To authors' knowledge there are only reports regarding commercial compounds against *C. parapsilosis*, *C. tropicalis* and *C. lusitanae*. The antifungal activity of flavonoids against *Candida* species is presented in Table 9. Commercial compounds displayed slight higher antifungal effects than flavonoids isolated from plant sources. Among the commercial compounds, silibinin (MIC = 4 µg/mL), a flavolignan, exerted the most pronounced antifungal activity, against *C. albicans*, followed by apigenin (a flavones), genistein (an isoflavone), naringin (a flavanone), quercetin (a flavonol) and silymarin (a flavolignan), which presented similar potential (MIC = 8 µg/mL). Concerning to the flavonoids derived from plant origin, 2'-*O*-methylabronisoflavone (16–25 µg/mL) and 9-

Table 7
Activity of other extracts against *Candida* species.

Extract	Plant source	<i>Candida</i> species	MIC	References
Dimethyl sulfoxide (DMSO)	<i>Eleutherine plicata</i>	<i>C. albicans</i>	259 mg/mL	Menezes et al. (2009)
	<i>Psidium guajava</i>		125 mg/mL	
	<i>Syzygium aromaticum</i>		62.5 mg/mL	
Diethyl ether	<i>Origanum vulgare</i>	<i>C. albicans</i>	5 mg/mL	Ličina et al. (2013)
Ethyl acetate	<i>Crossandra infundibuliformis</i>	<i>C. krusei</i>	125 mg/mL	Madhumitha and Saral (2011)
	<i>Curcuma zedoaria</i>	<i>C. albicans</i>	12.5 mg/mL	Zhu et al. (2005)
		<i>C. lusitanae</i>	12.5 mg/mL	
	<i>Hyssopus officinalis</i> subsp. <i>Pilifer</i>	<i>C. albicans</i>	7.00 mg/mL	Džamić et al. (2013)
	<i>Origanum vulgare</i>	<i>C. albicans</i>	5 mg/mL	Ličina et al. (2013)
Tinctures	<i>Anacardium occidentale</i>	<i>C. albicans</i>	100 mg/mL	Cardoso et al. (2012)
		<i>C. krusei</i>	100 mg/mL	
	<i>Malva sylvestris</i>	<i>C. albicans</i>	100 mg/mL	Cardoso et al. (2012)
		<i>C. krusei</i>	25 mg/mL	
		<i>C. tropicalis</i>	100 mg/mL	
	<i>Salvia officinalis</i>	<i>C. krusei</i>	100 mg/mL	Cardoso et al. (2012)

MIC – minimum inhibitory concentration.

Table 8
Activity of phenolics acids against *Candida* species.

Phenolic acids	Source	<i>Candida</i> species	MIC	References
Hydroxycinnamic acid derivatives				
Caffeic acid	Commercial compound	<i>C. albicans</i> <i>C. parapsilosis</i>	8 µg/mL 16 µg/mL	Ozçelik et al. (2011)
Chlorogenic acid	Commercial compound	<i>C. albicans</i> <i>C. parapsilosis</i>	8 µg/mL 16 µg/mL	Ozçelik et al. (2011)
<i>m</i> -coumaric acid (2-hydroxycinnamic acid)	Commercial compound	<i>C. albicans</i> ; <i>C. parapsilosis</i> ; <i>C. glabrata</i> ; <i>C. tropicalis</i> ; <i>C. krusei</i> ; <i>C. lusitaniae</i>	820 µg/mL ^a	Faria et al. (2011)
<i>o</i> -coumaric acid (3-hydroxycinnamic acid)	Commercial compound	<i>C. albicans</i> ; <i>C. parapsilosis</i> ; <i>C. glabrata</i> ; <i>C. tropicalis</i> ; <i>C. krusei</i> ; <i>C. lusitaniae</i>	820 µg/mL ^a	Faria et al. (2011)
<i>p</i> -coumaric acid (4-hydroxycinnamic acid)	Commercial compound	<i>C. albicans</i> <i>C. albicans</i> ; <i>C. parapsilosis</i> ; <i>C. glabrata</i> ; <i>C. tropicalis</i> ; <i>C. krusei</i> ; <i>C. lusitaniae</i>	40 µg/mL 820 µg/mL ^a	Siler et al. (2014) Faria et al. (2011)
Ferulic acid	Commercial compound	<i>C. albicans</i>	20 µg/mL	Siler et al. (2014)
4-hydroxy-3-methoxy cinnamic acid	<i>Halimodendron halodendron</i>	<i>C. albicans</i>	200 µg/mL	Wang et al. (2012)
Hydroxybenzoic acid derivatives				
Gallic acid	<i>Lythrum salicaria</i> <i>Paeonia rockii</i> <i>Pelargonium reniforme</i> subsp. <i>reniforme</i>	<i>C. albicans</i> <i>C. albicans</i> <i>C. albicans</i> <i>C. glabrata</i> <i>C. krusei</i>	2500 µg/mL 30 µg/mL 500 µg/mL 62 µg/mL 125 µg/mL	Manayi et al. (2013) Picerno et al. (2011) Latte and Kolodziej (2000)
	Commercial compound	<i>C. albicans</i> <i>C. parapsilosis</i>	8 µg/mL 16 µg/mL	Ozçelik et al. (2011)
Gentisic acid	<i>Stenoloma chusanum</i>	<i>C. albicans</i>	50 µg/mL	Ren et al. (2009)
2- <i>O</i> -β-D-(6- <i>O</i> -gentisoylglucosyl) gentisic acid	<i>Stenoloma chusanum</i>	<i>C. albicans</i>	100 µg/mL	Ren et al. (2009)
<i>p</i> -Hydroxybenzoic acid	<i>Halimodendron halodendron</i>	<i>C. albicans</i>	100 µg/mL	Wang et al. (2012)
Salicylic acid	<i>Halimodendron halodendron</i> Commercial compound	<i>C. albicans</i> <i>C. albicans</i> ; <i>C. parapsilosis</i> ; <i>C. glabrata</i> ; <i>C. tropicalis</i> ; <i>C. krusei</i> ; <i>C. lusitaniae</i>	150 µg/mL 690 µg/mL ^a	Wang et al. (2012) Faria et al. (2011)
Syringic acid	<i>Stenoloma chusanum</i>	<i>C. albicans</i>	50 µg/mL	Ren et al. (2009)
Vanillic acid	<i>Stenoloma chusanum</i>	<i>C. albicans</i>	50 µg/mL	Ren et al. (2009)
4- <i>O</i> -β-D-(6- <i>O</i> -gentisoylglucosyl) vanillic acid	<i>Stenoloma chusanum</i>	<i>C. albicans</i>	50 µg/mL	Ren et al. (2009)
4-Acetoxy vanillic acid	<i>Thecatoris annobonae</i>	<i>C. albicans</i>	156.25 µg/mL	Kuete et al. (2010)
Other phenolic acids/related compounds				
Albicanyl caffeate	Commercial compound	<i>C. albicans</i>	4 µg/mL	Sun et al. (2004)
Dihydro- <i>N</i> -caffeoyltyramine ^b	<i>Lycium chinense</i>	<i>C. albicans</i>	5 µg/mL	Lee et al. (2004)
<i>cis</i> - <i>N</i> -caffeoyltyramine ^b	<i>Lycium chinense</i>	<i>C. albicans</i>	40 µg/mL	Lee et al. (2004)
<i>trans</i> - <i>N</i> -Feruloyloctopamine ^b	<i>Lycium chinense</i>	<i>C. albicans</i>	10 µg/mL	Lee et al. (2004)
<i>trans</i> - <i>N</i> -Caffeoyltyramine ^b	<i>Lycium chinense</i>	<i>C. albicans</i>	5 µg/mL	Lee et al. (2004)
3,3',4-tri- <i>O</i> -methyl ellagic acid-4- <i>O</i> -β-D-glucopyranoside	<i>Lythrum salicaria</i>	<i>C. albicans</i>	312 µg/mL	Manayi et al. (2013)
Quinic acid	Commercial compound	<i>C. albicans</i> <i>C. parapsilosis</i>	8 µg/mL 16 µg/mL	Ozçelik et al. (2011)

MIC – Minimum inhibitory concentration.

^a Tested concentration and not MIC.

^b Phenolic amides.

O-methyl-4-hydroxyboeravinone B (48 µg/mL), two isoflavones isolated from *Mirabilis jalapa*, also presented a pronounced antifungal potential against *C. albicans*, followed by the flavonols kaempferol, myricetin, quercetin and rutin from *Origanum vulgare* subsp. *gracile* and *Origanum acutidens* (except quercetin and rutin). In general, flavonols (mainly represented by kaempferol, myricetin and quercetin), followed by flavanones and flavolignans, were the most effective antifungal flavonoids against *Candida* species. Once again, *C. albicans* was the main studied *Candida* species, being only tested commercial flavonoids against *C. parapsilosis* and compounds isolated from *Origanum* species against *C. glabrata*.

Table 10 shows the antifungal activity of hydrolyzable tannins, stilbenes and xanthones against *Candida* species. Comparing the different hydrolyzable tannins isolated from *Pelargonium reniforme* subsp. *reniforme*, it is possible to conclude that similar antifungal effects were achieved; the obtained MIC values were the same

for corilagin, glucogallin, methyl ester, pelargonin and phyllanthusin, against *C. albicans*, *C. glabrata* and *C. krusei*. Comparing *Pelargonium reniforme* subsp. *reniforme* with *Punica granatum*, the hydrolyzable tannin, punicalagin, exerted higher antifungal effect against *C. albicans* (~128 times higher). Regarding stilbenes, resveratrol and 3,4'-difluorostilbene gave similar MIC values against *C. albicans* (Sun et al., 2004), while for xanthones the commercial compound mangiferin presented a more pronounced effect against *C. albicans*. 8-Carboxy-methyl-1,3,5,6-tetrahydroxyxanthone gave significantly different MIC values for each *Candida* species; *C. krusei* and *C. parapsilosis* were the most sensible to the mentioned compound (15.7 µg/mL), while *C. albicans* showed higher resistance (62.5 µg/mL). Among tannins, stilbenes and xanthones, the latter seem to be the most efficient, exhibiting lower MICs against *Candida* species. Overall, commercial compounds seem to be more efficient than phenolic compounds isolated from natu-

Table 9Activity of flavonoids against *Candida* species.

Flavonoids	Source	<i>Candida</i> species	MIC	References
Flavonols				
Kaempferol	<i>Origanum vulgare</i> subsp. <i>gracile</i> ; <i>Origanum acutidens</i>	<i>C. albicans</i> ; <i>C. glabrata</i>	0.1 µg/g ^a 0.5 µg/g ^a	Kursat et al. (2011)
Morin	<i>Origanum vulgare</i> subsp. <i>gracile</i> ; <i>Origanum acutidens</i>	<i>C. albicans</i> ; <i>C. glabrata</i>	42.8 µg/g ^a 4.5 µg/g ^a	Kursat et al. (2011)
Myricetin	<i>Origanum vulgare</i> subsp. <i>gracile</i> ; <i>Origanum acutidens</i>	<i>C. albicans</i> ; <i>C. glabrata</i>	0.2 µg/g ^a 1.2 µg/g ^a	Kursat et al. (2011)
Quercetin	<i>Buddleja salviifolia</i> <i>Origanum vulgare</i> subsp. <i>gracile</i> <i>Halimodendron halodendron</i> Commercial compound	<i>C. albicans</i> <i>C. albicans</i> ; <i>C. glabrata</i> <i>C. albicans</i> <i>C. albicans</i> ; <i>C. parapsilosis</i>	125 µg/mL 0.3 µg/g ^a 250 µg/mL 8 µg/mL	Pendota et al. (2013) Kursat et al. (2011) Wang et al. (2012) Ozcelik et al. (2011)
3- <i>O</i> -methylquercetin	<i>Halimodendron halodendron</i>	<i>C. albicans</i>	200 µg/mL	Wang et al. (2012)
3,3'-di- <i>O</i> -methylquercetin	<i>Halimodendron halodendron</i>	<i>C. albicans</i>	200 µg/mL	Wang et al. (2012)
8- <i>O</i> -methylretusin	<i>Halimodendron halodendron</i>	<i>C. albicans</i>	250 µg/mL	Wang et al. (2012)
8- <i>O</i> -methylretusin-7- <i>O</i> -β-D-glucopyranoside	<i>Halimodendron halodendron</i>	<i>C. albicans</i>	250 µg/mL	Wang et al. (2012)
Rutin	<i>Origanum vulgare</i> subsp. <i>gracile</i> Commercial compound	<i>C. albicans</i> ; <i>C. glabrata</i> <i>C. albicans</i>	6.4 µg/g ^a 40 µg/mL	Kursat et al. (2011) Siler et al. (2014)
Flavones				
Apigenin	Commercial compound	<i>C. albicans</i> ; <i>C. parapsilosis</i>	8 µg/mL 16 µg/mL	Ozcelik et al. (2011)
Luteolin-6- <i>C</i> -β-D-glucopyranoside	<i>Leiothrix spiralis</i>	<i>C. parapsilosis</i>	125 µg/mL	(Araújo et al., 2011)
4'-methoxyluteolin-6- <i>C</i> -β-D-glucopyranoside	<i>Leiothrix spiralis</i>	<i>C. albicans</i>	125 µg/mL	(Araújo et al., 2011)
3,5,7,8,4'-pentahydroxy-3'-methoxy flavone	<i>Halimodendron halodendron</i>	<i>C. albicans</i>	250 µg/mL	
Flavanones				
Naringin	Commercial compound	<i>C. albicans</i> <i>C. parapsilosis</i>	8 µg/mL 16 µg/mL	Ozcelik et al. (2011)
Naringenin	<i>Origanum acutidens</i>	<i>C. albicans</i> ; <i>C. glabrata</i>	10.3 µg/g ^a	Kursat et al. (2011)
Isoflavones				
Genistein	Commercial compound	<i>C. albicans</i> ; <i>C. parapsilosis</i>	8 µg/mL	Ozcelik et al. (2011)
2'- <i>O</i> -methylabronisoflavone	<i>Mirabilis jalapa</i>	<i>C. albicans</i>	16–25 µg/mL	Yang et al. (2001)
9- <i>O</i> -methyl-4-hydroxyboeravinone B	<i>Mirabilis jalapa</i>	<i>C. albicans</i>	48 µg/mL	Yang et al. (2001)
Flavolignan				
Silibinin	Commercial compound	<i>C. albicans</i> <i>C. parapsilosis</i>	4 µg/mL 8 µg/mL	Ozcelik et al. (2011)
Silymarin	Commercial compound	<i>C. albicans</i> <i>C. parapsilosis</i>	8 µg/mL 16 µg/mL	Ozcelik et al. (2011)

MIC – minimum inhibitory concentration.

^a Compound concentration in the tested extract.**Table 10**Activity of hydrolyzable tannins, stilbenes and xanthones against *Candida* species.

Hydrolyzable tannins	Source	<i>Candida</i> species	MIC	References
Corilagin	<i>Pelargonium reniforme</i> subsp. <i>reniforme</i>	<i>C. albicans</i> <i>C. glabrata</i> <i>C. krusei</i>	500 µg/mL 31 µg/mL 125 µg/mL	Latte and Kolodziej (2000)
Glucogallin	<i>Pelargonium reniforme</i> subsp. <i>reniforme</i>	<i>C. glabrata</i> <i>C. krusei</i>	31 µg/mL 125 µg/mL	Latte and Kolodziej (2000)
Methyl ester	<i>Pelargonium reniforme</i> subsp. <i>reniforme</i>	<i>C. glabrata</i> <i>C. krusei</i>	31 µg/mL 125 µg/mL	Latte and Kolodziej (2000)
Pelargoniin	<i>Pelargonium reniforme</i> subsp. <i>reniforme</i>	<i>C. albicans</i> <i>C. glabrata</i> <i>C. krusei</i>	500 µg/mL 31 µg/mL 125 µg/mL	Latte and Kolodziej (2000)
Phyllantusiin	<i>Pelargonium reniforme</i> subsp. <i>reniforme</i>	<i>C. albicans</i> <i>C. glabrata</i> <i>C. krusei</i>	500 µg/mL 31 µg/mL 125 µg/mL	Latte and Kolodziej (2000)
Punicalagin	<i>Punica granatum</i>	<i>C. albicans</i> <i>C. parapsilosis</i>	3.9 µg/mL 1.9 µg/mL	Endo et al. (2010)
Not specified	<i>Rhizophora apiculata</i>	<i>C. albicans</i>	6250 µg/mL	Lim et al. (2006)
Stilbenes				
Resveratrol	Commercial compound	<i>C. albicans</i> <i>C. albicans</i>	16 µg/mL 20 µg/mL	Sun et al. (2004) Jung et al. (2005)
3,4'-difluorostilbene	Commercial compound	<i>C. albicans</i>	16 µg/mL	Sun et al. (2004)
Xanthones				
8-carboxy-methyl-1,3,5,6-tetrahydroxyxanthone	<i>Leiothrix spiralis</i>	<i>C. albicans</i> <i>C. krusei</i> <i>C. parapsilosis</i> <i>C. tropicalis</i>	62.5 µg/mL 15.7 µg/mL 15.7 µg/mL 31.25 µg/mL	Araújo et al. (2011)
Mangiferin	Commercial compound	<i>C. albicans</i>	40 µg/mL	Siler et al. (2014)

MIC – minimum inhibitory concentration.

ral matrices probably due to their higher purity grade. Regarding phenolic compounds from natural sources, punicalagin (isolated from *Punica granatum*), followed by dihydro-*N*-caffeoyltyramine and *trans-N*-caffeoyltyramine (isolated from *Lycium chinense*) were the most efficient against *C. albicans*. The hydrolyzable tannins, carolagin, glucogallin, methyl ester, pelargonin and phyllantusiin (isolated from *Pelargonium reniforme* subsp. *reniforme*) were the most efficient for *C. glabrata*, while 8-carboxy-methyl-1,3,5,6-tetrahydroxyxanthone, a xanthone isolated from *Leiothrix spiralis* was the most effective against *C. krusei* and *C. tropicalis*. Lastly, for *C. parapsilosis*, punicalagin (isolated from *Punica granatum*), followed by 8-carboxy-methyl-1,3,5,6-tetrahydroxyxanthone (isolated from *Leiothrix spiralis*) presented the most pronounced effects.

5. Conclusions

Being part of the commensal flora, *Candida* species are also related to several situations of overgrowth, and an increasing resistance to the current antifungal drugs has been observed. Despite the efforts in order to discover and/or synthesize new chemical molecules, more efficient and effective than the existents, it is necessary to consider that these compounds could be associated to a wide variety of side effects. Beyond that those experiments are very expensive and take a long time until the accreditation/commercialization of the products. Medicinal plants are used since ancient times and present several bioactive properties, due to their richness in some biomolecules, such as phenolic compounds. Several studies have been developed towards investigating the antifungal properties of phenolic compounds enriched extracts, against *Candida* species. For that, different extraction solvents have been used, namely, water, ethanol, methanol, methanol-dichloromethane and even mixtures of the previous. In parallel, and due to the high antifungal potential evidenced by some plant extracts, an increasing number of studies have been evaluating the antifungal potential of phenolic compounds isolated from those matrices.

Despite the existence of a tenuous variation on the antifungal activity according to *Candida* species, in general, the plant extracts obtained using methanol, water and mixtures of the previous (hydroalcoholic extracts) were more effective against *Candida* species than the extracts prepared using a unique solvent. This higher antifungal activity might be related with the solubility of the active principles in the solvents used; phenolic compounds are one of the most abundant bioactive molecules present in those polar extracts. Hydrolysable tannins, followed by some phenolic amides and flavonoids seem to exert the most pronounced anti-*Candida* effects. There are some evidences that the mentioned extracts/compounds affect fungal cells through interaction with the lipid bilayer of the cell membrane. They could act by disrupting the membrane integrity or by intercalation into cell wall and/or DNA. Nevertheless, much more studies are necessary not only to evaluate the antifungal potential of the reviewed and other medicinal plants, but also to establish clearly mechanisms of action involved in potential of phenolic compounds from plant origin against *Candida* species, as also validate the results by in vivo studies.

Acknowledgements

The authors are grateful to Foundation for Science and Technology (FCT, Portugal) for N. Martins grant (SFRH/BD/87658/2012), L. Barros researcher contract under “Programa Compromisso com Ciência – 2008” and financial support to the Research Centre CIMO (strategic project PEst-OE/AGR/UI0690/2011). This work was also supported by the Programa Operacional, Fatores de competitividade – COMPETE and by national funds through FCT – Fundação para a Ciência e a Tecnologia on the scope of the projects FCT

PTDC/SAU-MIC/119069/2010, RECI/EBB-EBI/0179/2012 and PEst-OE/EOB/LA0023/2013. The authors thank the Project “BioHealth – Biotechnology and Bioengineering approaches to improve health quality”, Ref. NORTE-07-0124-FEDER-000027, co-funded by the Programa Operacional Regional do Norte (ON.2-O Novo Norte), QREN, FEDER.

References

- Abdul, B.A., Hassan, A.M., Hassan, A.S., 2012. *In vitro* antimicrobial activity of *Thymus vulgaris*, *Origanum vulgare* and *Rosmarinus officinalis* against dental caries pathogens. *Haitham J. Pure Appl. Sci.* 25, 1–7.
- Abi-Said, D., Anaissie, E., Uzun, O., Raad, I., Pinzowski, H., Vartivarian, S., 1997. The epidemiology of hematogenous candidiasis caused by different *Candida* species. *Clin. Infect. Dis.* 24, 1122–1128.
- Adekunle, A.A., Ikumapayi, A.M., 2006. Antifungal property and phytochemical screening of the crude extracts of *Funtumia elastica* and *Mallotus oppositifolius*. *West Indian Med. J.* 55, 9–13.
- Agarwal, V., Lal, P., Pruthi, V., 2010. Effect of plant oils on *Candida albicans*. *J. Microbiol. Immunol. Infect.* 43, 447–451.
- Ali-Shtayeh, M.S., Ghdeib, S.I.A., 1999. Antifungal activity of plant extracts against dermatophytes. *Mycoses* 42, 665–672.
- Alves, M.J., Ferreira, I.C.F.R., Dias, J., Teixeira, V., Martins, A., Pintado, M., 2013. A review on antifungal activity of mushroom (basidiomycetes) extracts and isolated compounds. *Curr. Top. Med. Chem.* 13, 2648–2659.
- Alves-Silva, J.M., Santos, S.M.D., Pintado, M.E., Pérez-Álvarez, J.A., Fernández-López, J., Viuda-Martos, M., 2013. Chemical composition and in vitro antimicrobial, antifungal and antioxidant properties of essential oils obtained from some herbs widely used in Portugal. *Food Control* 32, 371–378.
- Amara, A.A., Shibl, A., 2013. Role of probiotics in health improvement, infection control and disease treatment and management. *Saudi Pharm. J.*, 1–8.
- Anibal, P.C., Peixoto, I.T.A., Foglio, M.A., Höfling, J.F., 2013. Antifungal activity of the ethanolic extracts of *Punica granatum* L. and evaluation of the morphological and structural modifications of its compounds upon the cells of *Candida* spp. *Braz. J. Microbiol.* 44, 839–848.
- Araújo, M.G.F., Hilário, F., Vilegas, W., Santos, L.C., Brunetti, I.L., Sotomayor, C.E., Bauab, T.M., 2012. Correlation among antioxidant, antimicrobial, hemolytic, and antiproliferative properties of *Leiothrix spiralis* leaves extract. *Int. J. Mol. Sci.* 13, 9260–9277.
- Araújo, M.G.D.F., Hilário, F., Nogueira, L.G., Vilegas, W., Santos, L.C., Bauab, T.M., 2011. Chemical constituents of the methanolic extract of leaves of *Leiothrix spiralis* ruhlund and their antimicrobial activity. *Molecules* 16, 10479–10490.
- Arif, T., Bhosale, J.D., Kumar, N., Mandal, T.K., Bendre, R.S., Lavekar, G.S., Dabur, R., 2009. Natural products – antifungal agents derived from plants. *J. Asian Nat. Prod. Res.* 11, 621–638.
- Arif, T., Mandal, T.K., Dabur, R., 2011. Natural Products: Anti-fungal Agents Derived from Plants, Opportunity, Challenge and Scope of Natural Products in Medicinal Chemistry. pp. 283–311.
- Asgarpanah, J., Kazemivash, N., 2012. Phytochemistry, pharmacology and medicinal properties of *Coriandrum sativum* L. *Afr. J. Pharm. Pharmacol.* 6, 2340–2345.
- Ashour, H.M., 2008. Antibacterial, antifungal, and anticancer activities of volatile oils and extracts from stems, leaves, and flowers of *Eucalyptus sideroxylon* and *Eucalyptus torquata*. *Cancer Biol. Ther.* 7, 399–403.
- Askun, T., Tumen, G., Satil, F., Kilic, T., 2008. Effects of some *Lamiaceae* species methanol extracts on potential mycotoxin producer fungi. *Pharm. Biol.* 46, 688–694.
- Asl, M.N., Hosseinzadeh, H., 2008. Review of pharmacological effects of *Glycyrrhiza* sp. and its bioactive compounds. *Phyther. Res.* 22, 709–724.
- Asmundsdóttir, L.R., Erlendsdóttir, H., Agnarsson, B.A., Gottfredsson, M., 2009. The importance of strain variation in virulence of *Candida dubliniensis* and *Candida albicans*: results of a blinded histopathological study of invasive candidiasis. *Clin. Microbiol. Infect.* 15, 576–585.
- Bakkali, F., Averbeck, S., Averbeck, D., Idaomar, M., 2008. Biological effects of essential oils – a review. *Food Chem. Toxicol.* 46, 446–475.
- Balch, P.A., 2006. Prescription For Nutritional Healing, Fourth ed. Penguin Group, London, England.
- Barros, L., Alves, C.T., Dueñas, M., Silva, S., Oliveira, R., Carvalho, A.M., Henriques, M., Santos-Buelga, C., Ferreira, I.C.F.R., 2013a. Characterization of phenolic compounds in wild medicinal flowers from Portugal by HPLC–DAD–ESI/MS and evaluation of antifungal properties. *Ind. Crops Prod.* 44, 104–110.
- Barros, L., Dueñas, M., Alves, C.T., Silva, S., Henriques, M., Santos-Buelga, C., Ferreira, I.C.F.R., 2013b. Antifungal activity and detailed chemical characterization of *Cistus ladanifer* phenolic extracts. *Ind. Crops Prod.* 41, 41–45.
- Basile, A., Rigano, D., Conte, B., Bruno, M., Rosselli, S., Sorbo, S., 2013. Antibacterial and antifungal activities of acetonic extract from *Paullinia cupana* Mart. Seeds. *Nat. Prod. Res.* 27, 2084–2090.
- Bhalodia, N.R., Nariya, P.B., Acharya, R.N., Shukla, V.J., 2012. *In vitro* antibacterial and antifungal activities of *Cassia fistula* Linn. fruit pulp extracts. *AYU* 33, 123–129.
- Blanco, J.L., Garcia, M.E., 2008. Immune response to fungal infections. *Vet. Immunol. Immunopathol.* 125, 47–70.
- Brunke, S., Hube, B., 2013. Two unlike cousins: *Candida albicans* and *C. Glabrata* infection strategies. *Cell. Microbiol.* 15, 701–708.

- Camacho-Hernández, I.L., Cisneros-Rodríguez, C., Uribe-Beltrán, M.J., Ríos-Morgan, A., Delgado-Vargas, F., 2004. Antifungal activity of fruit pulp extract from *Psidium sartorianum*. *Fitoterapia* 75, 401–404.
- Campos, L., 2008. Entender a bioquímica, 5th ed.
- Cardoso, A.M.R., Cavalcanti, Y.W., Almeida, L.F.D., Pérez, A.L.A.L., Padilha, W.W.N., 2012. Antifungal activity of plant-based tinctures on *Candida*. *RSBO* 9, 25–30.
- Carocho, M., Ferreira, I.C.F.R., 2013a. A review on antioxidants, prooxidants and related controversy: natural and synthetic compounds, screening and analysis methodologies and future perspectives. *Food Chem. Toxicol.* 51, 15–25.
- Carocho, M., Ferreira, I.C.F.R., 2013b. The role of phenolic compounds in the fight against cancer – a review. *Anticancer Agents Med. Chem.* 13, 1236–1258.
- Carrara, M.A., Donatti, L., Damke, E., Svidizinski, T.I.E., Consolaro, M.E.L., Batista, M.R., 2010. A new model of vaginal infection by *Candida albicans* in rats. *Mycopathologia* 170, 331–338.
- Çekiç, S.D., Cetinkaya, A., Avan, A.N., Apak, R., 2013. Correlation of total antioxidant capacity with reactive oxygen species (ROS) consumption measured by oxidative conversion. *J. Agric. Food Chem.* 61, 5260–5270.
- Chaturvedi, R.K., Beal, M.F., 2013. Mitochondrial diseases of the brain. *Free Radic. Biol. Med.* 63, 1–29.
- Choudhary, M.I., Atta-ur-Rahman, 1999. Recent studies on bioactive natural products. *Pure Appl. Chem.* 71, 1079–1081.
- Cooposamy, R.M., Magwa, M.L., 2007. Traditional use, antibacterial activity and antifungal activity of crude extract of *Aloe excelsa*. *Afr. J. Biotechnol.* 6, 2406–2410.
- Correa-Royero, J., Tangarife, V., Durán, C., Stashenko, E., Mesa-Arango, A., 2010. In vitro antifungal activity and cytotoxic effect of essential oils and extracts of medicinal and aromatic plants against *Candida krusei* and *Aspergillus fumigatus*. *Braz. J. Pharmacogn.* 20, 734–741.
- Dai, J., Mumper, R.J., 2010. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules* 15, 7313–7352.
- Darwish, R.M., Aburjai, T.A., 2011. Antimicrobial activity of some medicinal plants against different *Candida* species. *Jordan J. Pharm. Sci.* 4, 70–80.
- Devasagayam, T.P.A., Tilak, J.C., Boloor, K.K., Sane, K.S., Ghaskadbi, S.S., Lele, R.D., 2004. Free radicals and antioxidants in human health: current status and future prospects. *J. Assoc. Phys. India* 52, 794–804.
- Develoux, M., Bretagne, S., 2005. Candidoses et levures diverses. *EMC – Mal. Infect.* 2, 119–139.
- Dib, M.A., Djabou, N., Allali, H., Tabti, B., 2010. Identification of phenolic compounds and antimicrobial activity of roots of *Arbutus unedo* L. *Asian J. Chem.* 22, 4045–4053.
- Doddanna, S.J., Patel, S., Sundarrao, M.A., Veerabhadrapa, R.S., 2013. Antimicrobial activity of plant extracts on *Candida albicans*: an in vitro study. *Indian J. Dent. Res.* 24, 401–405.
- Džamić, A.M., Soković, M.D., Novaković, M., Jadranin, M., Ristić, M.S., Tešević, V., Marin, P.D., 2013. Composition, antifungal and antioxidant properties of *Hyssopus officinalis* L. subsp. *pilifer* (Pant.) Murb. essential oil and deodorized extracts. *Ind. Crops Prod.* 51, 401–407.
- Dzoyem, J.P., Kechia, F.A., Kuete, V., Pieme, A.C., Akak, C.M., Tangmouo, J.G., Lohoue, P.J., 2011. Phytotoxic, antifungal activities and acute toxicity studies of the crude extract and compounds from *Diospyros canaliculata*. *Nat. Prod. Res.* 25, 741–749.
- Dzoyem, J.P., Tangmouo, J.G., Lontsi, D., Etoa, F.X., Lohoue, P.J., 2007. In vitro antifungal activity of extract and plumbagin from the stem bark of *Diospyros crassiflora* Hiern (Ebenaceae). *Phyther. Res.* 21, 671–674.
- Dzoyem, J.P., Tchuengue, R.T., Kuete, J.R., Tekke, G.N., Kechia, F.A., Kuete, V., 2014. In vitro and in vivo antifungal activities of selected Cameroonian dietary species. *BMC Complement. Altern. Med.* 14 (1), 8.
- Eggimann, P., Garbino, J., Pittet, D., 2003a. Epidemiology of *Candida* species infections in critically ill non-immunosuppressed patients. *Lancet Infect. Dis.* 3, 685–702.
- Eggimann, P., Garbino, J., Pittet, D., 2003b. Management of *Candida* species infections in critically ill patients. *Lancet Infect. Dis.* 3, 772–785.
- Endo, E.H., Cortez, D.A.G., Ueda-Nakamura, T., Nakamura, C.V., Filho, B.P.D., 2010. Potent antifungal activity of extracts and pure compound isolated from pomegranate peels and synergism with fluconazole against *Candida albicans*. *Res. Microbiol.* 161, 534–540.
- Epstein, J.B., Polsky, B., 1998. Oropharyngeal candidiasis: a review of its clinical spectrum and current therapies. *Clin. Ther.* 20, 40–57.
- Ertürk, Ö., 2006. Antibacterial and antifungal activity of ethanolic extracts from eleven spice plants. *Biologia (Bratisl)* 61, 275–278.
- Ettefagh, K.A., Burns, J.T., Junio, H.A., Kaatz, G.W., Cech, N.B., 2011. Goldenseal (*Hydrastis canadensis* L.) extracts synergistically enhance the antibacterial activity of berberine via efflux pump inhibition. *Planta Med.* 77, 835–840.
- Fanello, S., Bouchara, J.P., Jousset, N., Delbos, V., LeFlohic, A.M., 2001. Nosocomial *Candida albicans* acquisition in a geriatric unit: epidemiology and evidence for person-to-person transmission. *J. Hosp. Infect.* 47, 46–52.
- Farhat, M., Ben, Jordán, M.J., Chaouech-Hamada, R., Landoulsi, A., Sotomayor, J.A., 2009. Variations in essential oil, phenolic compounds, and antioxidant activity of tunisian cultivated *Salvia officinalis* L. *J. Agric. Food Chem.* 57, 10349–10356.
- Farhat, M., Ben, Landoulsi, A., Chaouech-Hamada, R., Sotomayor, J.A., Jordán, M.J., 2013. Characterization and quantification of phenolic compounds and antioxidant properties of *Salvia* species growing in different habitats. *Ind. Crops Prod.* 49, 904–914.
- Faria, N.C.G., Kim, J.H., Gonçalves, L.A.P., Martins, M. de L., Chan, K.L., Campbell, B.C., 2011. Enhanced activity of antifungal drugs using natural phenolics against yeast strains of *Candida* and *Cryptococcus*. *Lett. Appl. Microbiol.* 52, 506–513.
- Fatima, A., Gupta, V.K., Luqman, S., Negi, A.S., Kumar, J.K., Shanker, K., Saikia, D., Srivastava, S., Darokar, M.P., Khanuja, S.P.S., 2009. Antifungal activity of *Glycyrrhiza glabra* extracts and its active constituent glabridin. *Phyther. Res.* 1193, 1190–1193.
- Fei, L.C., Chen, P.Y., Ahmad, Z., Othman, F., Pei, C., 2008. Antifungal properties of *Allium sativum* extracts on *Candida* species. *J. Trop. Med. Plants* 9, 43–48.
- Franz, R., Kelly, S.L., Lamb, D.C., Kelly, D.E., Ruhnke, M., Morschhäuser, J., 1998. Multiple molecular mechanisms contribute to a stepwise development of fluconazole resistance in clinical *Candida albicans* strains. *Antimicrob. Agents Chemother.* 42, 3065–3072.
- Geiger, A.M., Foxman, B., Sobel, J.D., 1995. Chronic vulvovaginal candidiasis: characteristics of women with *Candida albicans*, *C. Glabrata* and no *Candida*. *Genitourin. Med.* 71, 304–307.
- Ghannoun, M.A., Rice, L.B., 1999. Antifungal agents: mode of action, mechanisms of resistance, and correlation of these mechanisms with bacterial resistance. *Clin. Microbiol. Rev.* 12, 501–517.
- Ghasemi Pirbalouti, A., Hashemi, M., Ghahfarokhi, F.T., 2013. Essential oil and chemical compositions of wild and cultivated *Thymus daenensis* celak and *Thymus vulgaris* L. *Ind. Crops Prod.* 48, 43–48.
- Giada, M.D.L.R., 2013. Food phenolic compounds: main classes, sources and their antioxidant power. In: Morales-González, J.A. (Ed.), *Oxidative Stress and Chronic Degenerative Diseases – A Role for Antioxidants*. InTech, Rio de Janeiro – RJ, p. 501. Chapters.
- Greenberg, M.S., Glick, M., 2003. *Burket's Oral Medicine: Diagnosis and Treatment*, 10 Edition. B.C. Decker Inc., USA.
- Gubbins, P.O., Anaissie, E.J., 2009. Antifungal therapy. In: Livingstone, C. (Ed.), *Clinical Mycology*. Elsevier, pp. 165–199.
- Halberstein, R.A., 2005. Medicinal plants: historical and cross-cultural usage patterns. *Ann. Epidemiol.* 15, 686–699.
- Hashem, M., 2011. Antifungal properties of crude extracts of five Egyptian medicinal plants against dermatophytes and emerging fungi. *Mycopathologia* 172, 37–46.
- Hassawi, D., Kharmar, A., 2006. Antimicrobial activity of some medicinal plants against *Candida albicans*. *J. Biol. Sci.* 6, 109–114.
- Höfling, J.F., Mardegan, R.C., Anibal, P.C., Furlletti, V.F., Foglio, M.A., 2011. Evaluation of antifungal activity of medicinal plant extracts against oral *Candida albicans* and proteinases. *Mycopathologia* 172, 117–124.
- Inbaraj, J.J., Kukielićzak, B.M., Bilski, P., He, Y.-Y., Sik, R.H., Chignell, C.F., 2006. Photochemistry and photocytotoxicity of alkaloids from goldenseal (*Hydrastis canadensis* L.) 2. Palmatine, hydrastine, canadine, and hydrastinine. *Chem. Res. Toxicol.* 19, 739–744.
- Inbaraj, J.J., Kukielićzak, B.M., Bilski, P., Sandvik, S.L., Chignell, C.F., 2001. Photochemistry and photocytotoxicity of alkaloids from goldenseal (*Hydrastis canadensis* L.) 1. Berberine. *Chem. Res. Toxicol.* 14, 1529–1534.
- Irshad, Shreaz, S., Manzoor, N., Khan, L.A., Rizvi, M.M.A., 2011. Anticandidal activity of *Cassia fistula* and its effect on ergosterol biosynthesis. *Pharm. Biol.* 49, 727–733.
- Ishaq, M.S., Hussain, M.M., Afridi, M.S., Ali, G., Khattak, M., Ahmad, S., Shakirullah, 2014. In vitro phytochemical, antibacterial, and antifungal activities of leaf, stem, and root extracts of *Adiantum capillus veneris*. *Sci. World J.*, 1–7.
- Isolauri, E., Kirjavainen, P.V., Salminen, S., 2002. Probiotics: a role in the treatment of intestinal infection and inflammation. *Gut* 50, 54–59.
- Jaikittivong, A., Butsarakamruha, T., Langlais, R.P., 2009. Antifungal activity of *Morinda citrifolia* fruit extract against *Candida albicans*. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol. Endod.* 108, 394–398.
- Jasim, R.M., Al-khalil, I.M.A., 2011. Inhibitory effect of aqueous *Salvia officinalis*'s leaves in the growth of *Candida albicans* from infected women with vaginal candidiasis. *Al-Kindy Coll. Med. J.* 6, 47–49.
- Joe, M.M., Jayachitra, J., Vijayapriya, M., 2009. Antimicrobial activity of some common spices against certain human pathogens. *J. Med. Plants Res* 3, 1134–1136.
- Jung, H.J., Hwang, I.A., Sung, W.S., Kang, H., Kang, B.S., Seu, Y.B., Lee, D.G., 2005. Fungicidal effect of resveratrol on human infectious fungi. *Arch. Pharm. Res.* 28, 557–560.
- Junio, H.A., Sy-Cordero, A.A., Ettefagh, K.A., Burns, J.T., Micko, K.T., Graf, T.N., Richter, S.J., Cannon, R.E., Oberlies, N.H., Cech, N.B., 2011. Synergy-directed fractionation of botanical medicines: a case study with goldenseal (*Hydrastis canadensis*). *J. Nat. Prod.* 74, 1621–1629.
- Kaewpiboon, C., Lirdprapamongkol, K., Srisomsap, C., Winayanuwattikun, P., Yongvanich, T., Puwapisrisiran, P., Svasti, J., Assavalapsakul, W., 2012. Studies of the in vitro cytotoxic, antioxidant, lipase inhibitory and antimicrobial activities of selected Thai medicinal plants. *BMC Complement. Altern. Med.* 12, 1–8.
- Kanafani, Z.A., Perfect, J.R., 2008. Resistance to antifungal agents: mechanisms and clinical impact. *Clin. Infect. Dis. – Antimicrob. Resist.* 46, 120–128.
- Kariba, R.M., Siboe, G.M., Dossaji, S.F., 2001. In vitro antifungal activity of *Schizogygia coffaeoides* bail. (Apocynaceae) extracts. *J. Ethnopharmacol.* 74, 41–44.
- Karomi, M.F., Huda, S.A., Kadhim, S.I., Salman, Z.O., 2012. Effect of methanolic extracts of different parts of *Liquorice Glycyrrhiza glabra* L. on growth of some pathogenic bacteria and fungus *Candida albicans*. *Iraq Acad. Sci. J.* 4, 140–150.
- Kaur, I.P., Kuhad, A., Garg, A., Chopra, K., 2009. Probiotics: delineation of prophylactic and therapeutic benefits. *J. Med. Food* 12, 219–235.

- Kim, J., Sudbery, P., 2011. *Candida albicans*, a major human fungal pathogen. *J. Microbiol.* 49, 171–177.
- Kim, Y., Lee, H., 2012. Anticandidal effect of *Syzygium aromaticum* on biofilm formation, cell surface hydrophobicity, and cell cycle. *J. Med. Plants Res.* 6, 1926–1934.
- Kontoyiannis, D.P., Lewis, R.E., 2002. Antifungal drug resistance of pathogenic fungi. *Lancet* 359, 1135–1144.
- Korukluoglu, M., Gurbuz, O., Sahlan, Y., Yigit, A., Kacar, O., Rouseff, R., 2008. Chemical characterization and antifungal activity of *Origanum onites* L. essential oils and extracts. *J. Food Saf.* 29, 144–161.
- Kosalec, I., Pepelnjak, S., Kustrak, D., 2005. Antifungal activity of fluid extract and essential oil from anise fruits (*Pimpinella anisum* L., Apiaceae). *Acta Pharm.* 55, 377–385.
- Kostić, D.A., Velicković, J.M., Mitić, S.S., Mitić, M.N., Randelović, S.S., 2012. Phenolic content, and antioxidant and antimicrobial activities of *Crataegus oxyacantha* L. (Rosaceae) fruit extract from southeast Serbia. *Trop. J. Pharm. Res.* 11, 117–124.
- Krisch, J., Ördögh, L., Galgóczy, L., Papp, T., Vágvolgyi, C., 2008. Anticandidal effect of berry juices and extracts from *Ribes* species. *Cent. Eur. J. Biol.* 4, 86–89.
- Ksouri, R., Ksouri, W.M., Jallali, I., Debez, A., Magné, C., Hiroko, I., Abdelly, C., 2012. Medicinal halophytes: potent source of health promoting biomolecules with medical, nutraceutical and food applications. *Crit. Rev. Biotechnol.* 32, 289–326.
- Kuete, V., Poumale, H.M.P., Guedem, A.N., Shiono, Y., Randrianasolo, R., Ngadjui, B.T., 2010. Antimycobacterial, antibacterial and antifungal activities of the methanol extract and compounds from *Thecacoris annobonae* (Euphorbiaceae). *South Afr. J. Bot.* 76, 536–542.
- Kurdela, R.R., Lima, B., Tapia, A., Feresin, G.E., Gonzalez Sierra, M., Rodríguez, M.V., Zaccchini, S., Enriz, R.D., Freile, M.L., 2010. Antifungal activity of extracts and prenylated coumarins isolated from *Baccharis darwinii* Hook & Arn. (Asteraceae). *Molecules* 15, 4898–4907.
- Kursat, M., Emre, I., Yılmaz Ö, Erecevit, P., 2011. Antioxidant and antimicrobial activity in the seeds of *Origanum vulgare* L. subs. *gracile* (C. Koch) letswaart and *Origanum acutidens* (Hand. -Mazz.) letswaart from Turkey. *Grasas Aceites* 62, 410–417.
- Kwamin, F., Nartey, N.O., Codjoe, F.S., Newman, M.J., 2013. Distribution of *Candida* species among HIV-positive patients with oropharyngeal candidiasis in Accra, Ghana. *J. Infect. Dev. Ctries* 7, 41–45.
- Lamb, D., Kelly, D., Kelly, S., 1999. Molecular aspects of azole antifungal action and resistance. *Drug Resist. Updat.* 2, 390–402.
- Latte, K.P., Kolodziej, H., 2000. Antifungal effects of hydrolysable tannins and related compounds on dermatophytes, mould fungi and yeasts. *Zeitschrift Naturforsch. C – A J. Biosci.* 55, 467–472.
- Lee, D.G., Park, Y., Kim, M.-R., Jung, H.J., Seu, Y.B., Hahm, K.-S., Woo, E.-R., 2004. Anti-fungal effects of phenolic amides isolated from the root bark of *Lycium chinense*. *Biotechnol. Lett.* 26, 1125–1130.
- Lewis, L.E., Bain, J.M., Lowes, C., Gow, N.A.R., Erwig, L.-P., 2012. *Candida albicans* infection inhibits macrophage cell division and proliferation. *Fungal Genet. Biol.* 49, 679–680.
- Li, S.Y., Yang, Y.L., Chen, K.W., Cheng, H.H., Chiou, C.S., Wang, T.H., Lauderdale, T.L., Hung, C.C., Lo, H.J., 2006. Molecular epidemiology of long-term colonization of *Candida albicans* strains from HIV-infected patients. *Epidemiol. Infect.* 134, 265–269.
- Ličina, B.Z., Stefanović, O.D., Vasić, S.M., Radojević, I.D., Dekić, M.S., Čomić, L.R., 2013. Biological activities of the extracts from wild growing *Origanum vulgare* L. *Food Control* 33, 498–504.
- Lim, S.H., Darah, I., Jain, K., 2006. Antimicrobial activities of tannins extracted from *Rhizophora apiculata* barks. *J. Trop. For. Sci.* 18, 59–66.
- Liu, X.P., Fan, S.R., Bai, F.Y., Li, J., Liao, Q.P., 2009. Antifungal susceptibility and genotypes of *Candida albicans* strains from patients with vulvovaginal candidiasis. *Mycoses* 52, 24–28.
- Lott, T.J., Fundyga, R.E., Kuykendall, R.J., Arnold, J., 2005. The human commensal yeast, *Candida albicans*, has an ancient origin. *Fungal Genet. Biol.* 42, 444–451.
- Lu, Q., Jayatilake, J.A.M.S., Samaranyake, L.P., Jin, L., 2006. Hyphal invasion of *Candida albicans* inhibits the expression of human beta-defensins in experimental oral candidiasis. *J. Invest. Dermatol.* 126, 2049–2056.
- Lubian, C.T., Teixeira, J.M., Lund, R.G., Nascente, P.S., Del Pino, F.A.B., 2010. Antifungal activity of the aqueous extract from *Arctium minus* (Hill) Bernh. (Asteraceae) on oral *Candida* species. *Braz. J. Med. Plants* 12, 157–162.
- Lupetti, A., Danesi, R., Campa, M., Del Tacca, M., Kelly, S., 2002. Molecular basis of resistance to azole antifungals. *Trends Mol. Med.* 8, 76–81.
- Lupetti, A., Nibbering, P.H., Campa, M., Del Tacca, M., Danesi, R., 2003. Molecular targeted treatments for fungal infections: the role of drug combinations. *Trends Mol. Med.* 9, 269–276.
- Madhumitha, G., Saral, A.M., 2011. Preliminary phytochemical analysis, antibacterial, antifungal and anticandidal activities of successive extracts of *Crossandra infundibuliformis*. *Asian Pac. J. Trop. Med.* 4, 192–195.
- Manayi, A., Saeidnia, S., Faramarzi, M.A., Samadi, N., Jafari, S., Vazirian, M., Ghaderi, A., Mirnezami, T., Hadjiakhoondi, A., Ardekani, M.R.S., Khanavi, M., 2013. A comparative study of anti-*Candida* activity and phenolic contents of the calluses from *Lythrum salicaria* L. in different treatments. *Appl. Biochem. Biotechnol.* 170, 176–184.
- Mandeel, Q., Taha, A., 2005. Assessment of in vitro antifungal activities of various extracts of indigenous Bahraini medicinal plants. *Pharm. Biol.* 43, 340–348.
- Martins, N., Barros, L., Santos-Buelga, C., Henriques, M., Silva, S., Ferreira, I.C.F.R., 2015. Evaluation of bioactive properties and phenolic compounds in different extracts prepared from *Salvia officinalis* L. *Food Chem.* 170, 378–385.
- Martins, N., Ferreira, I.C.F.R., Barros, L., Silva, S., Henriques, M., 2014. Candidiasis: predisposing factors, prevention, diagnosis and alternative treatment. *Mycopathologia* 177, 223–240.
- Masoko, P., Mmushi, T.J., Mogashoa, M.M., Mokgotho, M.P., Mampuru, L.J., Howard, R.L., 2008. In vitro evaluation of the antifungal activity of *Sclerocarya birrea* extracts against pathogenic yeasts. *Afr. J. Biotechnol.* 7, 3521–3526.
- Masoko, P., Picard, J., Howard, R.L., Mampuru, L.J., Eloff, J.N., 2010. In vivo antifungal effect of *Combretum* and *Terminalia* species extracts on cutaneous wound healing in immunosuppressed rats. *Pharm. Biol.* 48, 621–632.
- Mathur, A., Singh, R., Yousuf, S., Bhardwaj, A., Verma, S.K., Babu, P., Gupta, V., Prasad, G.B.K.S., Dua, V.K., 2011. Antifungal activity of some plant extracts against clinical pathogens. *Adv. Appl. Sci. Res.* 2, 260–264.
- Maurya, I.K., Thota, C.K., Sharma, J., Tupe, S.G., Chaudhary, P., Singh, M.K., Thakur, I.S., Deshpande, M., Prasad, R., Chauhan, V.S., 2013. Mechanism of action of novel synthetic dodecapeptides against *Candida albicans*. *Biochim. Biophys. Acta* 1830, 5193–5203.
- Mayer, F.L., Wilson, D., Hube, B., 2013. *Candida albicans* pathogenicity mechanisms. *Virulence* 4, 119–128.
- McCullough, M.J., Ross, B.C., Reade, P.C., 1996. *Candida albicans*: a review of its history, taxonomy, epidemiology, virulence attributes, and methods of strain differentiation. *Int. J. Oral Maxillofac. Surg.* 25, 136–144.
- Menezes, T.O.A., Alves, A.C.B.A., Vieira, J.M.S., Menezes, S.A.F., Alves, B.P., Mendonça, L.C.V., 2009. In vitro evaluation of the antifungal activity of essential oils and plant extracts from amazonian region against *Candida albicans*. *J. Odontol. UNESP* 38, 184–191.
- Modaressi, M., Shahsavari, R., Ahmadi, F., Rahimi-Nasrabadi, M., Abiri, R., Mikaeli, A., Batoli, H., 2013. The evaluation of antibacterial, antifungal and antioxidant activity of methanolic extract of *Mindium laevigatum* (Vent.) Rech. F., from central part of Iran. *Jundishapur J. Nat. Pharm. Prod.* 8, 34–40.
- Murray, M.T., 2004. The Healing Power Of Herbs, 2nd ed. Random House, New York, NY.
- Murray, M.T., Pizzorno, J., 1998. Enciclopedia de Medicina Natural, 2nd ed. Ediciones Tutor S.A., Espanha.
- Nelson, D.L., Cox, M.M., 2000. *Lehninger Principles Of Biochemistry*, Third ed. Worth Publishers, Inc., New York.
- Ngameni, B., Kuete, V., Simo, I.K., Mbaveng, A.T., Awoussong, P.K., Patnam, R., Roy, R., Ngadjui, B.T., 2009. Antibacterial and antifungal activities of the crude extract and compounds from *Dorstenia turbinata* (Moraceae). *South Afr. J. Bot.* 75, 256–261.
- Niño, J., Narváez, D.M., Mosquera, O.M., Correa, Y.M., 2007. Antibacterial, antifungal and cytotoxic activities of eight asteraceae and two Rubiaceae plants from Colombian biodiversity. *Braz. J. Microbiol.*, 566–570.
- Noël, T., 2012. The cellular and molecular defense mechanisms of the *Candida* yeasts against azole antifungal drugs. *J. Mycol. Med.* 22, 173–178.
- Nordin, M.-A.-F., Harun, W.H.A.W., Razak, F.A., 2013. Antifungal susceptibility and growth inhibitory response of oral *Candida* species to *Brucea javanica* Linn. extract. *BMC Complement. Altern. Med.* 13, 1–8.
- Ozcelik, B., Kartal, M., Orhan, I., 2011. Cytotoxicity, antiviral and antimicrobial activities of alkaloids, flavonoids, and phenolic acids. *Pharm. Biol.* 49, 396–402.
- Papageorgiou, V., Gardeli, C., Mallouchos, A., Papaioannou, M., Komaitis, M., 2008. Variation of the chemical profile and antioxidant behavior of *Rosmarinus officinalis* L. And *Salvia fruticosa* Miller grown in Greece. *J. Agric. Food Chem.* 56, 7254–7264.
- Parekh, J., Chanda, S., 2008. In vitro antifungal activity of methanol extracts of some Indian medicinal plants against pathogenic yeast and moulds. *Afr. J. Biotechnol.* 7, 4349–4353.
- Parsaieimehr, A., Sargsyan, E., Javidnia, K., 2010. A comparative study of the antibacterial, antifungal and antioxidant activity and total content of phenolic compounds of cell cultures and wild plants of three endemic species of *Ephedra*. *Molecules* 15, 1668–1678.
- Pathak, A.K., 2012. Anti-*Candida* activity of aqueous extracts of some herbs. *Indian J. Fundam. Appl. Life Sci.* 2, 1–6.
- Pendota, S.C., Aderogba, M.A., Ndhala, A.R., Staden, J.V., 2013. Antimicrobial and acetylcholinesterase inhibitory activities of *Buddleja salviifolia* (L.) Lam. leaf extracts and isolated compounds. *J. Ethnopharmacol.* 148, 515–520.
- Pereira Gonzales, F., Maisch, T., 2012. Photodynamic inactivation for controlling *Candida albicans* infections. *Fungal Biol.* 116, 1–10.
- Perlin, D.S., 2009. Resistance to echinocandin-class antifungal drugs. *Drug Resist. Updat.* 10, 121–130.
- Perlin, D.S., 2014. Current perspectives on echinocandin class drugs. *Future Microbiol.* 6, 441–457.
- Pessini, G.L., Filho, B.P.D., Nakamura, C.V., Cortez, D.A.G., 2005. Antifungal activity of the extracts and neolignans from *Piper regnellii* (Miq.) C. DC. var. *pallenscens* (C. DC.) Yunck. *J. Braz. Chem. Soc.* 16, 1130–1133.
- Petrovska, B.B., 2012. Historical review of medicinal plants' usage. *Pharmacogn. Rev.* 6, 1–5.
- Picerno, P., Mencherini, T., Sansone, F., Gaudio, P., Del, Granata, I., Porta, A., Aquino, R.P., 2011. Screening of a polar extract of *Paeonia rockii*: composition and antioxidant and antifungal activities. *J. Ethnopharmacol.* 138, 705–712.
- Pierce, G.E., 2005. *Pseudomonas aeruginosa*, *Candida albicans*, and device-related nosocomial infections: implications, trends, and potential approaches for control. *J. Ind. Microbiol. Biotechnol.* 32, 309–318.
- Politeo, O., Juki, M., Milo, M., 2006. Chemical composition and antioxidant activity of essential oils of twelve spice plants. *Croat. Chem. Acta* 79, 545–552.

- Prabhakar, K., Kumar, L.S., Rajendran, S., Chandrasekaran, M., Bhaskar, K., Sajit Khan, A.K., 2008. Antifungal activity of plant extracts against *Candida* species from oral lesions. *Indian J. Pharm. Sci.* 70, 801–803.
- Proença da Cunha, A., Silva, A.P., Roque, O.R., 2012. Plantas e produtos vegetais em Fitoterapia, 4th ed. Lisboa.
- Proença da Cunha, A., Teixeira, F., Silva, A.P., Roque, O.R., 2010. Plantas na terapêutica: farmacologia e ensaios clínicos, 2nd ed. Lisboa.
- Qader, M.K., Khalid, N.S., Saadullah, A.A.M., 2013. Screening of some plant extracts for antifungal activity against *Candida* species. *Int. J. Microbiol. Immunol. Res.* 1, 65–67.
- Rajeh, M.A.B., Zuraini, Z., Sasidharan, S., Latha, L.Y., Amutha, S., 2010. Assessment of *Euphorbia hirta* L. leaf, flower, stem and root extracts for their antibacterial and antifungal activity and brine shrimp lethality. *Molecules* 15, 6008–6018.
- Raman, S.B., Nguyen, M.H., Cheng, S., Badrane, H., Iczkowski, K.A., Wegener, M., Gaffen, S.L., Mitchell, A.P., Clancy, C.J., 2013. A competitive infection model of hematogenously disseminated candidiasis in mice redefines the role of *Candida albicans* IRS4 in pathogenesis. *Infect. Immun.* 81, 1430–1438.
- Rana, I.S., Rana, A.S., Rajak, R.C., 2011. Evaluation of antifungal activity in essential oil of the *Syzygium aromaticum* (L.) by extraction, purification and analysis of its main component eugenol. *Braz. J. Microbiol.* 42, 1269–1277.
- Ren, B., Xia, B., Li, W., Wu, J., Zhang, H., 2009. Two novel phenolic compounds from *Stenoloma chusanum* and their antifungal activity. *Chem. Nat. Compd.* 45, 182–186.
- Rice-Evans, C.A., Miller, N.J., Paganga, G., 1996. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic. Biol. Med.* 20, 933–956.
- Robbins, R.J., 2003. Phenolic acids in foods: an overview of analytical methodology. *J. Agric. Food Chem.* 51, 2866–2887.
- Rose, J., 1999. Herbal and nutritional support for the immune system. *Clin. Nutr. Insights* 6, 1–4.
- Rubió, L., Motilva, M.-J., Romero, M.-P., 2013. Recent advances in biologically active compounds in herbs and spices: a review of the most effective antioxidant and anti-inflammatory active principles. *Crit. Rev. Food Sci. Nutr.* 53, 943–953.
- Saeed, S., Tariq, P., 2007. Antimicrobial activities of *Emblica officinalis* and *Coriandrum sativum* against gram positive bacteria and *Candida albicans*. *Pakistan J. Bot.* 39, 913–917.
- Safaei-Ghomi, J., Ahd, A.A., 2010. Antimicrobial and antifungal properties of the essential oil and methanol extracts of eucalyptus largiflorens and *Eucalyptus intertexta*. *Pharmacogn. Mag.* 6, 172–175.
- Saleem, M., Nazir, M., Ali, M.S., Hussain, H., Lee, Y.S., Riaz, N., Jabbar, A., 2010. Antimicrobial natural products: an update on future antibiotic drug candidates. *Nat. Prod. Rep.* 27, 238–254.
- Sangamwar, A.T., Deshpande, U.D., Pekamwar, S.S., 2008. Antifungals: need to search for a new molecular target. *Indian J. Pharm. Sci.* 70, 423–430.
- Sanglard, D., 2002. Resistance of human fungal pathogens to antifungal drugs. *Curr. Opin. Microbiol.* 5, 379–385.
- Sanglard, D., Odds Reviews, F.C., 2002. Resistance of candida species to antifungal agents: molecular mechanisms and clinical consequences. *Lancet Infect. Dis.* 2, 73–85.
- Santos-Buelga, C., Gonzalez-Manzano, S., Dueñas, M., Gonzalez-Paramas, A.M., 2012. Extraction and isolation of phenolic compounds. *Methods Mol. Biol.* 864, 427–464.
- Sardi, J.C.O., Scorzoni, L., Bernardi, T., Fusco-Almeida, A.M., Giannini, M.J.S.M., 2013. *Candida* species: current epidemiology, pathogenicity, biofilm formation, natural antifungal products and new therapeutic options. *J. Med. Microbiol.* 62, 10–24.
- Sertel, S., Fu, Y., Zu, Y., Rebacz, B., Konkimalla, B., Plinkert, P.K., Krämer, A., Gertsch, J., Efferth, T., 2011. Molecular docking and pharmacogenomics of *Vinca* alkaloids and their monomeric precursors, vindoline and catharanthine. *Biochem. Pharmacol.* 81, 723–735.
- Shamim, S., Ahmed, S.W., Azhar, I., 2004. Antifungal activity of *Allium*, *Aloe*, and *Solanum* species. *Arch. Physiol. Biochem.* 42, 491–498.
- Sher, A., 2009. Antimicrobial activity of natural products from medicinal plants. *Gomal J. Med. Sci.* 7, 72–78.
- Shojaii, A., Fard, M.A., 2012. Review of pharmacological properties and chemical constituents of *Pimpinella anisum*. *Int. Sch. Res. Netw.* – ISSN Pharm. 2012, 1–8.
- Siler, B., Živković, S., Banjanac, T., Cvetković, J., Živković, J.N., Cirić, A., Soković, M., Mišić, D., 2014. Centauries as underestimated food additives: antioxidant and antimicrobial potential. *Food Chem.* 147, 367–376.
- Silva, F., Ferreira, S., Duarte, A., Mendonça, D.L., Domingues, F.C., 2011a. Antifungal activity of *Coriandrum sativum* essential oil, its mode of action against *Candida* species and potential synergism with amphotericin B. *Phytomedicine* 19, 424–47.
- Silva, S., Henriques, M., Hayes, A., Oliveira, R., Azeredo, J., Williams, D.W., 2011b. *Candida glabrata* and *Candida albicans* co-infection of an in vitro oral epithelium. *J. Oral Pathol. Med.* 40, 421–427.
- de Silva, G.S., 2012. Study of the antimicrobial activity of the medicinal plant extracts against clinical isolates of *Candida* species. State University of Paraíba.
- Silva, N.C.C., Fernandes Júnior, A., 2010. Biological properties of medicinal plants: a review of their antimicrobial activity. *J. Venom. Anim. Toxins Incl. Trop. Dis.* 16, 402–413.
- Singh, A., Duggal, S., Kaur, N., Singh, J., 2010. Berberine: alkaloid with wide spectrum of pharmacological activities. *J. Nat. Prod.* 3, 64–75.
- Singh, R.P., Sharad, S., Kapur, S., 2004. Free radicals and oxidative stress in neurodegenerative diseases: relevance of dietary antioxidants. *J. Indian Acad. Clin. Med.* 5, 218–225.
- Sobel, J.D., 2007. Vulvovaginal candidosis. *Lancet* 369, 1961–1971.
- Souza, N.A.B., Lima, de, E.O., Guedes, D.N., Pereira, F.D.O., Souza, E.L., de, Sousa, F.B., 2010. Efficacy of *Origanum* essential oils for inhibition of potentially pathogenic fungi. *Braz. J. Pharm. Sci.* 46, 499–508.
- Sule, A., Ahmed, Q.U., Latip, J., Samah, O.A., Omar, M.N., Umar, A., Dogarai, B.B.S., 2012. Antifungal activity of *Andrographis paniculata* extracts and active principles against skin pathogenic fungal strains in vitro. *Pharm. Biol.* 50, 850–856.
- Sullivan, D.J., Moran, G.P., Pinjon, E., Almosaid, A., Stokes, C., Vaughan, C., Coleman, D.C., 2004. Comparison of the epidemiology, drug resistance mechanisms, and virulence of *Candida dubliniensis* and *Candida albicans*. *FEMS Yeast Res.* 4, 369–376.
- Sun, S., Lou, H., Gao, Y., Fan, P., Ma, B., Ge, W., Wang, X., 2004. Liquid chromatography–tandem mass spectrometric method for the analysis of fluconazole and evaluation of the impact of phenolic compounds on the concentration of fluconazole in *Candida albicans*. *J. Pharm. Biomed. Anal.* 34, 1117–1124.
- Tarry, W., Fisher, M., Shen, S., Mawhinney, M., 2005. *Candida albicans*: the estrogen target for vaginal colonization. *J. Surg. Res.* 129, 278–282.
- Tharkar, P.R., Tatiya, A.U., Shinde, P.R., Surana, S.J., Patil, U.K., 2010. Antifungal activity of *Glycyrrhiza glabra* Linn. and *Emblica officinalis* Gaertn. by direct bioautography method. *Int. J. PharmTech. Res.* 2, 1547–1549.
- Thirach, S., Tragoolpua, K., Punjaisee, S., Jatisatienr, C., Kharmanwan, C., Kunyanone, N., 2003. Antifungal activity of some medicinal plant extracts against *Candida albicans* and *Cryptococcus neoformans*. *Acta Hort.* 597, 217–221.
- Timothy, S.Y., Wazis, C.H., Adati, R.G., Maspalma, I.D., 2012. Antifungal activity of aqueous and ethanolic leaf extracts of *CCassia alata* Linn. *J. Appl. Pharm. Sci.* 2, 182–185.
- Tintino, S.R., Souza, C.E.S., Guedes, G.M.M., Costa, J.I.V., Duarte, F.M., Chaves, M.C.O., Silva, V.A., Pessôa, H.L.F., Lima, M.A., Garcia, C.A., Coutinho, H.D.M., 2014. Modulatory antimicrobial activity of piper arboreum extracts. *Acta Bot. Croat.* 73, 281–289.
- Tomczykowa, M., Tomczyk, M., Jakoniuk, P., Trzyniewska, E., 2008. Antimicrobial and antifungal activities of the extracts and essential oils of *bidens tripartita*. *Folia Histochem. Cytobiol.* 46, 389–393.
- Touré, A., Bahi, C., Ouattara, K., Djama, J.A., Coulibaly, A., 2011. Phytochemical screening and in vitro antifungal activities of extracts of leaves of *Morinda morindoides* (Morinda, Rubiaceae). *J. Med. Plants Res.* 5, 6780–6786.
- Tsai, P.-W., Chen, Y.-T., Hsu, P.-C., Lan, C.-Y., 2013. Study of *Candida albicans* and its interactions with the host: a mini review. *Biomedicine* 3, 51–64.
- Tsuzuki, J.K., Svidzinski, T.I.E., Shinobu, C.S., Silva, L.F.A., Rodrigues-Filho, E., Cortez, D.A.G., Ferreira, I.C.P., 2007. Antifungal activity of the extracts and saponins from *Sapindus saponaria* L. *Ann. Braz. Acad. Sci.* 79, 577–583.
- Unver, A., Arslan, D., Cetynkaya, Z., Ozcan, M.M., 2008. Antimycotic activity of methanol extracts of sage (*Salvia officinalis* L.), laurel (*Laurus nobilis* L.) and thyme (*Thymbra spicata* L.). *J. Essent. Oil Bear. Plants* 11, 90–95.
- Valko, M., Leibfritz, D., Moncol, J., Cronin, M.T.D., Mazur, M., Telser, J., 2007. Free radicals and antioxidants in normal physiological functions and human disease. *Int. J. Biochem. Cell Biol.* 39, 44–84.
- Vázquez-González, D., Perusquía-Ortiz, A.M., Hunderiker, M., Bonifaz, A., 2013. Opportunistic yeast infections: candidiasis, cryptococcosis, trichosporonosis and geotrichosis. *J. Ger. Soc. Dermatol.* 11, 381–394.
- Volleková, A., Kost'álová, D., Kettmann, V., Tóth, J., 2003. Antifungal activity of *Mahonia aquifolium* extract and its major protoberberine alkaloids. *Phyther. Res.* 17, 834–837.
- Wächter, B., Citiulo, F., Jablonowski, N., Förster, S., Dalle, F., Schaller, M., Wilson, D., Hube, B., 2012. *Candida albicans*–epithelial interactions: dissecting the roles of active penetration, induced endocytosis and host factors on the infection process. *PLoS One* 7, 1–10.
- Walch, S.G., Tinzoh, L.N., Zimmermann, B.F., Stühlinger, W., Lachenmeier, D.W., 2011. Antioxidant capacity and polyphenolic composition as quality indicators for aqueous infusions of *Salvia officinalis* L. (sage tea). *Front. Pharmacol.* 2, 1–6.
- Waldorf, A.R., Polak, A., 1983. Mechanisms of action of 5-fluorocytosine. *Antimicrob. Agents Chemother.* 23, 79–85.
- Wang, J., Lou, J., Luo, C., Zhou, L., Wang, M., Wang, L., 2012. Phenolic compounds from *Halimodendron halodendron* (Pall.) Voss and their antimicrobial and antioxidant activities. *Int. J. Mol. Sci.* 13, 11349–11364.
- Webster, D., Taschereau, P., Belland, R.J., Sand, C., Rennie, R.P., 2008. Antifungal activity of medicinal plant extracts; preliminary screening studies. *J. Ethnopharmacol.* 115, 140–146.
- Westwater, C., Schofield, D.A., Nicholas, P.J., Paulling, E.E., Balish, E., 2007. *Candida glabrata* and *Candida albicans*: dissimilar tissue tropism and infectivity in a gnotobiotic model of mucosal candidiasis. *FEMS Immunol. Med. Microbiol.* 51, 134–139.
- White, T.C., Holleman, S., Dy, F., Laurence, F., Stevens, D.A., Mirels, L.F., 2002. Resistance mechanisms in clinical isolates of *Candida albicans*. *Antimicrob. Agents Chemother.* 46, 1704–1713.
- White, T.C., Marr, K.A., Bowden, R.A., 1998. Clinical, cellular, and molecular factors that contribute to antifungal drug resistance. *Clin. Microbiol. Rev.* 11, 382–402.
- Wojdylo, A., Oszmianski, J., Czemyers, R., 2007. Antioxidant activity and phenolic compounds in 32 selected herbs. *Food Chem.* 105, 940–949.
- Wong, K.S., Tsang, W.K.P., 2009. In vitro antifungal activity of the aqueous extract of *Scutellaria baicalensis* georgi root against *Candida albicans*. *Int. J. Antimicrob. Agents* 34, 283–284.

- Wyk, C., Van, Botha, F.S., Steenkamp, V., 2009. *In vitro* antimicrobial activity of medicinal plants against oral *Candida albicans* isolates. *Int. J. Biomed. Pharm. Sci.* 3, 26–30.
- Xie, Q., Johnson, B.R., Wenckus, C.S., Fayad, M.I., Wu, C.D., 2012. Efficacy of berberine, an antimicrobial plant alkaloid, as an endodontic irrigant against a mixed-culture biofilm in an in vitro tooth model. *J. Endod.* 38, 1114–1117.
- Yang, S.W., Ubillas, R., McAlpine, J., A, Stafford, Ecker, D.M., Talbot, M.K., Rogers, B., 2001. Three new phenolic compounds from a manipulated plant cell culture, *Mirabilis jalapa*. *J. Nat. Prod.* 64, 313–317.
- Yazdani, D.M., Rezazadeh, S., Amin, G., Abidin, M.Z., Shahnazi, S., Jamalifar, H., 2009. Antifungal activity of dried extracts of anise (*Pimpinella anisum* L.) and star anise (*Illicium verum* Hook.) against dermatophyte and saprophyte fungi. *J. Med. Plants* 8, 24–29.
- Zhu, X.F., Zhang, H.X., Lo, R., 2005. Antifungal activity of *Cynara scolymus* L. extracts. *Fitoterapia* 76, 108–111.