

P1: 2

## An in vitro dynamic model of catheter-associated urinary tract infections to investigate the role of uncommon bacteria on the *Escherichia coli* microbial consortium

**Andreia Sofia Azevedo<sup>1,2</sup>, Carina Almeida<sup>1,3</sup>, Luciana Calheiros Gomes<sup>1</sup>, Carla Ferreira<sup>1</sup>, Filipe J. Mergulhão<sup>1</sup>, Luís F. Melo<sup>1</sup>, Nuno Filipe Azevedo<sup>1</sup>**

<sup>1</sup>Laboratory for Process Engineering, Environment, and Energy and Biotechnology Engineering (LEPABE), Department of Chemical Engineering, Faculty of Engineering, University of Porto, Rua Dr Roberto Frias, 4200-465 Porto, Portugal; <sup>2</sup>Nucleic Acid Center, Department of Physics, Chemistry and Pharmacy, University of Southern Denmark, 5230 Odense M, Denmark; <sup>3</sup>Institute for Biotechnology and Bioengineering (IBB), Centre of Biological Engineering, Universidade do Minho, Campus de Gualtar, 4710-057 Braga, Portugal; [asazevedo@fe.up.pt](mailto:asazevedo@fe.up.pt)

About 9% of nosocomial infections are attributed to catheter-associated urinary tract infections (CAUTIs). Uncommon bacteria (*Delftia tsuruhatensis*) have been isolated in CAUTIs in combination with well-established pathogenic bacteria such as *E. coli*. Nonetheless, the reason why *E. coli* coexists with other bacteria instead of outcompeting and completely eliminating them are unknown. As such, a flow cell reactor simulating the hydrodynamic conditions found in CAUTIs (shear rate of 15 s<sup>-1</sup>) was used to characterize the microbial physiology of *E. coli* and *D. tsuruhatensis* individually and in consortium, in terms of growth kinetics and substrate uptake. Single-species biofilms showed that up to 48 h the CFU counts significantly increased for both species ( $p < 0.05$ ). After 48 h, both species stabilized with similar CFU values reaching log 6.24 CFU.cm<sup>-2</sup> for *E. coli* and log 6.31 CFU.cm<sup>-2</sup> for *D. tsuruhatensis* ( $p > 0.05$ ). The assessment of spatial distribution of dual-species biofilms by LNA/2'OMe-FISH revealed that *E. coli* and *D. tsuruhatensis* coexist and tend to co-aggregate over time, which implies that bacteria are able to cooperate synergistically. Substrate uptake measurements revealed that in artificial urine medium the bacteria metabolized lactic acid, uric acid (*E. coli* and *D. tsuruhatensis*) and citric acid (*D. tsuruhatensis*). In the consortium, *D. tsuruhatensis* consumed citric acid more rapidly, presumably leaving more uric acid available in the medium to be used by *E. coli*. In conclusion, metabolic cooperation between *E. coli* and

uncommon species seems to occur when these species share the same environment, leading to the formation of a stable microbial community.