

ON ROTH'S PSEUDO EQUIVALENCE OVER RINGS*

R.E. HARTWIG[†] AND PEDRO PATRICIO[‡]

Abstract.

We characterize the pseudo-equivalence of a block lower triangular matrix $T = [T_{ij}]$ over a regular ring, and its block diagonal matrix $D(T) = [T_{ii}]$, in terms of suitable Roth consistency conditions. The latter can in turn be expressed in terms of the solvability of certain matrix equations of the form $T_{ii}X - YT_{jj} = U_{ij}$.

Key words. Regular rings, Roth equivalence.

AMS subject classifications. 16A30, 15A21

1. Introduction and Definitions. Let \mathcal{R} be a ring with unity 1, let $\mathcal{R}_{m \times n}$ be the set of $m \times n$ matrices over \mathcal{R} , and shorten $\mathcal{R}_{n \times n}$ to \mathcal{R}_n . Throughout all our rings will have an identity.

An element $a \in \mathcal{R}$ is said to be *regular* if $a = axa$, for some x , which is denoted by $x = a^-$. \mathcal{R} is said to be regular if all of its elements are regular. A reflexive inverse a is an element x , such that $axa = a, xax = x$. We shall denote such an inverse of a by a^+ . The sets of inner and reflexive inverses of a , if any, will be respectively denoted by $a\{1\}$ and $a\{1, 2\}$.

DEFINITION 1.1. $m, n \in \mathcal{R}$ are *pseudo-equivalent*, $m \approx n$, provided there exist regular elements p, q and p^-, q^- such that

$$n = pmq, \quad m = p^-nq^-.$$

We may without loss of generality replace the inner inverses p^-, q^- by reflexive inverses p^+, q^+ .

A ring \mathcal{R} is called (von Neumann) *finite* if $ab = 1$ implies $ba = 1$, and it is called *stably finite* if \mathcal{R}_n is finite for all $n \in \mathbb{N}$.

A ring \mathcal{R} is called *unit regular* if for every a in \mathcal{R} , $aua = a$, for some unit u in \mathcal{R} . When $p > q$ then R_p finite implies that R_q is finite.

Matrices A and B are said to be *equivalent*, denoted by $A \sim B$, if $A = PBQ$ for some invertible matrices P, Q . Likewise, matrices A and B are said to be *pseudo-equivalent*, denoted by $A \approx B$, if $A = PBQ$ and $B = P^+BQ^+$ for some square matrices P and Q , with reflexive inverses P^+, Q^+ .

If $A = [A_{ij}]$ is a block matrix over \mathcal{R} with A_{ij} of size $p_i \times q_j$, and $i, j = 1, \dots, n$, then A_k with $k \leq n$, is the *leading* principal block-submatrix $[A_{ij}]$ with $i, j = 1, \dots, k$. The *trailing* principal submatrix is given by $\hat{A}_k = [A_{ij}]$ with $i, j = n-k+1, \dots, n-1, n$,

*Received by the editors on ... Accepted for publication on Handling Editor: ...

[†]Mathematics Department, N.C.S.U. Raleigh, NC 27695-8205, U.S.A. (hartwig@unity.ncsu.edu).

[‡]Departamento de Matemática, Universidade do Minho, 4710-057 Braga, Portugal (pedro@math.uminho.pt). Member of the research unit CMAT – Centro de Matemática da Universidade do Minho, Portugal.

i.e.,

$$A = \begin{bmatrix} A_k & ? \\ ? & \hat{A}_{n-k} \end{bmatrix}.$$

DEFINITION 1.2. For a block matrix $A = [A_{ij}]$ we define its *diagonal* as the block matrix $D(A) = \text{diag}(A_{11}, \dots, A_{nn})$. We further set $D_k(A) = \text{diag}(A_{11}, \dots, A_{kk})$.

2. Roth Conditions. Consider the matrix

$$M = \begin{bmatrix} A & 0 \\ B & D \end{bmatrix}.$$

When the matrix equation $DX - YA = B$ has a solution pair (X, Y) , one checks that

$$(2.1) \quad \begin{bmatrix} I & 0 \\ Y & I \end{bmatrix} \begin{bmatrix} A & 0 \\ B & D \end{bmatrix} \begin{bmatrix} I & 0 \\ -X & I \end{bmatrix} = \begin{bmatrix} A & 0 \\ 0 & D \end{bmatrix},$$

i.e., M is equivalent to its block diagonal matrix $N = \text{diag}(A, D)$.

On the other hand, for regular matrices A and D , the consistency of $DX - YA = B$ is equivalent to the condition $(1 - DD^-)B(1 - A^-A) = 0$, for some and hence all D^-, A^- .

Given consistency, it was shown in [4] that over a regular ring, the general solution to $DX - YA = B$ is given by

$$\begin{aligned} X &= D^-B + D^-ZA + (I - DD^-)W \\ Y &= -(I - DD^-)BA^- + Z - (I - DD^-)ZAA^-, \end{aligned}$$

where W and Z are arbitrary.

In 1952, W.E. Roth proved the converse of 2.1 for matrices over a field \mathbb{F} [16], i.e.,

$$(2.2) \quad DX - YA = B \text{ has a solution pair if and only if } \begin{bmatrix} A & 0 \\ B & D \end{bmatrix} \sim \begin{bmatrix} A & 0 \\ 0 & D \end{bmatrix}.$$

A ring R is said to have *Roth's equivalence property* if the equivalence 2.2 is valid for all matrices over R . Roth's equivalence property was extended in [10], where it was shown that over a unit regular ring,

$$\begin{bmatrix} a & 0 \\ b & d \end{bmatrix} \sim \begin{bmatrix} a & 0 \\ 0 & d \end{bmatrix} \Leftrightarrow dx - ya = b \text{ has a solution pair } (x, y).$$

This result implies that such rings must be finite, and have Roth's equivalence property.

This result was later extended to regular rings by Guralnick [6], who showed that over a regular ring, Roth's equivalence property holds if and only if \mathcal{R} is stably finite. In a

parallel paper [7], Gustafson proved that a commutative ring also must have Roth's equivalence property.

We shall show that in Roth's equivalence property, we may replace equivalence by *pseudo*-equivalence, provided the diagonal blocks A_i are regular and \mathcal{R}_τ is finite, for suitable τ .

3. Lemmata. We begin by deriving some simple consequences of pseudo equivalence.

LEMMA 3.1. If $n \approx m$, then

- (i) \approx is symmetric.
- (ii) m is regular if and only if n is regular.
- (iii) $nR = pmR, Rn = Rmq$.
- (iv) $mR \cong nR$.

Proof. (iv) Let $\phi(mx) = nqx$. Then $nqx_1 = nqx_2 \Rightarrow mx_1 = pnqx_1 = pnqx_2 = mx_2$. Also for any $s, ns = nqq^+s = \phi(mq^+s)$. As such ϕ is a one-one onto module isomorphism, and therefore mR and nR are isomorphic. \square

We may at once apply these to the matrix rings over R , and state

COROLLARY 3.2. If $M \approx D$ and $M \sim M'$, then $R(D) \cong R(M')$.

The following lemma was proved in [11] and characterizes the finiteness of \mathcal{R}_n .

LEMMA 3.3. Let $e, f \in \mathcal{R}$ with $e^2 = e$ and $f^2 = f$. The following conditions are equivalent:

- 1. \mathcal{R} is finite.
- 2. $e\mathcal{R} \subseteq f\mathcal{R}, e\mathcal{R} \cong f\mathcal{R} \implies e\mathcal{R} = f\mathcal{R}$.
- 3. $\mathcal{R}e \subseteq \mathcal{R}f, \mathcal{R}e \cong \mathcal{R}f \implies \mathcal{R}e = \mathcal{R}f$.

We shall apply this to regular matrix rings.

A key result in our reduction is the following "corner" lemma.

LEMMA 3.4.

(a) When $yd^+ = 0$ then $\begin{bmatrix} a & 0 \\ b & d+y \end{bmatrix} \sim \begin{bmatrix} a & 0 \\ (1-dd^+)b(1-a^+) & d \end{bmatrix}$.

(b) If $M = \begin{bmatrix} a & 0 \\ r & d \end{bmatrix}$, with $r = (1-dd^+)b(1-a^+)$, then there exists a 1-2 inverse

$$M^+ = \begin{bmatrix} a^+ & (1-a^+)r^+(1-dd^+) \\ 0 & d^+ \end{bmatrix}$$

such that

$$MM^+ = \begin{bmatrix} aa^+ & 0 \\ 0 & dd^+ + rr^+(1-dd^+) \end{bmatrix},$$

$$M^+M = \begin{bmatrix} a^+a + (1-a^+)r^+r & 0 \\ 0 & d^+d \end{bmatrix}.$$

Proof. (a)

$$\begin{bmatrix} 1 & 0 \\ -ba^+ & 1 \end{bmatrix} \begin{bmatrix} a & 0 \\ b & d+y \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -d^+b(1-a^+a) & 1 \end{bmatrix} = \begin{bmatrix} a & 0 \\ (1-dd^+)b(1-a^+a) & d \end{bmatrix}.$$

(b) This fundamental result was shown in [9], pg 211 eq. (3.5). \square

4. The Cornered Canonical Forms. We next turn to the block triangular case. Let R be a regular ring and let

$$(4.1) \quad M = \begin{bmatrix} A_1 & & & & 0 \\ B_2 & A_2 & & & \\ & & \ddots & & \\ \boxed{B_k} & A_k & & & \\ & \vdots & & \ddots & \\ \boxed{B_n} & & & & A_n \end{bmatrix} = D + B$$

be a lower triangular block matrix with $D = D(M) = \text{diag}(A_1, \dots, A_n)$.

We shall assume that A_i is $p_i \times q_j$ and M is $p \times q$, where $p = \sum_{i=1}^n p_i$ and $q = \sum_{j=1}^n q_j$.

Our aim is to address the question of how to characterize $M \sim D$ and $M \approx D$. The former was done in [12], with aid of the canonical form

$$(4.2) \quad PMQ = P(D + B)Q = N = D + Y = \begin{bmatrix} A_1 & & & & 0 \\ Y_2 & A_2 & & & \\ & & \ddots & & \\ \boxed{Y_k} & A_k & & & \\ & \vdots & & \ddots & \\ \boxed{Y_n} & & & & A_n \end{bmatrix},$$

where

$$(4.3) \quad P = P_n \begin{bmatrix} P_{n-1} & 0 \\ 0 & 1 \end{bmatrix} \cdots \begin{bmatrix} P_2 & 0 \\ 0 & I_{n-2} \end{bmatrix} = \begin{bmatrix} \Delta_k & 0 \\ 0 & I_{n-k} \end{bmatrix},$$

$$Q = \begin{bmatrix} Q_2 & 0 \\ 0 & I_{n-2} \end{bmatrix} \cdots \begin{bmatrix} Q_{n-1} & 0 \\ 0 & 1 \end{bmatrix} Q_n = \begin{bmatrix} \pi_k & 0 \\ 0 & I_{n-k} \end{bmatrix},$$

$$\Delta_k = P_k \begin{bmatrix} P_{k-1} & 0 \\ 0 & 1 \end{bmatrix} \cdots \begin{bmatrix} P_2 & 0 \\ 0 & I_{k-2} \end{bmatrix} = P_k \begin{bmatrix} \Delta_{k-1} & 0 \\ 0 & 1 \end{bmatrix}$$

and

$$\pi_k = \begin{bmatrix} Q_2 & 0 \\ 0 & I_{k-2} \end{bmatrix} \cdots \begin{bmatrix} Q_{k-1} & 0 \\ 0 & 1 \end{bmatrix} Q_k = \begin{bmatrix} \pi_{k-1} & 0 \\ 0 & 1 \end{bmatrix}.$$

In these expressions

$$(4.4) \quad \begin{aligned} P_k &= \begin{bmatrix} I_{k-1} & 0 \\ -B_k \pi_{k-1} D_{k-1}^+ & 1 \end{bmatrix} \text{ and} \\ Q_k &= \begin{bmatrix} I_{k-1} & 0 \\ -A_k^+ B_k \pi_{k-1} (I - D_{k-1}^+ D_{k-1}) & 1 \end{bmatrix} = \begin{bmatrix} I_{k-1} & 0 \\ -\mathbf{q}_k & 1 \end{bmatrix} \end{aligned}$$

are both $k \times k$.

In addition, the submatrices Y_k of N , are defined by

$$(4.5) \quad Y_k = (1 - A_k A_k^+) B_k \pi_{k-1} (I - D_{k-1}^+ D_{k-1}).$$

The reduction as given in 4.2 is equivalent to the ‘‘horizontal’’ reduction

$$(4.6) \quad V^{-1} M (U^{-1} Z) = N = D + Y$$

in which $U = I + D^+ B$, $V = I + B D^+$ and $Z = I + D^+ B D D^+$ are all invertible.

In order to solve the pseudo-equivalence problem, we shall need a second *parallel canonical form*, which again uses Lemma 3.4,

$$(4.7) \quad P' M Q' = P' (D + B) Q' = N' = \begin{bmatrix} A_1 & & & 0 \\ R_2 & A_2 & & \\ & \ddots & & \\ \boxed{R_k} & & A_k & \\ & \vdots & & \ddots \\ \boxed{R_n} & & & A_n \end{bmatrix} = D + R.$$

The steps in this reduction are identical to those used to obtain N , except that we replace D_k by the principal block N'_k at each stage. This gives

$$(4.8) \quad \begin{aligned} P' &= P'_n \cdots \begin{bmatrix} P'_k & 0 \\ 0 & I_{n-k} \end{bmatrix} \cdots \begin{bmatrix} P'_2 & 0 \\ 0 & I_{n-2} \end{bmatrix} = \begin{bmatrix} \Delta'_k & 0 \\ 0 & I_{n-k} \end{bmatrix}, \\ Q' &= \begin{bmatrix} Q'_2 & 0 \\ 0 & I_{n-2} \end{bmatrix} \cdots \begin{bmatrix} Q'_k & 0 \\ 0 & I_{n-k} \end{bmatrix} \cdots Q'_n = \begin{bmatrix} \pi'_k & 0 \\ 0 & I_{n-k} \end{bmatrix}, \end{aligned}$$

and

$$(4.9) \quad \begin{aligned} \Delta'_k &= P'_k \begin{bmatrix} P'_{k-1} & 0 \\ 0 & I \end{bmatrix} \cdots \begin{bmatrix} P'_2 & 0 \\ 0 & I_{k-2} \end{bmatrix} \text{ and} \\ \pi'_k &= \begin{bmatrix} Q'_2 & 0 \\ 0 & I_{k-2} \end{bmatrix} \cdots \begin{bmatrix} Q'_{k-1} & 0 \\ 0 & I \end{bmatrix} Q'_k. \end{aligned}$$

In these products,

$$(4.10) \quad P'_k = \begin{bmatrix} I_{k-1} & 0 \\ -B_k \pi'_{k-1} (N'_{k-1})^+ & I \end{bmatrix} \text{ and}$$

$$Q'_k = \begin{bmatrix} I_{k-1} & 0 \\ -A_k^+ B_k \pi'_{k-1} (I - (N'_{k-1})^+ N'_{k-1}) & I \end{bmatrix} = \begin{bmatrix} I_{k-1} & 0 \\ -\mathbf{q}'_k & I \end{bmatrix}$$

are both are $k \times k$.

It should be noted that

$$\pi'_2 = \pi_2 = Q_2 = \begin{bmatrix} I & 0 \\ -B_2 A_1^+ & I \end{bmatrix}$$

and

$$\Delta'_2 = \Delta_2 = \begin{bmatrix} I & 0 \\ -A_2^+ B_2 (I - A_1^+ A_1) & I \end{bmatrix}.$$

The submatrices R_k of N' in 4.7 are defined by

$$(4.11) \quad R_k = (1 - A_k A_k^+) B_k \pi'_{k-1} (I - (N'_{k-1})^+ N'_{k-1}).$$

There does not seem to be an obvious “horizontal” reduction (using D , B and $(\cdot)^+$) that is equivalent to this “total” block reduction!

We next take advantage of the special form of N' . Using Lemma 3.4 we obtain the following

THEOREM 4.1. Let N' be as in 4.7. Then there exists a reflexive inverse $(N')^+$ such that

$$N'(N')^+ = \text{diag}(A_1 A_1^+, A_2 A_2^+ + R_2 R_2^+ (1 - A_2 A_2^+), \dots, A_n A_n^+ + R_n R_n^+ (1 - A_n A_n^+)) = \mathfrak{D}$$

Proof. If $N'_k = \begin{bmatrix} N'_{k-1} & 0 \\ R_k & A_k \end{bmatrix}$ then by Lemma 3.4 we can find a reflexive inverse

$$(N')_k^+ = \begin{bmatrix} (N')_{k-1}^+ & (I - (N')_{k-1}^+ N'_{k-1}) R_k^+ (1 - A_k A_k^+) \\ 0 & A_k^+ \end{bmatrix}.$$

Recalling that $A_k^+ R_k = 0$ and $R_k (N')_{k-1}^+ = 0$, we may conclude that

$$(4.12) \quad (N')_k (N')_k^+ = \begin{bmatrix} (N')_{k-1} (N')_{k-1}^+ & 0 \\ 0 & A_k A_k^+ + R_k (R_k)^+ (1 - A_k A_k^+). \end{bmatrix}$$

We note in passing that $R_k D_k^+ \neq 0$ in general.

It now follows by induction, that if $N'_{k-1} (N')_{k-1}^+$ is diagonal, then so is $N'_k (N')_k^+$, and has the form

$$N'_k (N')_k^+ = \text{diag}[A_1 A_1^+, A_2 A_2^+ + R_2 (R_2)^+ (1 - A_2 A_2^+), \dots, A_k A_k^+ + R_k (R_k)^+ (1 - A_k A_k^+)]$$

for $k = 1, \dots, n$.

When $k = n$, we arrive at $\mathfrak{D} = N'_n(N'_n)^+$, as desired. \square

Because the product of $A_k A_k^+$ and $I - A_k A_k^+$ is zero, we see that

$$R[A_k A_k^+ + R_k(R_k)^+(1 - A_k A_k^+)] = R[A_k A_k^+] \dot{+} R[R_k(R_k)^+(1 - A_k A_k^+)]$$

as an internal direct sum, and hence that

$$R(A_k) = R(A_k A_k^+) \subseteq R[A_k A_k^+ + R_k(R_k)^+(1 - A_k A_k^+)].$$

This allows us to conclude that

COROLLARY 4.2.

$$R(D) = \begin{bmatrix} R(A_1) \\ \vdots \\ R(A_n) \end{bmatrix} \subseteq \begin{bmatrix} R(A_1) \\ R(A_2) \dot{+} R[R_2(R_2)^+(1 - A_2 A_2^+)] \\ \vdots \\ R(A_n) \dot{+} R[R_n(R_n)^+(1 - A_n A_n^+)] \end{bmatrix} = R(\mathfrak{D}) = R(N'_n)$$

We may now combine Corollaries 3.2 and 4.2 to conclude that

- (i) $R(D) \subseteq R(\mathfrak{D})$.
- (ii) $R(M) \cong R(N') = R[N'(N')^+] = R(\mathfrak{D})$.
- (iii) If $M \approx D$ then $R(D) \cong R(M) \cong R(\mathfrak{D})$.

Now if $R_{p \times p}$ is finite then by Theorem 1 of [10], we may conclude that $R(D) = R(\mathfrak{D})$ and thus $R[R_k R_k^+(1 - A_k A_k^+)] = 0$. This means that $R_k = R_k(R_k)^+ R_k = 0$, for $k = 1, 2, \dots, n$ and we have the Roth Consistency Conditions $R = 0$, i.e.,

$$(4.13) \quad R_k = (1 - A_k A_k^+) B_k \pi'_{k-1} (I - (N')_{k-1}^+ N'_{k-1}) = 0.$$

In order to relate these conditions to the condition that $Y = 0$, we shall need the following

LEMMA 4.3. Let N' be as in 4.7. For each $1 \leq t \leq n$, the following are equivalent.

- (i) $N'_k(N'_k)^+ = D_k D_k^+$, $k = 1, \dots, t$.
- (ii) $R_k = 0$, $k = 2, \dots, t$.
- (iii) $(N'_k)^+ N'_k = D_k^+ D_k$, $k = 1, \dots, t$.
- (iv) $Y_k = 0$, $k = 2, \dots, t$.

Proof. We shall use induction in all four cases.

- (i) \Rightarrow (ii) From Theorem 4.1 we see that for $k \leq t$,

$$N'_k(N'_k)^+ = \begin{bmatrix} (N')_{k-1}^+ (N')_{k-1}^+ & 0 \\ 0 & A_k A_k^+ + R_k(R_k)^+(1 - A_k A_k^+) \end{bmatrix}.$$

If this equals $D_k D_k^+$, then we must have $R_k(R_k)^+(1 - A_k A_k^+) = 0$ for $k \leq t$. Post-multiplication by $B_k \pi'_{k-1} (I - (N')_{k-1}^+ N'_{k-1})$ then shows that $R_k = 0$.

- (ii) \Rightarrow (i). Indeed, if $N'_{k-1}(N')_{k-1}^+ = D_{k-1} D_{k-1}^+$ then setting $R_k = 0$ in 4.12, we see that

$$N'_k(N'_k)^+ = \begin{bmatrix} N'_{k-1}(N')_{k-1}^+ & 0 \\ 0 & A_k A_k^+ \end{bmatrix} = \begin{bmatrix} D_{k-1} D_{k-1}^+ & 0 \\ 0 & A_k A_k^+ \end{bmatrix} = D_k D_k^+.$$

(ii) \Rightarrow (iii). One checks that $N_2^+ N_2 = D_2^+ D_2$. Next we assume it holds for $k = r - 1$. Then from Lemma 3.4 we see that for $N_r^+ N_r = \begin{bmatrix} N_{r-1}^+ N_{r-1} & 0 \\ 0 & A_r^+ A_r \end{bmatrix} = \begin{bmatrix} D_{r-1}^+ D_{r-1} & 0 \\ 0 & A_r^+ A_r \end{bmatrix} = D_r^+ D_r$.

(iii) \Rightarrow (ii). If $N_k^+ N_k = \begin{bmatrix} N_{k-1}^+ N_{k-1} + (I - N_{k-1}^+ N_{k-1}) R_k^+ R_k & 0 \\ 0 & A_k^+ A_k \end{bmatrix} = D_k^+ D_k$, then

$$N_{k-1}^+ N_{k-1} + (I - N_{k-1}^+ N_{k-1}) R_k^+ R_k = D_{k-1}^+ D_{k-1}.$$

For $k = 2$ we have $D_2^+ D_2 = N_2^+ N_2$ and thus $(I - A_1^+ A_1) R_2^+ R_2 = 0$. This gives $R_2 = 0$. Assuming $R_k = 0$ for $k < r$, shows that

$$\begin{aligned} D_r^+ D_r = N_r^+ N_r &= \begin{bmatrix} (N')_{r-1}^+ (N')_{r-1} (I - N_{r-1}^+ N_{r-1}) R_r^+ R_r & 0 \\ 0 & A_r^+ A_r \end{bmatrix} \\ &= \begin{bmatrix} D_{r-1}^+ D_{r-1} + (I - D_{r-1}^+ D_{r-1}) R_r^+ R_r & 0 \\ 0 & A_r^+ A_r \end{bmatrix}. \end{aligned}$$

This gives $(I - D_{r-1}^+ D_{r-1}) R_r^+ R_r = 0$ and hence that $R_r = 0$.

(ii) \Rightarrow (iv). If $N_k^+ N_k = D_k^+ D_k$ for $k \leq t$ then $\pi'_k = \pi_k$ and $R_k = Y_k$.

(iv) \Rightarrow (ii) If $Y_2 = 0$ then clearly $R_2 = 0$. So assume that $R_i = Y_i = 0$ for $i = 2, \dots, k - 1$. Then for these values of i , $N_i^+ N_i = D_i^+ D_i$ and $\pi'_i = \pi_i$. Consequently $R_k = (I - A_k A_k^+) B_k \pi'_{k-1} (I - N_k^+ N_k) = (I - A_k A_k^+) B_k \pi_{k-1} (I - D_k^+ D_k) = Y_k = 0$. \square

We may now combine all the above in

THEOREM 4.4. Let M be a block triangular matrix as in 4.1 and suppose that $R_{p \times p}$ is finite regular, then the following are equivalent:

- (i) $M \sim D(M)$.
- (ii) $M \approx D(M)$.
- (iii) $R_k = 0$, $k = 2, \dots, n$.
- (iv) $Y_k = 0$, $k = 2, \dots, n$.

5. Back to the Roth Conditions. Let us now turn the Roth conditions $Y = 0$, into matrix equations. It should be noted that the equivalent condition $R = 0$ is not so transparent. When $k = 2$, we see that the first Roth consistency condition becomes

$$R_2 = (1 - A_2 A_2^+) B_1 (I - A_1^+ A_1) = 0,$$

which return us to $A_2 X - Y A_1 = B_2$.

For general k , we recall the consistency condition [12]

$$(I - E)BU^{-1}(I - F) = 0$$

in which $E = DD^+$ and $F = D^+D$. Using path products this gives for the (p, q) block,

$$(5.1) \quad (I - A_p A_p^+) [(BU^{-1})_{pq}] (I - A_q^+ A_q),$$

where

$$(BU^{-1})_{pq} = B_{pq} + \sum_{k=1}^r (-1)^k \sum_{p > i_1 > i_2 > \dots > i_k > q} B_{p, i_1} A_{i_1}^+ B_{i_1, i_2} A_{i_2}^+ \dots B_{i_{k-1}, i_k} A_{i_k}^+ B_{i_k, q}.$$

This leads at once to the Roth-matrix consistency condition

$$(5.2) \quad A_p X - Y A_q = (BU^{-1})_{pq} \quad p, q = 1, \dots, n.$$

We may now combine the above with the results of [6] and [7].

THEOREM 5.1. Let R be a regular ring. The following are equivalent:

- (i) R has Roth's pseudo equivalence property, i.e., $M \approx D(M) \Rightarrow Y_k = 0$ ($R_k = 0$).
- (ii) R has Roth's equivalence property, i.e., $M \sim D(M) \Rightarrow Y_k = 0$ ($R_k = 0$).
- (iii) R is stably finite.

Proof. (i) \Rightarrow (ii). This always holds.

(ii) \Rightarrow (iii). This was shown in [6].

(iii) \Rightarrow (i). If R is stably finite and regular, then R_N is finite for all N . Now if in addition, $M \approx D(M)$ then by Theorem 4.4, we see that the Roth consistency conditions $Y_k = 0$ hold for $k = 1, \dots, n$. \square

6. The Row-space Case. The key condition in Theorem 4.1 was that \mathcal{R}_p is finite. This condition can be weakened when $q < p$, to \mathcal{R}_q being finite. To do this we have to repeat the above procedure with *column spaces* instead of row spaces.

The canonical form in this case can be obtained by using Lemma 3.4 starting from the lower right corner rather from the upper left corner and using *trailing* principal submatrices \hat{M}_k . Again we have two canonical forms. One involves D_k , which corresponds to the second horizontal factorization $WV^{-1}MU^{-1} = N = D + Y$. In this case it is more convenient to *reverse* the numbering of the blocks.

$$(6.1) \quad M = \left[\begin{array}{c|c|c|c|c} A_n & & & & 0 \\ \hline & A_{n-1} & & & \\ \hline & & \ddots & & \\ \hline C_n & C_{n-1} & \cdots & A_2 & \\ \hline & & \cdots & \boxed{C_2} & A_1 \end{array} \right] = D + C$$

Even though the new Consistency conditions take a different form from the original Roth conditions, we shall show that we actually do get the *same canonical* matrix M' ! This will become clear once we identify this reduction with the factorization

$$WV^{-1}MU^{-1} = M'.$$

Our aim is to show that the reduction in this case can actually be obtained from the first procedure. We need two concepts.

DEFINITION 6.1. If $A = [a_{ij}]$ is $m \times n$ then

- (i) $\bar{A} = [\bar{a}_{ij}]$, where $a_{ij} = a_{m+1-j, n+1-i}$
- (ii) $A^\sim = [a_{ij}^\sim]$ where in a_{ij}^\sim we *reverse* all products, if any.

The former can be thought of as $\bar{A} = (FAF)^T$, where F is the ‘flip’ matrix

$$\begin{bmatrix} 0 & & 1 \\ & \ddots & \\ 1 & & 0 \end{bmatrix},$$

provided we block transpose.

In particular if $D = \text{diag}(A_1, \dots, A_n)$ then $\bar{D} = \text{diag}(A_n, \dots, A_1)$.

We shall need

THEOREM 6.2. Consider the matrices $A_{m \times k}$, $B_{k \times \ell}$ and $C_{\ell \times n}$ over an arbitrary ring. Then $\overline{(ABC)^\sim} = \bar{C}\bar{B}\bar{A}$

Proof. $(\bar{C}\bar{B}\bar{A})_{ij} = \sum_{u=1}^{\ell} \sum_{v=1}^k (\bar{C})_{iu} (\bar{B})_{uv} (\bar{A})_{vl}$ which in turn equals

$$\sum_{u=1}^{\ell} \sum_{v=1}^k c_{\ell+1-u, n+1-i} b_{k+1-v, \ell+1-u} a_{m+1-j, k+1-v}.$$

Now set $r = \ell + 1 - u$ and $s = k + 1 - v$. This gives

$$(\bar{C}\bar{B}\bar{A})_{ij} = \sum_{r=1}^{\ell} \sum_{s=1}^k c_{r, n+1-i} b_{s, r} a_{m+1-j, s}.$$

On the other hand, $(ABC)_{ij} = \sum_{s=1}^k \sum_{r=1}^{\ell} a_{is} b_{sr} c_{rj}$ and hence

$$(ABC^\sim)_{ij} = \sum_{s=1}^k \sum_{r=1}^{\ell} c_{rj} b_{sr} a_{is}.$$

Next we have $\overline{((ABC)^\sim)}_{ij} = \sum_{s=1}^k \sum_{r=1}^{\ell} c_{r, n+1-i} b_{sr} a_{m+1-j, s}$, which is the (i, j) entry in the RHS. \square

We shall apply this to the reduction $P'MQ' = N'$, as given in 4.8-4.11.

Theorem 4.4 ensures that $\bar{Q}'\bar{M}\bar{P}' = \bar{N}'$, in which

$$(6.2) \quad \bar{N}' = \begin{bmatrix} A_n & & & & 0 \\ \vdots & A_{n-1} & & & \\ \vdots & \vdots & \ddots & & \\ \bar{R}_n & \bar{R}_{n-1} & \cdots & A_2 & \\ \vdots & \vdots & \cdots & \bar{R}_2 & A_1 \end{bmatrix},$$

with $\bar{R}_k = (I - N'_{k-1}(N')_{k-1}^+) \bar{\pi}_{k-1} C_k (I - A_k^+ A_k)$ and

$$\bar{P}' = \begin{bmatrix} I_{n-2} & 0 \\ 0 & \bar{P}'_2 \end{bmatrix} \cdots \bar{P}'_n = \begin{bmatrix} I_{n-k} & 0 \\ 0 & \bar{\Delta}_k \end{bmatrix},$$

with $\bar{\Delta}_k = \begin{bmatrix} I_{k-2} & 0 \\ 0 & \bar{P}'_2 \end{bmatrix} \cdots \begin{bmatrix} 1 & 0 \\ 0 & \bar{P}'_{k-1} \end{bmatrix} \bar{P}'_k$ and

$$\bar{Q}' = \bar{Q}'_n \cdots \begin{bmatrix} I_{n-2} & 0 \\ 0 & \bar{Q}'_2 \end{bmatrix} = \begin{bmatrix} I_{n-k} & 0 \\ 0 & \bar{\pi}_k \end{bmatrix},$$

with $\bar{\pi}_k = \bar{Q}'_k \begin{bmatrix} 1 & 0 \\ 0 & \bar{Q}'_{k-1} \end{bmatrix} \cdots \begin{bmatrix} I_{k-2} & 0 \\ 0 & \bar{Q}'_{k-1} \end{bmatrix}$, in which

$$\bar{P}'_k = \begin{bmatrix} 1 & 0 \\ -(N')_{k-1}^+ \bar{\pi}'_{k-1} C_k & I_{k-1} \end{bmatrix}$$

and

$$\bar{Q}'_k = \begin{bmatrix} 1 & 0 \\ -(I - N'_{k-1}(N')_{k-1}^+) \bar{\pi}'_{k-1} C_k A_k^+ & I_{k-1} \end{bmatrix}.$$

To identify this canonical form we recall that

$$(6.3) \quad \begin{aligned} \Delta'_{k+1} &= P'_{k+1} \begin{bmatrix} \Delta'_k & 0 \\ 0 & 1I \end{bmatrix} = \begin{bmatrix} \Delta'_k & 0 \\ -B_{k+1} \pi'_k D_k^+ \Delta'_k & 1 \end{bmatrix} \text{ and} \\ \pi'_{k+1} &= \begin{bmatrix} \pi'_k & 0 \\ 0 & I \end{bmatrix} Q'_{k+1} = \begin{bmatrix} \pi'_k & 0 \\ -\mathbf{q}'_{k+1} & 1 \end{bmatrix} \end{aligned}$$

Again using Theorem 4.4 we now obtain

$$(6.4) \quad \bar{\pi}'_{k+1} = \begin{bmatrix} 1 & 0 \\ -\bar{\mathbf{q}}'_{k+1} & \bar{\pi}'_k \end{bmatrix} \text{ and } \bar{\Delta}'_{k+1} = \begin{bmatrix} 1 & 0 \\ -\bar{\Delta}_k D_k^+ \bar{\pi}'_k C_{k+1} & \bar{\Delta}'_k \end{bmatrix}.$$

In particular,

$$(6.5) \quad \bar{Q}' = \begin{bmatrix} 1 & & & & 0 \\ \hline & 1 & & & \\ \hline & & \ddots & & \\ \hline -\bar{\mathbf{q}}'_n & -\bar{\mathbf{q}}'_{n-1} & \cdots & 1 & \\ \hline & & \cdots & -\bar{\mathbf{q}}'_2 & 1 \end{bmatrix}.$$

From the form of \bar{Q}' we see that $\bar{Q}' = I - (I - DD^+) \bar{Q}' CD^+$, in which $DD^+ \bar{Q}' = DD^+$ (Here D is labeled backwards!). This gives $\bar{Q}'(I + CD^+) = I + DD^+ CD^+$ from which we see that $\bar{Q}' = (I + DD^+ CD^+)(I + CD^+)^{-1}$. Since $M = D + B = D + C$ (with column partitioning and reverse numbering), we can identify these matrices as

$\bar{Q}' = WV^{-1}$, where $W = I + DD^+CD^+$ and $V = I + CD^+$.

Lastly, if $U = I + D^+C = \begin{bmatrix} ? & ? \\ ? & U_k \end{bmatrix}$ then

$$U_{k+1} = \begin{bmatrix} 1 & 0 \\ D_k^+C_{k+1} & U_k \end{bmatrix}$$

and consequently,

$$U_{k+1}^{-1} = \begin{bmatrix} 1 & 0 \\ -U_k^{-1}D_k^+C_{k+1} & U_k^{-1} \end{bmatrix}.$$

On the other hand, we may match this with 6.4 in which $U_2^{-1} = \Delta_2$. Note that this uses $D_k^+\bar{\pi}'_k = D_k^+$. We have thus shown that $U^{-1} = P'$.

Remarks

(i) The “horizontal” reductions $V^{-1}MU^{-1}Z$ and $WV^{-1}MU^{-1}$, respectively, correspond to the “row” and “column” partitioned cases.

(ii) Since $V^{-1}MU^{-1}Z = WV^{-1}MU^{-1} = D + Y = D + \bar{Y}$, we see that we have two sets of consistency conditions for $M \sim D$, i.e., $Y = 0$ and $\bar{Y} = 0$. That is,

$$(6.6) \quad Y_k = (1 - A_kA_k^+)B_k\pi_{k-1}(I - D_{k-1}^+D_{k-1}) = 0, \quad k = 2, \dots, n$$

and

$$(6.7) \quad \bar{Y}_k = (1 - D_{k-1}D_{k-1}^+)\bar{\pi}_{k-1}C_k(I - A_k^+A_k) = 0, \quad k = 2, \dots, n.$$

These respectively correspond to the *rows* or *columns* of Y being zero.

In conclusion, let us return to Theorem 4.1.

Consider

$$\bar{N}' = \begin{bmatrix} ? & ? \\ ? & \bar{N}'_k \end{bmatrix}$$

and \bar{R}_k as given in 6.2. We may again apply Lemma 3.4 together with $\bar{R}_kA_k^+ = 0$ and $\bar{N}'_{k-1}^+\bar{R}_k = 0$, to obtain a reflexive inverse \bar{N}'_k such that

$$\bar{N}'_k\bar{N}'_k = \begin{bmatrix} A_k^+A_k + (I - A_k^+A_k)\bar{R}_k^+\bar{R}_k & 0 \\ 0 & \bar{N}'_{k-1}^+\bar{N}'_{k-1} \end{bmatrix}$$

It now follows by induction, that if $(\bar{N}')_{k-1}^+\bar{N}'_{k-1}$ is diagonal, then so is $\bar{N}'_k\bar{N}'_k$, and has the form

$$\bar{N}'_k^+\bar{N}'_k = \text{diag}[A_k^+A_k + (I - A_k^+A_k)\bar{R}_k^+\bar{R}_k, \dots, A_2^+A_2 + (1 - A_2^+A_2)\bar{R}_2^+\bar{R}_2, A_1^+A_1]$$

for $k = 1, \dots, n$.

When $k = n$, we arrive at $\bar{\mathcal{D}} = \bar{N}'_n \bar{N}'_n$, as desired.

As before, because the product of $A_k^+ A_k$ and $I - A_k^+ A_k$ is zero, we see that

$$RS[A_k^+ A_k + (1 - A_k^+ A_k) \bar{R}_k^+ \bar{R}_k] = RS(A_k^+ A_k) \dot{+} RS[(1 - A_k^+ A_k) \bar{R}_k^+ \bar{R}_k]$$

as an internal direct sum, and hence that

$$RS(A_k) = RS(A_k^+ A_k) \subseteq RS[A_k^+ A_k + (1 - A_k^+ A_k) \bar{R}_k^+ \bar{R}_k].$$

This allows us to conclude that

COROLLARY 6.3.

$$RS(\bar{D}) = \begin{bmatrix} RS(A_1) \\ \vdots \\ RS(A_n) \end{bmatrix} \subseteq \begin{bmatrix} RS[A_n^+ A_n \dot{+} (1 - A_n^+ A_n) \bar{R}_n^+ \bar{R}_n] \\ \vdots \\ RS[A_2^+ A_2 \dot{+} (1 - A_2^+ A_2) \bar{R}_2^+ \bar{R}_2] \\ RS(A_1^+ A_1) \end{bmatrix} = RS(\bar{\mathcal{D}})$$

We may now combine Corollaries 3.2 and 6.3 to conclude that

- (i) $RS(\bar{D}) \subseteq RS(\bar{\mathcal{D}})$
- (ii) $RS(\bar{M}) \cong RS(\bar{N}') = RS[\bar{N}'^+ \bar{N}'] = RS(\bar{\mathcal{D}})$
- (iii) If $\bar{M} \approx \bar{D}$ then $RS(\bar{D}) \cong RS(\bar{M}) \cong RS(\bar{\mathcal{D}})$.

Now if $R_{q \times q}$ is finite, then by Theorem 1 of [10], we may conclude that $RS(D) = RS(\bar{\mathcal{D}})$ and thus $RS[(1 - A_k^+ A_k) \bar{R}_k^+ \bar{R}_k] = 0$. This means that $\bar{R}_k = \bar{R}_k \bar{R}_k^+ \bar{R}_k = 0$, for $k = 1, 2, \dots, n$ and we have the dual Roth Consistency Conditions $\bar{R} = 0$, i.e.,

$$(6.8) \quad \bar{R}_k = (I - \bar{N}'_{k-1} \bar{N}'_{k-1}^+) \bar{\pi}'_{k-1} C_k (1 - A_k^+ A_k) = 0,$$

with reverse numbering.

As before, we can use the ‘barred’ version of Lemma 4.3 to show that these are equivalent to the simpler consistency conditions that $\bar{Y}_k = 0$ of 6.7.

7. Questions. We close this section with several open questions.

1. Can we improve on the matrix consistency conditions of 5.2 in terms of matrix M ?
2. Can we use the above to derive consistency conditions for the Stein equation?
3. What is the block form of $N^+ N$? Can we use induction?
4. Are there any other applications of the “bar” lemma?
5. What horizontal consistency conditions does the second canonical form correspond to?
6. In Theorem 6.2 do two factors suffice?

Acknowledgement. This research was supported by CMAT – Centro de Matemática da Universidade do Minho, Portugal, and by the Portuguese Foundation for Science and Technology – FCT through the research program POCTI.

REFERENCES

- [1] P. Ara, private communication.
- [2] P. Ara, K.R. Goodearl, K.C. O'Meara and E. Pardo, Diagonalization of matrices over regular rings. *Linear Algebra and Its Applications*, 265:147-163, 1997.
- [3] P. Ara, K.R. Goodearl, K.C. O'Meara and E. Pardo, Separative cancellation for projective modules over exchange rings. *Israel Journal of Mathematics*, 105:105-137, 1998.
- [4] J.K. Baksalary and R. Kala, The matrix equation $AX - YB = C$. *Linear Algebra and Its Applications*, 25:41-43, 1979.
- [5] A. Ben-Israel and T.N.E. Greville, *Generalized Inverses: Theory and Applications*, Wiley, New York, 1974.
- [6] R.M. Guralnick, Roth's theorems and decomposition of modules. *Linear Algebra and Its Applications*, 39:155-165, 1980.
- [7] W.H. Gustafson, Roth's theorems over commutative rings. *Linear Algebra and Its Applications*, 23:245-251, 1979.
- [8] D. Handelman, Perspectivity and cancellation in regular rings. *Journal of Algebra*, 48:1-16, 1977.
- [9] R.E. Hartwig, Block Generalized Inverses, *Arch. Rat. Mech Anal*, 61(3): 13-44, 1976.
- [10] R.E. Hartwig, Roth's equivalence problem in unit regular rings. *Proc. Amer. Math. Soc.*, 59:39-44, 1976.
- [11] R.E. Hartwig and J. Luh, On finite regular rings. *Pacific Journal of Mathematics*, 69(1):73-95, 1977.
- [12] R.E. Hartwig, P. Patrício and R. Puystjens, Diagonalizing triangular matrices via orthogonal Pierce decompositions, *Linear Algebra and its Applications*, 401:381-391, 2005.
- [13] M. Henriksen, On a class of regular rings that are elementary divisor rings. *Archiv der Mathematik*, 24:133-141, 1973.
- [14] D. Huylebrouck, The generalized inverse of a sum with a radical element: Applications. *Linear Algebra and Its Applications*, 246:159-175, 1996.
- [15] P. Patrício and R. Puystjens, About the von Neumann regularity of triangular block matrices. *Linear Algebra and Its Applications*, 332-334:485-502, 2001.
- [16] W.E. Roth, The equations $AX - YB = C$ and $AX - XB = C$, in matrices, *Proc. Amer. Math Soc.* 3: 392-396 (1952); MR 13,900