

Cointegration and the joint confirmation hypothesis

Vasco J. Gabriel^{a,b,*}

^a*Department of Economics, University of Minho, Gualtar, 4710-057, Braga, Portugal*

^b*Department of Economics, Birkbeck College, Oxford, UK*

Abstract

In this paper, the discussion concerning the joint use of unit root and stationarity tests is extended to the case of cointegration. Critical values for testing the joint confirmation hypothesis of no cointegration are computed and a small Monte Carlo experiment evaluates the relative performance of this approach.

Keywords: Cointegration; Joint confirmation hypothesis; Monte Carlo simulations

JEL classification: C12; C15; C22

1. Introduction

The issues of unit roots and cointegration have generated a vast literature in the past few years. More recently, it has been argued that confirmatory analysis (i.e. applying unit root tests in conjunction with stationarity tests) may in some cases lead to a better description of the series, improving upon the separate use of each type of test (see, for example, Amano and Van Norden (1992) and the discussion in Maddala and Kim (1998)). If the two approaches give consistent results, i.e. there is an acceptance *and* a rejection of the nulls, one may conclude whether a given series is stationary or not. On the other hand, if both tests either reject or accept their respective null hypotheses, the results are inconclusive.

Some practical aspects concerning the joint use of unit root and stationarity tests have been addressed by Charemza and Syczewska (1998) and Carrion et al. (2001). The first authors suggest that, instead of conventional individual critical values for each type of test, one should use symmetric critical power values. These correspond to the probability with which the two types of tests make a

*Tel.: +351-253-604-510; fax: +351-253-676-375.

E-mail address: vjgabriel@eeg.uminho.pt (V.J. Gabriel).

wrong decision when both cumulative marginal distributions are equal. Charemza and Syczewska (1998), using Monte Carlo methods, tabulated the new critical values needed for the joint confirmation hypothesis (JCH) of stationarity when the augmented Dickey–Fuller (ADF) and the Kwiatkowski et al. (1992), KPSS henceforth) tests are to be used. However, this approach depends on the parameterization of the autocorrelation in the errors. Hence, Carrion et al. (2001) recommend that the JCH of a unit root should be tested instead, providing a new set of critical values.

In this paper, we study the application of this methodology to cointegration testing. Following Charemza and Syczewska (1998) and Carrion et al. (2001), we show how the testing procedure may be implemented and the related critical values obtained for tests with null hypothesis of no cointegration, as well as null of cointegration. We address the cases where Engle–Granger’s ADF and Phillips–Ouliaris Z_α and Z_τ tests are used in conjunction with the KPSS-type test for the null hypothesis of cointegration developed by McCabe et al. (1997) (see Gabriel (2001) for a comparative study of the properties of null of cointegration tests). Furthermore, the application of the joint confirmation procedure is assessed by means of a set of Monte Carlo experiments, establishing some comparisons with the separate use of each type of test. This is of great interest, since joint testing will be an alternative approach only if it is able to produce better results than individual testing.

The paper proceeds as follows. The next section establishes the notation for the JCH in the context of cointegration, while Section 3 presents the critical values for the JCH of no cointegration. The Monte Carlo study is undertaken in Section 4 and Section 5 concludes.

2. Joint confirmation hypothesis and cointegration

The first step in order to implement the joint use of null of no cointegration and null of cointegration tests is to decide whether one wishes to test the JCH of cointegration or no cointegration. A simple cointegrated model is generally formulated as

$$y_t = x_t' \beta + u_t, \quad (1)$$

where y_t is a scalar $I(1)$ process and x_t is a vector $I(1)$ process of dimension k . The variables y_t and x_t are said to be cointegrated if u_t is $I(0)$, whereas if u_t is $I(1)$ there is no long run equilibrium relationship between y_t and x_t .

A common parameterization for the error process is to assume that u_t is an autoregressive process $u_t = \rho u_{t-1} + \omega_t$, $\omega_t \sim n.i.d.(0, \sigma_\omega^2)$, with $|\rho| < 1$ in the case of cointegration and $\rho = 1$ when there is no cointegration. Another possibility is to consider that under the hypothesis of no cointegration the disturbance u_t may be decomposed into the sum of a random walk and stationary component,

$$u_t = \gamma_t + \varepsilon_t, \quad (2)$$

where the random walk is $\gamma_t = \gamma_{t-1} + \eta_t$, with $\gamma_0 = 0$ and η_t distributed as $i.i.d.(0, \sigma_\eta^2)$, while the stationary part ε_t is distributed as $i.i.d.(0, \sigma_\varepsilon^2)$ and is assumed independent of η_t . Cointegration stems from this formulation when $\sigma_\eta^2 = 0$, so that $\gamma_t = 0$ and no longer is a random walk. Note that in this case the $i.i.d.$ assumption of the errors ($u_t = \varepsilon_t$) is not very realistic, since in empirical applications we should expect some degree of serial correlation. Thus, we may relax this assumption and assume that $\varepsilon_t = \pi \varepsilon_{t-1} + \zeta_t$, ζ_t being $i.i.d.(0, \sigma_\zeta^2)$.

If one chooses to test the JCH of cointegration (meaning $I(0)$ errors), the critical values would

always depend on the value of the autoregressive parameter of the error term, be it ρ if we specify the null hypothesis of cointegration as $H_0: |\rho| < 1$, or π if $H_0: \sigma_\eta^2 = 0$, allowing for autocorrelation in ε_t . It would involve extensive tabulations for a few particular values of ρ (or π), very likely to be different from the actual, unknown value in the empirical situation the researcher is dealing with. Note that this is a similar problem to that pointed out by Carrion et al. (2001) for the univariate case. Therefore, a way to circumvent this obstacle is to specify the JCH of no cointegration (i.e. the residuals have a unit root).

We closely follow the notation of Charemza and Syczewska (1998) and Carrion et al. (2001) by defining the probability of joint confirmation (PJC) of no cointegration as

$$\int_{\tilde{z}_D^{\text{PJC}}}^{\infty} \int_{\tilde{z}_K^{\text{PJC}}}^{\infty} f_{D,K}(z_D, z_K; \Theta, T | H_0^D, H_1^K) dz_D dz_K = \text{PJC}. \quad (3)$$

Here, z_j ($j = D, K$) represents the test statistics (in which we maintain the original notation), D for the ADF t -statistic and K for the KPSS cointegration version of McCabe et al. (1997). The vector of DGP parameters is denoted as Θ , T is the sample size, $f_{D,K}$ is the joint density function, while \tilde{z}_j^{PJC} are the critical values from the joint distribution for a given PJC significance level. As discussed in the above mentioned papers, for each PJC significance level the number of possible critical values is infinite. However, if we impose the restriction that the marginal probabilities (MPr) should be equal, then there is a unique pair $(\tilde{z}_D^{\text{PJC}}, \tilde{z}_K^{\text{PJC}})$ satisfying

$$\int_{\tilde{z}_D^{\text{PJC}}}^{\infty} f_D(z_D; \Theta, T | H_0^D) dz_D = \int_{\tilde{z}_K^{\text{PJC}}}^{\infty} f_K(z_K; \Theta, T | H_1^K) dz_K = \text{MPr}. \quad (4)$$

This restriction means that the probability of deciding wrongly when applying each statistic is equal, that is, when the ADF statistic does not reject the null of no cointegration (type II error) and the KPSS-type test rejects a true null of cointegration (type I error). Such pairs $(\tilde{z}_D^{\text{PJC}}, \tilde{z}_K^{\text{PJC}})$ are dubbed symmetric critical power values (SCPV). Therefore, we find cointegration at a PJC significance level if the joint ADF–KPSS statistic is in the interval $\{(-\infty, \tilde{z}_D^{\text{PJC}}), (0, \tilde{z}_K^{\text{PJC}})\}$, whereas the converse situation leads to a non-rejection of the JCH of no cointegration. In principle, this strategy avoids prioritizing either the cointegration or no cointegration hypotheses, although the practical implications may turn out to be different, as the simulation results in Carrion et al. (2001) show. We will return to this below.

Also note that we may also consider the JCH with other pairs of tests, changing the notation conformably. In fact, we will also consider the joint application of the Phillips–Ouliaris Z_α and Z_t tests, and KPSS-type test. In the next section, critical values for these cases are presented.

3. Critical values for the JCH of no cointegration

As known, critical values for cointegration testing depend not only on the number of regressors k , but also on the deterministic components that may be present in the cointegration space. We will restrict our attention to single equation models with a single cointegration vector. Generalizing (1) as

$$y_t = \alpha + \delta t + x_t' \beta + u_t, \quad (5)$$

where t denotes a time trend, we consider three cases: no constant ($\alpha = \delta = 0$), constant with no trend ($\alpha \neq 0, \delta = 0$) and the model with trend component ($\alpha \neq 0, \delta \neq 0$), up to three regressors.¹ Since we are considering the JCH of no cointegration, then $u_t = u_{t-1} + \omega_t$ ($\rho = 1$), ω_t is assumed to be *n.i.d.*(0, 1) and $u_0 = 0$. We also set $\alpha = 1$ and $\delta = 1$ for the relevant cases. After generating $n = 50,000$ replications for sample sizes $T = 50, 100$ and 250 , pairs of ADF–KPSS, Z_α –KPSS and Z_t –KPSS tests are computed. Using OLS, an appropriate lag length for the ADF test is obtained with a t -test downward selection procedure, by setting the maximum lag equal to 6 and then testing downward until a significant last lag is found, at the 5% level. Concerning Z_α and Z_t , the long run variance is estimated by means of a prewhitened quadratic spectral kernel with an automatically selected bandwidth estimator, using a first-order autoregression as a prewhitening filter, as recommended by Andrews and Monahan (1992). As for the KPSS cointegration statistic, we use Saikkonen's (1991) dynamic least squares estimator and filter the residuals with an ARIMA($p, 1, 1$) model, then using the variance estimator suggested by Leybourne and McCabe (1999) (see McCabe et al. (1997) and Gabriel (2001) for more details on the computation of the statistic).

Again, we follow the methodology of Charemza and Syczewska (1998) and Carrion et al. (2001) to obtain the critical values. Thus, the n pairs of observations are sorted according to the ADF (or Z -type) test and then 250 fractiles are computed. For each of these ADF (Z -type) fractiles, another 250 fractiles were obtained for the KPSS statistic, which means that we get a 250×250 table of empirical joint frequencies. After cumulating these frequencies and thus obtaining the joint distribution function, we may tabulate critical values for the desired significance levels. These are shown in Table 1. The computer routine to obtain these critical values was written in GAUSS² and is an adaptation of the program used by Charemza and Syczewska (1998). Since the ADF and Z_t share the same (marginal) asymptotic distribution and given that the results obtained in the simulations for these two tests are practically the same, we only show the critical values for the ADF test.

4. Monte Carlo experiment

In order to assess the performance of the JCH of no cointegration in terms of classifying the model as cointegrated or not, we devised a set of Monte Carlo simulations. The DGP is similar to the one in Carrion et al. (2001) and is the same as that of the previous section, although the errors are allowed to follow an ARMA(1, 1) process of the form

$$u_t = \rho u_{t-1} + \omega_t + \theta \omega_{t-1}, \quad (6)$$

where ρ takes the values $\{0.5, 0.9, 1\}$ and $\theta = \{-0.8, 0\}$. For simplicity, we only consider a model with a single regressor and a constant term, setting the sample size as $T = 100$ and 250 , computing 2500 replications.

¹Critical values for $k = 4$ and 5 were also computed and are available upon request.

²Available upon request.

Table 1
Critical values

	PJC	No constant			Constant			Trend		
		ADF	MLS	Z_α	ADF	MLS	Z_α	ADF	MLS	Z_α
$T = 50$	0.99	-4.054	0.044	-25.213	-4.703	0.032	-29.256	-5.212	0.061	-35.144
$k = 1$	0.95	-3.301	0.073	-18.721	-3.946	0.042	-23.029	-4.498	0.11	-29.081
	0.90	-2.983	0.107	-15.731	-3.592	0.052	-19.869	-4.184	0.186	-26.181
$k = 2$	0.99	-4.588	0.03	-28.964	-5.095	0.024	-34.365	-5.552	0.033	-38.768
	0.95	-3.879	0.037	-22.376	-4.426	0.029	-29.169	-4.885	0.043	-32.735
	0.90	-3.572	0.044	-19.32	-4.062	0.033	-29.914	-4.547	0.051	-29.968
$k = 3$	0.99	-5.041	0.022	-32.315	-5.477	0.018	-38.227	-5.895	0.023	-41.641
	0.95	-4.332	0.026	-26.127	-4.826	0.021	-32.514	-5.216	0.027	-36.267
	0.90	-3.99	0.029	-22.981	-4.496	0.024	-29.367	-4.902	0.031	-33.454
$T = 100$	0.99	-3.76	0.049	-25.977	-4.434	0.03	-31.871	-4.937	0.072	-38.553
$k = 1$	0.95	-3.185	0.109	-19.118	-3.758	0.047	-23.617	-4.312	0.229	-31.059
	0.90	-2.891	0.241	-16.039	-3.452	0.065	-20.091	-4.017	0.522	-27.416
$k = 2$	0.99	-4.336	0.03	-31.086	-4.866	0.023	-38.252	-5.265	0.038	-44.307
	0.95	-3.738	0.046	-23.508	-4.19	0.03	-29.60	-4.662	0.06	-36.002
	0.90	-3.427	0.063	-19.998	-3.893	0.036	-25.665	-4.368	0.088	-32.40
$k = 3$	0.99	-4.791	0.023	-35.89	-5.244	0.018	-43.747	-5.565	0.026	-48.861
	0.95	-4.177	0.03	-27.931	-4.61	0.022	-35.036	-4.971	0.034	-40.891
	0.90	-3.857	0.036	-24.342	-4.286	0.025	-31.219	-4.699	0.042	-36.914
$T = 250$	0.99	-3.689	0.065	-26.856	-4.255	0.035	-32.179	-4.709	0.136	-40.002
$k = 1$	0.95	-3.118	0.37	-19.473	-3.661	0.073	-23.887	-4.161	0.929	-31.275
	0.90	-2.841	1.262	-16.111	-3.371	0.141	-19.99	-3.894	1.73	-27.549
$k = 2$	0.99	-4.204	0.036	-31.956	-4.717	0.023	-39.71	-5.107	0.053	-47.115
	0.95	-3.631	0.082	-23.857	-4.064	0.039	-29.998	-4.533	0.153	-37.605
	0.90	-3.347	0.221	-20.333	-3.773	0.057	-25.772	-4.234	0.544	-32.874
$k = 3$	0.99	-4.652	0.025	-37.666	-5.05	0.018	-45.734	-5.43	0.034	-52.589
	0.95	-4.085	0.042	-28.826	-4.438	0.026	-35.907	-4.827	0.06	-42.496
	0.90	-3.779	0.07	-24.863	-4.157	0.035	-31.477	-4.566	0.105	-38.156

The results from this simulation exercise are shown in Tables 2 and 3 for $T = 100$ and $T = 250$, respectively. We considered different testing approaches. Firstly, computing each test individually³ and using the respective marginal distributions (i.e. the standard critical values), we gauge the proportion of times that the tests classify a given DGP as being cointegrated (line C) or not cointegrated (line NC), at the 5% level of significance. This corresponds to the usual power-size analysis. Secondly, and still resorting to the 5% critical values from the marginal distributions, we count the frequency a realization of the DGP is classified as cointegrated or not in the following way: (i) if tests for the null of no cointegration (ADF, Z_α and Z_t) reject their null and the KPSS null of cointegration test does not, the process is considered to be cointegrated (C); (ii) if tests for the null of

³This alternative was not considered in Carrion et al. (2001) for the univariate case.

Table 2

Monte Carlo results for ADF, Z_α and KPSS tests ($T = 100$)

(ρ, θ)		ADF	Z_α	KPSS	$D-K$	$Z-K$	$JU(D-K)$	$JU(Z-K)$	$JS(D-K)$	$JS(Z-K)$
(0.5, 0)	C	0.963	1.00	0.793	0.199	0.207	0.04	0.041	0.986	0.994
	NC	0.037	0.00	0.207	0.029	0.00	0.057	0.00	0.00	0.00
	Inc. A	0.00	0.00	0.00	0.764	0.793	0.902	0.959	0.006	0.006
	Inc. B	0.00	0.00	0.00	0.008	0.00	0.001	0.00	0.009	0.00
(0.5, -0.8)	C	1.00	1.00	0.97	0.03	0.03	0.001	0.001	1.00	1.00
	NC	0.00	0.00	0.03	0.00	0.00	0.004	0.00	0.00	0.00
	Inc. A	0.00	0.00	0.00	0.97	0.97	0.995	0.999	0.00	0.00
	Inc. B	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
(0.9, 0)	C	0.29	0.282	0.624	0.102	0.11	0.014	0.017	0.648	0.40
	NC	0.71	0.718	0.376	0.514	0.53	0.832	0.797	0.054	0.096
	Inc. A	0.00	0.00	0.00	0.188	0.172	0.126	0.16	0.082	0.04
	Inc. B	0.00	0.00	0.00	0.196	0.188	0.029	0.026	0.216	0.464
(0.9, -0.8)	C	0.927	1.00	0.938	0.06	0.062	0.007	0.008	0.976	0.994
	NC	0.073	0.00	0.062	0.07	0.00	0.106	0.00	0.00	0.00
	Inc. A	0.00	0.00	0.00	0.868	0.938	0.886	0.992	0.006	0.006
	Inc. B	0.00	0.00	0.00	0.002	0.00	0.001	0.00	0.018	0.00
(1, 0)	C	0.072	0.061	0.062	0.008	0.007	0.00	0.00	0.138	0.054
	NC	0.928	0.939	0.938	0.874	0.883	0.971	0.97	0.405	0.534
	Inc. A	0.00	0.00	0.00	0.063	0.054	0.027	0.028	0.187	0.058
	Inc. B	0.00	0.00	0.00	0.054	0.056	0.002	0.002	0.27	0.354
(1, -0.8)	C	0.595	0.977	0.061	0.041	0.06	0.004	0.006	0.588	0.76
	NC	0.405	0.023	0.939	0.385	0.022	0.475	0.024	0.053	0.003
	Inc. A	0.00	0.00	0.00	0.554	0.917	0.519	0.969	0.18	0.23
	Inc. B	0.00	0.00	0.00	0.02	0.00	0.002	0.00	0.179	0.007

no cointegration do not reject their null and the null of cointegration test does, the process is considered not to be cointegrated (NC); (iii) if both types of tests reject their nulls (inconclusive type A) or do not reject the respective nulls (inconclusive type B), no conclusion is achieved. These joint tests are labeled as $D-K$ for ADF and KPSS tests and $Z-K$ for Z_α and KPSS tests. Finally, a similar exercise is carried out, this time using the 5% critical values from the joint distribution as displayed in Table 1, with the tests denoted as $JU(D-K)$ and $JU(Z-K)$.

From the analysis of Tables 2 and 3, we observe that testing the JCH of no cointegration with $JU(D-K)$ and $JU(Z-K)$ leads to a very small number of correct decisions when the errors are stationary. This is also the case for joint testing with standard critical values. Indeed, most of the times an inconclusive response is obtained, namely rejections by both tests (type A inconclusive answers). Moreover, the results do not seem to improve for larger sample sizes, when we compare Tables 2 and 3. On the other hand, when the DGP is truly non-cointegrated, the JCH approach with $JU(D-K)$ is generally the most accurate in delivering the correct answer, except when a negative MA component is present. In this situation, $JU(D-K)$ still performs better than $D-K$, although the $JU(Z-K)$ version is greatly affected by this error structure. Overall, these results are in accordance with the simulations for univariate testing in Carrion et al. (2001), although with a much poorer performance.

Table 3
 Monte Carlo results for ADF, Z_α and KPSS tests ($T = 250$)

(ρ, θ)		ADF	Z_α	KPSS	$D-K$	$Z-K$	$JU(D-K)$	$JU(Z-K)$	$JS(D-K)$	$JS(Z-K)$
(0.5, 0)	C	1.00	1.00	0.951	0.049	0.049	0.011	0.011	0.806	0.808
	NC	0.00	0.00	0.049	0.00	0.00	0.00	0.00	0.00	0.00
	Inc. A	0.00	0.00	0.00	0.951	0.951	0.989	0.989	0.192	0.192
	Inc. B	0.00	0.00	0.00	0.00	0.00	0.001	0.00	0.002	0.00
(0.5, -0.8)	C	1.00	1.00	0.998	0.002	0.002	0.00	0.00	0.264	0.264
	NC	0.00	0.00	0.002	0.00	0.00	0.00	0.00	0.00	0.00
	Inc. A	0.00	0.00	0.00	0.998	0.998	1.00	1.00	0.736	0.736
	Inc. B	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
(0.9, 0)	C	0.858	0.92	0.702	0.336	0.36	0.121	0.139	0.369	0.061
	NC	0.142	0.08	0.298	0.102	0.064	0.262	0.171	0.194	0.264
	Inc. A	0.00	0.00	0.00	0.522	0.56	0.581	0.672	0.073	0.004
	Inc. B	0.00	0.00	0.00	0.04	0.016	0.036	0.018	0.364	0.671
(0.9, -0.8)	C	0.997	1.00	0.991	0.009	0.009	0.00	0.00	0.381	0.389
	NC	0.003	0.00	0.009	0.003	0.00	0.01	0.00	0.019	0.00
	Inc. A	0.00	0.00	0.00	0.988	0.991	0.99	1.00	0.592	0.611
	Inc. B	0.00	0.00	0.00	0.002	0.00	0.001	0.00	0.008	0.00
(1, 0)	C	0.068	0.06	0.018	0.002	0.002	0.00	0.00	0.002	0.00
	NC	0.932	0.94	0.982	0.916	0.924	0.969	0.967	0.90	0.91
	Inc. A	0.00	0.00	0.00	0.066	0.058	0.03	0.032	0.01	0.00
	Inc. B	0.00	0.00	0.00	0.016	0.016	0.002	0.001	0.088	0.09
(1, -0.8)	C	0.40	0.98	0.004	0.002	0.004	0.001	0.001	0.053	0.157
	NC	0.60	0.02	0.996	0.598	0.02	0.679	0.03	0.65	0.126
	Inc. A	0.00	0.00	0.00	0.399	0.976	0.32	0.969	0.181	0.705
	Inc. B	0.00	0.00	0.00	0.001	0.00	0.00	0.00	0.116	0.012

Comparing this performance with that of individual tests, we see that the latter have a much more reliable behaviour in terms of providing the correct decision, both when there is cointegration and when there is not. The performance of the KPSS cointegration test should be highlighted, given its relative robustness to serial correlation and most especially to the introduction of negative MA components in the errors. In fact, the performance of ADF and Z_α tests, as well as that of joint tests, seems to suffer a great deal with a negative MA error structure, which confirms previous results in the literature.⁴ On the other hand, if we stick to a particular (individual) test, we will not get inconclusive answers, as it happens with the JCH methodology.

Given these results, it would also be interesting to investigate what the outcome would be if one tested the JCH of cointegration. As explained earlier, there is the problem with the critical values depending on the degree of correlation of the errors. However, the researcher could choose an intermediate, though non-optimal, approximation by fixing ρ at an empirically plausible value and use the corresponding critical values. Such a value could be $\rho = 0.75$, which is also recommended and tabulated by Charemza and Syczewska (1998). Of course, if the true ρ is larger than 0.75, the critical

⁴This could eventually be overcome by using the procedure of Ng and Perron (1998), for example.

values would be too conservative, while the converse would lead to overrejecting the JCH of cointegration. Nevertheless, despite the arbitrariness of such a choice, this seems a fairly realistic way to proceed.

Therefore, adapting the methodology discussed in Sections 2 and 3 to the JCH of cointegration, we computed the 5% critical values for the DGP in this section⁵ and evaluated its performance using the same set of simulation experiences. The results are also displayed in Tables 2 and 3, under the columns $JS(D-K)$ and $JS(Z-K)$. We observe that this strategy clearly improves upon that of JCH of no cointegration, since a lot more correct decisions are achieved when the DGP is cointegrated. However, this behaviour is not sustained asymptotically, as the results for $T = 250$ are in general worse. On the other hand, the ability to detect non-cointegrated models improves with the sample size and attains very reasonable levels. Still, this approach does not seem to beat the conventional one, with individual testing.

5. Concluding remarks

In this paper, we extended the joint confirmation hypothesis approach to the context of cointegration. Following Charemza and Syczewska (1998) and Carrion et al. (2001), we tabulated critical values for the JCH of no cointegration. However, our subsequent Monte Carlo simulations, despite its limitations, question the usefulness of such a methodology, as they lead us to conclude that it seems preferable to use the standard individual testing approach, which consistently gave better (or at least as good) results. Indeed, the joint application of different types of tests may obscure, rather than clarify, the process of deciding whether a given model is cointegrated or not. In particular, testing the JCH of no cointegration with the critical values derived here is to be avoided, as it mainly leads to inconclusive answers when the DGP is truly cointegrated. By reversing the JCH to be tested (that is, cointegration), slightly better results are achieved. Further research is required, however, as there are issues that should be addressed. For example, it would be interesting to characterize and compare the behaviour of both types of tests (for the null of cointegration and no cointegration) under different types of DGPs. This line of research is already under study.

Acknowledgements

I wish to thank Wojciech Charemza for kindly making his computer code available. I am also indebted to Josep Lluís Carrion-i-Silvestre, Ron Smith, Haris Psaradakis and Luis Martins for helpful comments, although retaining responsibility for any remaining errors. Financial support from the Fundação para a Ciência e Tecnologia is gratefully acknowledged.

⁵These are -2.482 , -17.023 and 155.421 ($T=100$) and -4.065 , -46.76 and 14.712 ($T=250$) for the ADF, Z_α and KPSS tests, respectively, for $(\rho, \theta)=(0.75, 0)$. A more extensive tabulation is available upon request.

References

- Amano, R., Van Norden, S., 1992. Unit-Root Tests and the Burden of Proof. Bank of Canada.
- Andrews, D.W.K., Monahan, J.C., 1992. An improved heteroskedasticity and autocorrelation consistent covariance matrix estimator. *Econometrica* 60, 953–966.
- Carrion, J.L.I., Sansó, A.I., Ortuño, M.A., 2001. Unit root and stationarity tests' wedding. *Economics Letters* 70, 1–8.
- Charemza, W.W., Syczewska, E.M., 1998. Joint application of the Dickey–Fuller and KPSS tests. *Economics Letters* 61, 17–31.
- Gabriel, V.J., 2001. Tests for the null hypothesis of cointegration: a Monte Carlo comparison, Manuscript.
- Kwiatkowski, D., Phillips, P.C.B., Schmidt, P., Shin, Y., 1992. Testing the null hypothesis of stationarity against the alternative of a unit root. *Journal of Econometrics* 54, 159–178.
- Leybourne, S.J., McCabe, B.P.M., 1999. Modified stationarity tests with data-dependent model-selection rules. *Journal of Business and Economic Statistics* 17, 264–270.
- Maddala, G.S., Kim, I.M., 1998. *Unit Roots, Cointegration and Structural Change*. Cambridge University Press, Cambridge.
- McCabe, B.P.M., Leybourne, S.J., Shin, Y., 1997. A parametric approach to testing for the null of cointegration. *Journal of Time Series Analysis* 18 (4), 395–413.
- Ng, S., Perron, P., 1998. An autoregressive spectral density estimator at frequency zero for nonstationarity tests. *Econometric Theory* 14, 560–603.
- Saikkonen, P., 1991. Asymptotically efficient estimation of cointegrating regressions. *Econometric Theory* 7, 1–21.