Topic: Ecosystem processes

Long-term effects of riparian-plant diversity loss on a stream invertebrate shredder

Isabel Fernandes<sup>\*</sup>, Sofia Duarte, Cláudia Pascoal, Fernanda Cássio Centre of Molecular and Environmental Biology (CBMA), Department of Biology, University of Minho, Braga, Portugal <u>\*isabelrodriguesfernandes@bio.uminho.pt</u>

We used a pool of 3 common riparian-plant species in Portugal (alder, oak and eucalyptus) to examine the potential long-term impacts of riparian diversity loss on the feeding behaviour and body composition of a stream invertebrate shredder (Limnephilidae). Fine-mesh bags containing mixtures of the 3 leaf species were immersed in a mixed-forested stream to allow microbial colonization. After 2 weeks, colonized leaves were transferred to microcosms containing stream water and all combinations of 1 to 3 non-colonized leaf species enclosed in mesh containers. The experiment ran for 6 months. In each month, a portion of leaf material was used to inoculate new microcosms containing non-colonized leaf material, keeping the leaf species composition constant. After 2 and 6 months, those leaves were used to feed invertebrate shredders for a period of 8 days. Leaf consumption and FPOM production by the shredder were affected by leaf species diversity, with higher values in mixtures with 3 leaf species. Leaf mass loss and FPOM production were also affected by leaf species identity, but not by time of leaf diversity loss. C:N ratio of FPOM was affected by leaf identity and showed a positive linear relationship with initial leaf C:N ratio. Invertebrate C:N ratio increased along time of leaf diversity loss. Results suggest that the riparian-plant diversity loss can affect leaf consumption, FPOM production and invertebrate body composition, but effects did not appear to change at longer times.

FEDER-POFC-COMPETE and FCT supported this study (PEst-C/BIA/UI4050/2011, PTDC/AAC-AMB/117068/2010), IF (SFRH/BD/42215/2007) and SD (SFRH/BPD/47574/2008).