



**Universidade do Minho**

Escola de Ciências

Pedro Humberto Araújo Ramos Fernandes Castro

**Functional analysis of the SUMO  
conjugation/deconjugation system during  
the development and stress response of  
*Arabidopsis thaliana***

Tese de Doutoramento em Ciências  
Especialidade em Biologia

Trabalho realizado sob a orientação do:

**Prof. Doutor Herlânder Azevedo**

**Prof. Doutor Rui Manuel Tavares**

**Prof. Doutor Eduardo Rodríguez Bejarano**

Junho de 2013

## **DECLARAÇÃO**

**Nome:** Pedro Humberto Araújo Ramos Fernandes Castro

**Endereço Eletrónico:** pedro1berto@gmail.com

**Telefone:** +351 963546507

**Número do Cartão de Cidadão:** 12532609

### **Título da Tese de Doutoramento:**

Functional analysis of the SUMO conjugation/deconjugation system during the development and stress response of *Arabidopsis thaliana*

### **Orientadores:**

Prof. Doutor Herlânder Azevedo  
(Departamento de Biologia, Escola de Ciências, Universidade do Minho, Portugal)

Prof. Doutor Rui Manuel Tavares  
(Departamento de Biologia, Escola de Ciências, Universidade do Minho, Portugal)

Prof. Doutor Eduardo Rodríguez Bejarano  
(Departamento de Biología Celular, Genética y Fisiología, Universidad de Málaga, Espanha)

**Ano de Conclusão:** 2013

**Designação do Ramo de Conhecimento do Doutoramento:** Ciências

**Especialidade:** Biologia

É AUTORIZADA A REPRODUÇÃO PARCIAL DESTA TESE APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.

Universidade do Minho, cinco de junho de 2013

Assinatura: \_\_\_\_\_

## ACKNOWLEDGMENTS

These years as a PhD student were intense, full of hard work, yet lived with enthusiasm. This opportunity made me grow as scientist and as person. Now is the time to express my gratitude to those who contributed to the success of this step in my academic career.

My sincere acknowledgments go to my supervisor Herlânder Azevedo and my co-supervisors Rui Tavares and Eduardo Bejarano for giving me the opportunity to become their PhD student.

Once more, Herlânder thank you very much for your guidance, dedication, enthusiasm, and friendship. Most of all, thank you for believing more than anyone in my ideas and work capacity. I will never be able to thank you enough. I am sure that all the work will be translated into great contributions to the scientific progress. My gratitude to Rui, who may not know this, but was one of the causes I have decided to follow research in Plant Biology; thank you for the academic discussions and for helping me with all the never-ending bureaucratic procedures.

During these years, I had the opportunity to interchange knowledge and share lab life with many colleagues. To those who contributed directly or indirectly, I would like to express my appreciation: Teresa Lino-Neto and her group (Cátia, Cláudia, Daniela, Francisca, Joana, Jules, Manu, Óscar and Rute); Manuela Costa and her group (Eduarda, Helena, Inês, Joana, João and Rómulo); Hernâni and his group (Alice, Artur, Carlos Conde, Enrique, Natacha, Paulo, Richard and Viviana); and Alberto Dias and his group (Franklin, Mafalda and Marta).

A special mention goes to my past and more recent pupils in the lab: Daniel Couto, Eva Dias, Sara Freitas, Miguel Ângelo and Vânia Passos. Daniel my first full-time lab pupil, you learned fast and well. No doubt you will have a bright scientific future. I miss you around, all the freak-out mixed with Seinfeld moments. Eva Dias, I'll take this opportunity to confess that it is impossible to get mad at you, no matter how chaotic things go. You're very funny and I'll always remember nostalgically the Eva Dias supermarkets. Sara Freitas, my laboratorial "granddaughter", it is a pleasure to witness your great improvement in the lab. I believe that you will reach high in science. To Miguel Ângelo and Vânia Passos, I wish all the best for your future projects.

In addition, I want to acknowledge all the staff and colleagues at the Biology Department for all their support.

My special thanks to my very good friend Vítor, for all the partnership during these academic years. I do not have enough words to express my gratitude for your faithful friendship and willingness to help. Many career choices were unavoidably done under your influence. Everyone should learn with you how to see the optimistic angle of every detail. João Neves, you're like a brother that I could, in fact, choose. That was highlighted in all your support and encouragement during these years. Sara (沙蘭) Laranjeira, I admire your sincerity and I'm grateful for your truthful friendship and for the reciprocal encouragement during the Malaga periods. To my 4<sup>th</sup> of November friends, Isabel and Jorge, my gratitude for all the partnership and may it last for many many years.

A special thanks to all my friends and colleagues in Malaga that made my stay enjoyable and filled with good memories. First of all, I would like to express my deepest appreciation to my supervisor at University of Malaga, Eduardo, that always supported my experiments and made me feel comfortable at work. Several non-official supervisors were always helpful, with inputs and advices for my work, for that I would like to express my gratitude to my “second supervisors” from Malaga, Javier Ruiz-Albert, Araceli Castillo and Miguel Angel Botella.

My deepest gratitude to my group in Malaga, “the tower geneticists”: Manolo and Tabata were like my siblings in Spain; they saved me in moments of great need. Manolo you are an amazing teacher, many of the results in here were possible thanks to your lessons and advices. Tabata you have treated me like a little brother and that was also true in the lab, where you were many times an invaluable help. Rosa and Alberto, I have to say that you are a big inspiration to me, and you’re (although you’re ahead of common beings) examples of how a scientist should be. Ana Luna, you were always a good-mood emitter and a great friend. My partner in the lab and office, Natasa, thank you for your warm friendship. Edgar, the gene expression master and my sport fellow, thank you for your proverbs and scientific discussions. Thanks Zaira for taking care of us. Carlos Guevara was my lab boss during the first stay in Malaga, who guided me in the lab. Migue, thank you for teaching me how to sumoylate. Best wishes to the young students in the lab: Álvaro, Carlos, Cristina, Esther, Inma, Manu, Martín, Mauricio and Tamara.

My acknowledgments to members of the *Departamento de Biología Celular, Genética y Fisiología*, with special focus to Carmen Beuzón and her group (Adela, Ainhoa, Alberto, Diego, Inma, Jose, Juanjo and Spiridoula) and the technicians Mayte and Silvia. A special thanks to Alberto Macho and Jose Rufián for all the help during the bacterial infection experiments. Adela, thank you for sharing your experience with RT-PCR.

My acknowledgments to all the members of the *Laboratorio de Bioquímica y Biotecnología Vegetal*: Abel, Ali, Ana Chiqui, Arni, Carmen, Cristina, Cristobal, David, Edu, Eli, Fabiana, Irene Araguez, Irene Nevado, Jessi, Jose, Julio, Karen, Ludwig, Mayte, Naoufal, Paqui, Sonia, Vero, Viviana, and of course to Professors Victoriano Valpuesta.

Vero and Ian were my friends at first sight, and did everything possible to keep me from feeling lonely, making Malaga feel as home. I predict an amazing scientific future for both of you. Ana “Chiqui”, you’re a hard worker and I hope the future makes you justice. Thank you for all the innumerable and unpredictable good moments. Edu thank you for all the helpful advices and companionship. Sonia and Jose Vallarino thank you for the many recommendations and insights. Eli thank for your companion in the lab, you’re very funny. The “king” Naoufal to set the sportive calendar and for being always so positive. Ali and Paqui, thank you for your availability to help and support.

A very special thanks to my parents, without whom my scientific career would not be possible. For the unlimited support, belief, and encouragement to achieve my goals.

Last but certainly not least, Catarina, thank you for being my peace in this world, to you my heart and soul.

I would like to dedicate this thesis to my grandfather David Ramos,

*“El-rei não manda chover, manda marchar”*

O presente trabalho beneficiou do seguinte apoio da Fundação para a Ciência e a Tecnologia:

**Bolsa de Doutoramento - SFRH/BD/44484/2008**

Bolsa de Investigação no âmbito do QREN - POPH - Tipologia 4.1 - Formação Avançada, participado pelo Fundo Social Europeu e por fundos nacionais do MCTES.





# Functional analysis of the SUMO conjugation/deconjugation system during the development and stress response of *Arabidopsis thaliana*

## ABSTRACT

Living organisms are subjected to constantly changing environmental conditions that compromise survival. Perception of these changes precedes the triggering of signaling events that ultimately activate an adequate response. Since plants are sessile organisms they are considerably more exposed to an instable environment, having adopted a wide range of strategies to adapt, avoid or tolerate stress. Knowledge on the molecular basis of these strategies will be key for future crop improvement, particularly with a rapidly changing environment. Several studies have singled out how fast and reversible responses to stress correlate with post-translational modification (PTM) of key proteins. An increasingly important mechanism involves protein modification by small peptides such as ubiquitin and other ubiquitin-like modifiers (UBLs). Focus on small ubiquitin-related modifier (SUMO), a UBL family member, has increased massively in the last years. SUMO attachment, or sumoylation, may exert different effects depending on the target protein, controlling its conformation, or even creating or blocking interacting interfaces. To be attached to a target, pre-SUMO peptides are first processed by SUMO proteases (ULP/SENPs family), and then conjugated to a target's lysine via SUMO E1 activases and SUMO E2 conjugases, aided by SUMO E3 ligases. Deconjugation of the SUMO peptide can be carried out by SUMO proteases. In plants, SUMO homeostasis is critical because mutations in pathway components result in embryonic lethality or pleiotropic phenotypes. One interesting feature of SUMO is that SUMO-conjugates accumulate rapidly upon stress imposition, placing SUMO in the forefront of the plant response to stress, most likely associated to transcriptional re-programming. Studies of SUMO function in plants have been based mostly on reverse genetics approaches in the plant model *Arabidopsis thaliana*. Similarly, we employed T-DNA insertion mutants to characterize the role of the major SUMO E3 ligase SIZ1 and SUMO proteases ULP1c, ULP1d, ULP2a and ULP2b in the plant stress response.

Reactive oxygen species (ROS) are known internal signals caused by stress that can induce SUMO-conjugate accumulation. In the present work we addressed SUMO-ROS interplay, and report that SIZ1 is essential for SUMO-conjugate induction in response to ROS, and is involved in oxidative stress tolerance. Additionally, *siz1* mutants displayed altered ROS homeostasis, constitutively accumulating hydrogen peroxide, superoxide ion and singlet oxygen. The *siz1* phenotype was partially dependent of salicylic acid (SA) levels since several *siz1* defects were greatly recovered by the expression of the transgenic salicylate hydroxylase *NahG*. The *siz1* mutant displays constitutive autoimmune phenotypes, including SA-accumulation. Analysis of oxidative stress-responsive targets of sumoylation suggests a major role for transcription remodeling proteins in the SUMO-dependent response to oxidative stress.

Bioinformatic analysis revealed that the transcriptome profile of adult *siz1* is similar to that of mitogen-activated protein kinase (MAPK) pathway mutants. These also share common features of autoimmune responses, such as the accumulation of ROS. Results suggest a crosstalk between the SUMO and MAPK pathways, but a direct interaction between MKK2-MPK4 and SUMOs or in vitro sumoylation of MKK2-MPK4 was not observed. Nonetheless, we introgressed *mkk1/2* and *mpk4*

mutants into *siz1*, and sumoylation profiles suggest that MPK4 acts as a negative regulator while MKK1/2 is seemingly epistatic to SIZ1.

In contrast to the low number of components involved in SUMO conjugation, there are several SUMO proteases in plants. SUMO proteases are sources of selectivity, since they can discriminate different SUMO isoforms and targets to be de-sumoylated. They also display different expression patterns and subcellular localizations. We report how phylogenetic reconstruction of the Arabidopsis ULP family outlines the existence of two major branches, and within these, four phylogenetic subgroups. We characterized two of these subgroups, one composed by ULP1c/ULP1d and another by ULP2a/ULP2b. We show how ULP1c/d function redundantly to regulate several developmental traits. Using GUS report assays we also established their spatial expression pattern. A microarray analysis of differentially expressed genes in a double T-DNA insertion mutant was carried out, suggesting a deregulation of drought and abscisic acid-related genes. Taking this into consideration, a characterization of related phenotypes was performed in the double mutant.

We additionally demonstrated how ULP1c/d are negative regulators of the plant response against the hemibiotroph pathogen *Pseudomonas syringae* pv. *tomato* (*Pst*) DC3000. The response is however complex, since infection triggered down-regulation of *ULP1c* and *ULP1d* transcripts, which may correlate with the observed increase in SUMO-conjugates that follows infection. Analysis of the transcriptome of *ulp1c/d* after *Pst* DC3000 challenging revealed the existence of many de-regulated genes involved in both biotic stress responses and hormonal signaling, including auxin-responsive genes. We subsequently observed *ulp1c/d* sensitivity to exogenous auxins. Overall results suggest a ULP1c/d-dependent modulation of auxin in response to *Pst* DC3000.

ULP2a and ULP2b were structural and phylogenetically characterized, revealing a set of common features with ULP2-type proteases of other biological models. T-DNA insertion mutants for this ULP pair were characterized, showing diverse phenotypic defects and constitutive accumulation of SUMO-conjugates. ULP2a and ULP2b displayed unequal redundancy, placing ULP2b as the most preponderant. Microarray analysis evidenced a specific transcriptional signature that suggests the involvement of ULP2s in secondary metabolism, cell wall remodelling and nitrogen assimilation. The *ulp2a/b* mutant also displayed an antagonistic morphological phenotype to that of the well characterized SUMO E3 ligase mutant *siz1*. Most significantly, the triple mutant *ulp2a/b siz1* was phenotypically *siz1*-like, which places ULP2a/b as epistatic and downstream of SIZ1.

The current work highlights the importance of both SIZ1 and various SUMO proteases in development and the plant response to external challenges. Subsequent functional studies will help unravel the fundamental role that SUMO dynamics seems to play at a molecular level, in the control of plant development, hormone regulation, stress response and transcription programming.

## **Análise funcional do sistema de conjugação/desconjugação do SUMO no desenvolvimento e na resposta ao stresse em *Arabidopsis thaliana***

### **RESUMO**

Os organismos vivos estão constantemente sujeitos a alterações ambientais que comprometem a sua sobrevivência. A percepção destas alterações precede eventos de sinalização que culminam na activação de uma resposta adequada. Dado as plantas serem organismos sésseis, elas estão consideravelmente mais expostas a ambientes instáveis, tendo desenvolvido um vasto leque de estratégias para adaptar, evitar ou tolerar o stresse. O conhecimento das bases moleculares subjacentes a estas estratégias é chave para o melhoramento de cultivares de interesse, especialmente estando estes sujeitos a ambientes instáveis. Diversos estudos têm evidenciado como respostas rápidas e reversíveis se correlacionam com mecanismos de modificação pós-tradução (PTM) de proteínas-chave. Mecanismos de reconhecida importância envolvem modificações por pequenos péptidos como a ubiquitina e modificadores semelhantes à ubiquitina (UBLs). O estudo do pequeno modificador relacionado com ubiquitina (*small ubiquitin-related modifier*, SUMO), um membro da família UBL, tem sido crescente nos últimos anos. A ligação do SUMO a uma proteína (sumoilação) poderá exercer diferentes efeitos consoante a proteína-alvo, controlando a sua conformação, e criando, ou bloqueando, interfaces de interacção. Para ser ligado a um alvo, o péptido pre-SUMO terá que ser inicialmente processado por SUMO proteases (família ULP/SENPs), e então conjugado ao resíduo de lisina do alvo através de SUMO E1 activases e SUMO E2 conjugases, auxiliadas por SUMO E3 ligases. A desconjugação do SUMO poderá ser levada a cabo por SUMO proteases. Em plantas, a homeostasia do SUMO é fundamental uma vez que mutações em componentes da via resultam em letalidade embrionária ou em fenótipos pleiotrópicos. Um aspecto interessante do SUMO é o facto dos seus conjugados serem acumulados rapidamente em resposta ao stresse, posicionando o SUMO nos primeiros passos da resposta da planta ao stresse, muito possivelmente associada à reprogramação transcripcional. Estudos da função do SUMO em plantas assentam maioritariamente em estratégias de genética inversa onde é utilizada a espécie modelo *Arabidopsis thaliana* (*Arabidopsis*). De forma semelhante, o presente estudo utilizou mutantes de *Arabidopsis* por inserção de T-DNA para caracterizar o papel da SUMO E3 ligase SIZ1 e das SUMO proteases ULP1c, ULP1d, ULP2a e ULP2b na resposta da planta ao stresse.

As espécies reactivas de oxigénio (ROS) estão descritas como sendo sinalizadores internos originados pelo stresse que podem induzir a acumulação de conjugados de SUMO. No presente trabalho foi abordada a interacção SUMO-ROS, tendo-se obtido evidências de que SIZ1 é essencial para a indução de conjugados de SUMO em resposta a ROS, estando SIZ1 envolvida na tolerância ao stresse oxidativo. Adicionalmente, os mutantes *siz1* revelam alterações na homeostasia das ROS, acumulando constitutivamente peróxido de hidrogénio, ião superóxido e oxigénio singlete. O fenótipo de *siz1* é parcialmente dependente dos níveis de ácido salicílico (SA) uma vez que os defeitos fenotípicos de *siz1* são grandemente recuperados pela expressão do transgene salicilato hidrolase *NahG*. O mutante *siz1* apresenta um fenótipo de auto-imunidade constitutiva, incluindo a acumulação de SA. A análise dos alvos de sumoilação que respondem a stresse oxidativo sugere um papel preponderante de proteínas reguladoras da transcrição na resposta ao stresse oxidativo dependente do SUMO.

A análise bioinformática revelou que o padrão do transcriptoma de *siz1* é semelhante ao de mutantes da via das MAP cinases (MAPK). Estes apresentam também características de resposta auto-imune, como a acumulação de ROS. Os resultados sugerem uma intercepção entre as vias do SUMO e MAPK, porém não foram observadas interações entre MPK4 e péptidos SUMO, ou sumoilação in vitro de MKK2-MPK4. Contudo, as plantas resultantes da introgressão de mutantes *mkk1/2* e *mpk4* com *siz1* apresentam padrões de sumoilação alterados, permitindo sugerir a MPK4 como sendo um regulador negativo da sumoilação, enquanto as MKK1/2 poderão ser epistáticas de SIZ1.

Em contraste com o número limitado de componentes da via de conjugação do SUMO, em plantas existem várias SUMO proteases. As SUMO proteases são uma fonte de selectividade, uma vez que são capazes de discriminar isoformas de SUMOs e discriminar alvos a serem dessumoilados, reconhecem diferencialmente isoformas do SUMO, possuem diferentes padrões de expressão e diferente localização subcelular. A reconstrução filogenética da família ULP de Arabidopsis evidenciou a existência de dois grandes ramos, divididos em quatro subgrupos filogenéticos. Procedemos à caracterização de dois destes subgrupos, um composto por ULP1c/ULP1d e outro por ULP2a/ULP2b. Foi demonstrado que as ULP1c/d funcionam redundantemente na regulação de diversos aspectos do desenvolvimento. Recorrendo a ensaios pelo gene repórter GUS, foi possível estabelecer o padrão de expressão espacial destes genes. Análise por *microarray* de genes diferencialmente expressos no duplo mutante de inserção por T-DNA para ambos genes identificou uma desregulação de genes envolvidos na secura e ácido abscísico. Tendo este resultado em consideração, uma caracterização de fenótipos relacionados com estes processos foi realizada no duplo mutante.

Demonstrou-se igualmente que as ULP1c/d são reguladores negativos na defesa da planta ao microorganismo patogénico hemibiotrófico *Pseudomonas syringae* pv. *tomato* (*Pst*) DC3000. No entanto esta resposta é complexa já que a infecção diminui os níveis de transcritos de *ULP1c* e *ULP1d*, o que poderá estar correlacionado com o aumento de conjugados de SUMO que se seguem à infecção. Uma análise do transcriptoma do *ulp1c/d* após eliciação com *Pst* DC3000 revelou a existência de muitos genes desregulados envolvidos nas respostas ao stresse biótico e sinalização hormonal, incluindo genes de resposta a auxinas. Em suma, os resultados apontam para um papel das ULP1c/d na modelização das auxinas em resposta a *Pst* DC3000.

ULP2a e ULP2b foram caracterizadas estrutural e filogeneticamente, revelando um conjunto de características comuns às proteases do tipo ULP2 de outros modelos biológicos. Mutantes por inserção de T-DNA para este par de ULP mostraram diversos defeitos fenotípicos e uma acumulação constitutiva de conjugados de SUMO. ULP2a e ULP2b apresentam redundância desigual, sendo ULP2b a mais preponderante. A análise por *microarray* evidenciou uma assinatura transcripcional específica que sugere o envolvimento das ULP2s no metabolismo secundário, remodelação da parede celular e assimilação de azoto. O mutante *ulp2a/b* apresenta também um fenótipo morfológico antagónico ao fenótipo bem caracterizado do mutante da SUMO E3 ligase SIZ1. Foi interessante constatar que o triplo mutante *ulp2a/b siz1* é fenotipicamente semelhante a *siz1*, colocando as ULP2a/b numa posição epistática e a jusante de SIZ1.

O presente trabalho salienta a importância tanto de SIZ1 como de várias SUMO proteases na resposta da planta a desafios externos. Estudos funcionais subsequentes permitirão revelar, ao nível molecular, papéis fundamentais da dinâmica do SUMO no controlo do desenvolvimento, da regulação hormonal, da resposta ao stresse e da programação transcripcional.

# TABLE OF CONTENTS

<b>ACKNOWLEDGMENTS</b>	iii
<b>ABSTRACT</b>	vii
<b>RESUMO</b>	ix
<b>TABLE OF CONTENTS</b>	xi
<b>ABBREVIATIONS AND SYMBOLS</b>	xv

## CHAPTER 1

---

### **SUMO, a heavyweight player in plant abiotic stress responses**

<b>1.1. INTRODUCTION</b>	3
<b>1.2. A PRIMER OF THE SUMOYLATION PATHWAY</b>	4
<b>1.3. THE SUMO-ABIOTIC STRESS ASSOCIATION</b>	8
<b>1.4. IDENTIFICATION OF SUMO TARGETS</b>	10
<b>1.5. MOLECULAR BASIS OF SUMO REGULATION OF ABIOTIC STRESS TOLERANCE</b>	14
Extreme temperatures	14
Drought and salt stresses	17
Nutrient imbalance	18
<b>1.6. ADDITIONAL INSIGHTS INTO SUMO FUNCTION AND REGULATION BY STRESS</b>	20
<b>1.7. FINAL CONSIDERATIONS AND FUTURE PERSPECTIVES</b>	22
<b>1.8. OBJECTIVES AND OUTLINE OF THE THESIS</b>	23
<b>1.9. REFERENCES</b>	26

## CHAPTER 2

---

### ***Arabidopsis thaliana* SUMO E3 ligase SIZ1 is a key modulator of reactive oxygen species homeostasis**

<b>2.1. INTRODUCTION</b>	33
<b>2.2. RESULTS</b>	35
Endogenous and exogenous ROS induce SIZ1-dependent sumoylation	35
The <i>siz1</i> mutant displays altered responses to oxidative stress	35
The <i>siz1</i> mutant accumulates superoxide ion and hydrogen peroxide	36
SA levels correlate with ROS homeostatic levels	38
SIZ1 mutant seedlings are not affected in major ROS scavenging enzyme activities	39
Sumoylation interplays with key components of the ROS homeostatic network	40
SIZ1 controls singlet oxygen and chlorophyll levels independently of SA	43
<b>2.3. DISCUSSION</b>	45
ROS positively control the SUMO-conjugate pool	45
SUMO controls ROS homeostatic levels and oxidative stress responses via SIZ1	46
SUMO is likely to interplay with ROS-scavenging mechanisms	47
ROS accumulation involves SA signaling	48
Oxidative stress-dependent SUMO-conjugates suggest interplay between SUMO, ROS and SA at the nuclear level	49
The <i>siz1</i> mutant displays a conditional phenotype	50

Final considerations	51
2.4. MATERIALS AND METHODS	51
2.5. REFERENCES	54

## CHAPTER 3

### Crosstalk between SUMO and MAPK signalling cascades

3.1. INTRODUCTION	61
3.2. RESULTS	63
3.3. DISCUSSION	70
3.4. MATERIALS AND METHODS	73
3.5. REFERENCES	75

## CHAPTER 4

### SUMO proteases ULP1c and ULP1d are required for development and drought stress responses in *Arabidopsis thaliana*

4.1. INTRODUCTION	81
4.2. RESULTS	83
<i>ULP1c</i> and <i>ULP1d</i> show a similar expression pattern	83
ULP1c and ULP1d have a role in plant growth and seed germination	85
Microarray transcript profiling of <i>ulp1c/d</i>	87
ULP1c/d are negative regulators of drought tolerance	88
The <i>ulp1c/d</i> mutant shows altered stomatal response and density	92
The <i>ulp1c/d</i> mutant displays altered SUMO-conjugate levels	93
The triple mutant <i>siz1 ulp1c ulp1d</i> displays an accumulative phenotype	94
4.3. DISCUSSION	95
ULP1c/d control growth and seed germination	96
ULP1c/d affect SUMO conjugation and play a role in drought tolerance	97
ULP1c/d influence responses to low water potential	99
Final considerations	100
4.4. MATERIALS AND METHODS	101
4.5. REFERENCES	104

## CHAPTER 5

### The ULP1c/ULP1d SUMO protease pair negatively regulates *Arabidopsis thaliana* defence against *Pseudomonas syringae* pv. *tomato* DC3000

5.1. INTRODUCTION	111
5.2. RESULTS	114
The double mutant <i>ulp1c/d</i> is less susceptible to <i>Pst</i> DC3000 infection	114
SUMO-conjugate levels are affected by <i>Pst</i> DC3000	116
ULP1c and ULP1d overexpression lines diminish SUMO-conjugates levels but do not display an obvious <i>Pst</i> DC3000 response phenotype	117
Microarray analysis of the <i>ulp1c/d</i> double mutant in response to <i>Pst</i> DC3000	117
Analysis of promoter regions for <i>cis</i> -element enrichment	124
The <i>ulp1c/d</i> mutant displays altered auxin responses	124

<b>5.3. DISCUSSION</b>	126
ULP1c/d are negative regulators of <i>Pst</i> DC3000 resistance	126
ULP1c/d triggers transcriptional reprogramming in response to <i>Pst</i> DC3000	129
ULP1c/d are implicated in the auxin response	130
Identification of ULP1c/d potential targets	131
Future perspectives	131
<b>5.4. MATERIALS AND METHODS</b>	132
<b>5.5. REFERENCES</b>	134

## **CHAPTER 6**

---

### ***Arabidopsis thaliana* ULP2a and ULP2b are SUMO proteases involved in plant development**

<b>6.1. INTRODUCTION</b>	141
<b>6.2. RESULTS</b>	143
Phylogenetic reconstruction and topological analysis of ULP2s	143
ULP2a and ULP2b mutants are developmentally compromised	145
ULP2a and ULP2b have SUMO protease activity	148
ULP2a and ULP2b subcellular localization	149
Microarray transcript profiling of <i>ulp2a/b</i>	149
ULP2 mutants do not recover the <i>siz1</i> phenotype to wild-type	154
<b>6.3. DISCUSSION</b>	156
ULP2a/b are ULPs with likely isopeptidase activity	156
ULP2a/b control plant development downstream of SIZ1	157
ULP2a/b are nuclear components playing a role in transcription regulation	159
<b>6.4. MATERIALS AND METHODS</b>	160
<b>6.5. REFERENCES</b>	163

## **CHAPTER 7**

---

### **Concluding remarks and future perspectives**

<b>7.1. SUMO PROTEASES ARE A SOURCE OF SPECIFICITY</b>	171
<b>7.2. SUMO COMPONENTS ARE ESSENTIAL FOR PLANT GROWTH AND DEVELOPMENT</b>	174
<b>7.3. SUMO CONTROLS PLANT HORMONE HOMEOSTASIS AND HORMONAL RESPONSES</b>	176
<b>7.4. SUMO DYNAMICS IS IMPORTANT FOR AN ADEQUATE RESPONSE TO STRESS</b>	178
<b>7.5. SUMO CONTROLS THE TRANSCRIPTOME BY MODIFYING TRANSCRIPTION REGULATORS</b>	178
<b>7.6. REFERENCES</b>	181

## **APPENDIXES**

---

<b>Appendix I</b>	185
<b>Appendix II</b>	189
<b>Appendix III</b>	193
<b>Appendix IV</b>	197
<b>Appendix V</b>	205
<b>Appendix VI</b>	215



## ABBREVIATIONS AND SYMBOLS

$^1\text{O}_2$	singlet oxygen	LB	lysogeny broth medium
2D	two dimensional	M	molar
3D	three dimensional	m	meter
$^{\circ}\text{C}$	degrees Celsius	MAPK	mitogen activated protein kinase
$\Psi_w$	water potential	MDA	malondialdehyde
ABA	abscisic acid	MEKK	MAPK kinase kinase
ABRE	abscisic acid responsive elements	MES	2-(N-morpholino)ethanesulfonic acid sodium salt
AD	activation domain	min	minute
amiR	artificial microRNA	MJ	methyl jasmonate
APX	ascorbate peroxidase	MKK	MAPK kinase
ATG	autophagy peptide	mol	mole
AUX	auxin	mRNA	messenger RNA
BD	binding domain	MS	Murashige and Skoog culture medium
BL	brassinolide	MV	methyl viologen
bp	base pair	MW	molecular weight
BR	brassinosteroid	NADPH	nicotinamide adenine dinucleotide phosphate
CAT	catalase	<i>NahG</i>	salicylate hydrolase transgene
cDNA	complementary DNA	NBT	nitroblue tetrazolium
CFU	colony forming unit	NDSM	negatively charged amino acid-dependent SUMO motif
CoIP	co-immunoprecipitation	$\text{O}_2^{\cdot-}$	superoxide ion
Col	Colombia-0	OE	over-expression
Da	Dalton	ORF	open reading frame
DAB	3,3-diaminobenzidine	Pa	Pascal
DEG	differentially expressed gene	PAGE	polyacrylamide gel electrophoresis
DNA	deoxyribonucleic acid	PAMP	pathogen-associated molecular pattern
DNase	deoxyribonuclease	PBS	phosphate buffered saline
DPI	days post-inoculation	PBST	phosphate buffered saline tween
E	Einstein	PCA	principal component analysis
EDTA	ethylenediaminetetraacetic acid	PCR	polymerase chain reaction
EFR	EF-Tu receptor	PDSM	phosphorylation dependent SUMO motif
ERF	ethylene response factor	PEG	polyethylene glycol
ET	ethylene	Pi	inorganic phosphate
ETI	effector-triggered immunity	PPI	protein-protein interaction
FLG	flagellin	PPOD	pyrogallol peroxidase
FW	fresh weight	PR	pathogen related gene
g	gram	PRX	peroxidase
<i>g</i>	relative centrifuge force	PS	photosystem
GFP	green fluorescent protein	<i>Pst</i>	<i>Pseudomonas syringae</i> pv. <i>tomato</i>
GO	gene ontology	PTI	PAMP-triggered immunity
GUS	beta-glucuronidase	PTM	post-translation modification
h	hour	qPCR	quantitative real-time PCR
$\text{H}_2\text{O}_2$	hydrogen peroxide	R-protein	resistance protein
HL	high-light	RBOH	respiratory burst oxidase homologue
HMWC	high molecular weight SUMO conjugate	RGN	ROS gene network
$\text{HO}\cdot$	hydroxyl radical	RNA	ribonucleic acid
HS	heat shock	RNase	ribonuclease
HSF	heat shock factor	ROS	reactive oxygen species
HSP	heat shock protein	RT	room temperature
IgG	immunoglobulin G	RUB	related to ubiquitin
IAA	indole-3-acetic acid	s	second
JA	jasmonate		
KIU	Kaonashi ULP-like		
L	Litre		

SA	salicylic acid	TBA	2-thiobarbituric acid
SAE	SUMO activating enzyme	TF	transcription factor
SAUR	small auxin up-regulated	T <sub>m</sub>	melting temperature
SAR	systemic acquired resistance	Ub	ubiquitin
SCE	SUMO conjugating enzyme	UBL	ubiquitin-like modifier
SDS	sodium dodecyl sulphate	ULP	ubiquitin-like protease
SEM	standard error of the means	UPS	ubiquitin proteasome system
SENP	sentrin-specific protease	UTR	untranslated region
SIM	SUMO interacting motif	V	volt
SOD	superoxide dismutase	v/v	volume per volume
SOSG	singlet oxygen sensor green	w/v	weight per volume
STUbL	SUMO targeted ubiquitin ligase	Wt	wild-type
SUMO	small ubiquitin-like modifier	XTH	xyloglucan endotransglucosylase/ hydrolase
T3SS	type III secretion system	Y2H	yeast two-hybrid
T-DNA	transfer DNA		
<i>Taq</i>	<i>Thermus aquaticus</i> polymerase		

### Amino acids

A	Ala	Alanine
C	Cys	Cysteine
D	Asp	Aspartate
E	Glu	Glutamate
F	Phe	Phenylalanine
G	Gly	Glycine
H	His	Histidine
I	Ile	Isoleucine
K	Lys	Lysine
L	Leu	Leucine
M	Met	Methionine
N	Asn	Asparagine
P	Pro	Proline
Q	Gln	Glutamine
R	Arg	Arginine
S	Ser	Serine
T	Thr	Threonine
V	Val	Valine
W	Trp	Tryptophane
Y	Tyr	Tyrosine
X		unspecific amino acid
ψ		large hydrophobic residue

### Nucleotides

A	Adenine	
C	Cytosine	
G	Guanine	
T	Thymine	
U	Uracil	
R	A or G	Purine
Y	C or T	Pyrimidine
W	A or T	
S	C or G	
M	A or C	
K	G or T	
B	C, G or T not A	
D	A, G or T not C	
H	A, C or T not G	
V	A, C or G not T	
N	A, C, G or T	
AMP	Adenosine monophosphate	
ATP	Adenosine-5'-triphosphate	
dATP	2'-deoxyadenosine-5'-triphosphate	
dCTP	2'-deoxycytidine-5'-triphosphate	
dGTP	2'-deoxyguanosine-5'-triphosphate	
dNTP	2'-deoxynucleotide-5'-triphosphate	
dTTP	2'-deoxythymidine-5'-triphosphate	
GDP	Guanosine-5'-diphosphate	
GTP	Guanosine-5'-triphosphate	