

Factors that affect physicochemical and acid-base properties of compost and its potential use as a soil amendment

R. López^{1,*}, J. Antelo¹, A.C. Silva^{1,2}, F. Bento², S. Fiol^{1,3}

¹CRETUS-Institute. Cross-disciplinary Research in Environmental Technologies
University of Santiago de Compostela, 15782, Santiago de Compostela, Spain

²Department of Chemistry, Center of Chemistry, University of Minho, Campus Gualtar,
4710-057 Braga, Portugal

³Department of Physical Chemistry. University of Santiago de Compostela, 15782,
Santiago de Compostela, Spain

*Corresponding author

e-mail address: rociolopez.monforte@gmail.com

Abstract

Composting and vermicomposting have attracted attention in relation to both waste management and the potential to produce organic amendments that could improve soil quality. The main differences between compost depend on the feedstock, the production process, and the degree of maturity. In the present study, samples of compost of different origin or subjected to different composting methods were collected for analysis. Additionally, samples collected at various stages of the composting process were compared. Different analysis and techniques were used to characterize the compost samples and the organic matter extracts. The correlations obtained would allow for predicting the compost behaviour based on certain characteristics, and a reduction in the number of parameters determined experimentally, thus facilitating comparisons between different compost. In addition, the potential value of the compost as amendment was tested with a Haplic Cambisol from a mining area. The application of compost increased the pH, the organic matter and nutrient content, and promoted seed germination and root growth.

Keywords: Compost, feedstock, organic waste, humic substances, soil amendment

1. Introduction

Waste management is required to maintain natural resources and to protect the environment. Recycling, reuse and valorization of waste are possible solutions potentiating the circular economy (Hollins et al., 2017). Although most of the waste generated by anthropogenic activities is biodegradable, it should not be directly applied because its uncontrolled decomposition can have negative effects on soil systems (Hanc et al., 2019a). Organic matter from waste can be decomposed and stabilized in a process in which microorganisms play an important role. Thus, biodegradable organic residues can be stabilized and repurposed via composting and vermicomposting processes. Both processes are aerobic and bio-oxidative but differ in the involvement of earthworms in vermicomposting (Fornes et al., 2012). The involvement of earthworms in the composting process promotes the aeration and fragmentation of the organic matter, thus increasing the rate of mineralization (Hanc and Vasak, 2015)

The use of compost and vermicompost to improve soil quality has increased in recent years. Numerous studies have investigated the use of compost as a soil amendment with the aim of immobilising potentially toxic elements or increasing nutrient or organic matter content (Forjan et al., 2017; Palansooriya et al., 2020). To use the compost as a soil amendment it should have reached a certain degree of maturity and stability. Maturity is associated with the capacity of a compost to favour plant growth and with the absence of phytotoxic compounds, whereas stability is associated with a decrease in microbial activity during composting (Cerdá et al., 2018). The stability is related to the transformation of fresh organic matter into recalcitrant aliphatic compounds and highly humified aromatic structures (Droussi et al., 2009). Different physical, chemical and biological parameters can be used to determine the degree of maturity and stability of compost, including odour, cation exchange capacity, nitrogen and carbon contents or degree of humification (Bernal et al., 1998; Droussi et al., 2009).

The degree of humification of organic matter is related to the production of humic substances and to the degree of maturity or stability reached during composting (Adani et al., 1999). As the aim is to yield a stable product rich in humic substances, these substances are often extracted from compost for characterization (Amir et al., 2003). Some studies indicate that the humic substances extracted from compost improve the soil properties more effectively than other more conventional amendments (Spaccini and Piccolo, 2009). Humic substances can benefit soil systems by regulating pH, acting as redox agents, complexing metallic ions, and stimulating plant growth (Fuentes et al., 2018; Guo et al., 2019). Thus, humic substances show great potential for elaborating biofertilizers and for removing heavy metals from soils. The contribution of humic substances to the fate and bioavailability of macro and micronutrients and toxic species, such as heavy metals and pesticides, can be attributed to the presence of acidic functional groups in their structure (Ritchie and Perdue, 2008). The acid-base properties of HS-like substances and dissolved organic matter can provide useful information to improve the performance of the compost for agronomic purposes in soils (Plaza et al., 2005).

Various studies have focused on analysing compost and vermicompost derived from animal and plant waste (Bernal et al., 2009; Kucbel et al., 2019; Lv et al., 2013). Other studies have compared the composting systems (Soobhany et al., 2017), while others have characterized the HS extracted (Silva et al., 2013), monitored the changes in organic matter throughout the composting process (Jouraiphy et al., 2005) and, eventually, evaluated the effects of incorporating the resulting material on soils (Forjan et al., 2017; Teodoro et al., 2020). Although there are multiple studies of compost and

vermicompost from different feedstock, of the fate of the organic matter during the composting process, and even of the humic substances present in the composted materials, it is not easy to find studies comprising all these aspects and establishing comparisons among them.

The aim of the present research was to conduct an integral study of the aforementioned aspects, in order to obtain more detailed information about the products generated by composting and vermicomposting. For this purpose, samples of compost and vermicompost were selected to compare both the effect of the composting process and the effect of the feedstock material. The evolution of the organic matter was also monitored at different stages of the composting process. Although soil-compost interactions mainly occur in aqueous media, information about humic-like substances and dissolved organic carbon of these types of material is scarce. We therefore decided to compare the composition and acid-base behaviour of the dissolved organic carbon in the equilibrium solution with the corresponding humic-like extracts. Finally, with the aim of evaluating the effects of the application of stabilized organic residues on degraded soils, compost and vermicompost samples were applied as amendments to an unproductive mine soil.

2. Materials and Methods

2.1. Samples

Six compost samples were analysed: four were commercial products (CLW (livestock waste), CUW (urban waste), CSS (sewage sludge) and CVA (algae) and two were produced in traditional composting or vermicomposting systems (CDDW, from domestic waste and CVDW, from domestic waste using earthworms). In addition, with the aim of studying the changes in the organic matter throughout the composting process, samples were obtained at different stages (UW0 (fresh urban waste), CUW15 (after 15 days composting) and CUW30 (after 30 days of composting)), as outlined in Figure S1 (Supporting Information). A detailed description of the samples is included in Table S1 (Supporting Information).

The compost samples were used to extract the dissolved organic carbon (equilibrium solutions) by adding 150 mL of water to 7.5 g of compost. The resulting suspension was homogenized by an initial vigorous shaking and then equilibrated for 5 days (open atmosphere). The solution was then separated by centrifugation at 6000 rpm.

The humified organic matter was extracted from compost to yield the humic substance-like (HS-like) fractions. The extraction was carried out following the protocol established by the International Humic Substances Society (IHSS) (Swift, 1996).

2.2. Characterization

The compost, equilibrium solutions and HS-like fractions were characterized using common methods for these types of materials. Briefly, C, H, N and S contents were determined in an element analyzer (TruSpec CHN-1000, LecoSC-144DR); ash content was determined gravimetrically by combustion at 650 °C; oxygen content was determined as $O = 100 - (C + N + H + S + \text{ash})$; oxidizable C was determined by Sauerlandt's method (Walkley, 1947) and C in solution was determined using a total organic carbon analyzer (TOC-L CSN Shimadzu); concentrations of the major elements were determined by ICP-OES (Perkin-Elmer Optima 3300DV); P concentration was determined in the compost

and in the equilibrium solution by the molybdenum blue method (Murphy and Riley, 1962); concentration of ammonium in the equilibrium solution was determined with a selective electrode (Elit electrode Nico 2000); pH and electrical conductivity were measured in the equilibrium solution with a combined electrode (Crison, pH 0-14) and a conductivity meter (Crison GLP 32), respectively.

2.3. Acid-base properties

Acid-base titrations were conducted with the equilibrium solutions (~ 5 g solid/L) and of the different HS-like fractions (0.1 g HS/L), following the protocol proposed by López et al. (2003). The titration of the equilibrium solutions was conducted at 0.1 M ionic strength in KNO₃. The titration of the HS-like fractions was also conducted at 0.01 M to obtain the intrinsic parameters of protonation. Because of the high pH of the equilibrium solutions, two aliquots were taken from the same sample to study protonation (titration with 0.1M HCl) and deprotonation (titration with 0.1 M KOH).

2.4. Application of the amendments

Compost samples were applied to an unproductive mine soil, with the aim of evaluating the efficacy of these samples as soil amendments. The soil was a Haplic Cambisol (Ch2) collected in a uranium mine site (Mina Fé, Saelices el Chico, Spain) and characterized by Arán et al. (2020). The compost samples were applied as individual amendments by manually mixing the soil and compost. Thus, 150-200 g of soil were amended with the amount of compost required to reach a variable content between 2% and 10% (w:w). The soil-compost samples were then incubated at room temperature for 2-3 weeks, during which the moisture content was maintained at between 50 and 70% of the water holding capacity. At the end of the incubation period, the samples were dried, and the pH, C and N contents, total P, Olsen P, and the exchange cations were determined by standard methods (Sparks et al., 1996). A sample of unamended soil was used as a control.

Aqueous equilibrium solutions were made from the amended samples by the same method as outlined above. The resulting equilibrium solutions were used in germination assays on filter paper, following a similar protocol to that described by Di Salvatore et al. (2008). This assay provides information about the maturity and potential phytotoxicity of compost or other types of biomaterial (Roe et al., 1997). The ryegrass *Lolium perenne* was selected for use in the assays since it is relatively insensitive to high concentrations of metal, as usually observed in mine soils (Teodoro et al., 2020). Twenty *Lolium perenne* seeds were placed on three layers of filter paper held in Petri dishes (120 x 20 mm) and watered with 8 mL of equilibrium solution or distilled water (control). The Petri dishes were then sealed and placed in a germination chamber, where they were held at 22 °C, in darkness, for 7 days. Then, the germinated seeds were counted, and the root length was measured. Seeds were considered to have germinated when the root was at least 1 mm long. Each treatment was carried out in triplicate. To facilitate comparison of the results of the germination assays, the percentages of relative radicle growth (RRG), relative seed germination (RSG) and germination index (GI) was calculated (Walter et al., 2006). The normalized residual elongation (NRE) index was also determined (Bagur-González et al., 2011):

$$RRG(\%) = \frac{\text{Mean root length in sample}}{\text{Mean root length in control}} \times 100 \quad (1)$$

$$RSG(\%) = \frac{\text{Number of seed germinated in sample}}{\text{Number of seed germinated in control}} \times 100 \quad (2)$$

$$GI(\%) = \frac{RRG \times RSG}{100} \quad (3)$$

$$NRE = \frac{\text{Mean root length in sample} - \text{Mean root length in control}}{\text{Mean root length in control}} \quad (4)$$

3. Results and discussion

The results and discussion of physicochemical characterization, acid-base properties and application of the compost as soil amendment are presented in different sections. Within each section, different sub-sections are devoted to the results from compost, equilibrium solution and HS-like fractions, respectively. In each sub-section, results are displayed into three discussion blocks:

- Origin: Samples of different feedstock subjected to a similar composting process: CLW, CUW, CDDW and CSS; CVA and CVDW
- Process: Samples of the same feedstock subjected to different composting processes: CDDW and CVDW
- Maturation: Samples obtained at different times throughout the composting process: UW0, CUW15, CUW30 and CUW

3.1. Characterization

3.1.1. Compost

The results of the chemical analysis are included, along with the values of other relevant parameters, in Table 1. The metal contents of the different compost are shown in Table S2 (Supporting Information). The percentage of carbon in the samples produced by a traditional composting process with different types of waste material varied between 22.8 and 34.0%. The sample with the highest content of carbon corresponded to that obtained from a combination of green and food waste (CUW). The sample with the lowest percentage of carbon was produced from livestock manure (CLW). Comparison of samples CDDW and CVDW, produced from similar feedstock and subjected to traditional composting or vermicomposting, respectively, revealed a substantial increase in both the total and oxidizable carbon due to the vermicomposting process. The vermicomposting process reduces the proportion of carbon that can be oxidised relative to traditional composting: 75.4% and 81.0%, respectively. Analysis of the changes in the carbon content during the composting process (UW0, CUW15, CUW30 and CUW) revealed a decrease in both the total and oxidizable carbon, particularly during the maturation stage, i.e. oxidizable carbon was 86.1 % of the total carbon at the end of the process, while it remained above 95% during the initial decomposition stage.

The C/N ratio (Table 1) is widely used as an indicator of compost maturity. Thus, various studies report that values below 20 indicate that the maturity of the compost is acceptable, but only those compost showing a C/N ratio below 15 are recommended for

agronomic use (Soobhany et al., 2017). The C/N ratio ranged between 8.6 (CSS) and 12.8 (CLW) for the samples produced from different types of feedstock. A value of 8.5 was found for sample CVDW, lower than that determined for sample CDDW (10.3) produced by a different composting process. This finding appears to be consistent with previous studies and can be attributed to the earthworms, which supply N and may mineralize the organic nitrogen in the form of mucus excretions and nitrogenous excrements during vermicomposting (Kaushik and Garg, 2003; Suthar, 2010). As the composting process advanced, the values of the C/N ratio decreased due to the loss of carbon. As indicated by Bernal et al. (1998), the greatest decrease occurred during the active phase, while the maturation stage scarcely influenced the ratio. Although the C/N ratio in sample UW0 is low relative to the values suggested by various authors as optimal for a composting process to be considered effective ($20 < C/N < 30$) (Kumar et al., 2010; Wang et al., 2019), they show that green and food waste, for which the initial values of this ratio are low, may be effectively broken down by the composting process.

The cation exchange capacity (CEC) has been widely used to evaluate the maturity of the compost and the degree of humification of the organic matter, as functional groups that increased the CEC are generated during this process. Thus, high CEC values may be related to the increase of humic substances in the compost. Generally, values of CEC > 60 cmol $^+$ /kg indicate that the maturity of the compost is adequate for agronomic use (Harada and Inoko, 1980). Sample CSS, derived from sewage sludge, was the only sample showing a value below this reference. Results obtained showed that the CEC is positively correlated with the C/N ratio when compost of different origin were compared (Figure 1). Thus, samples with the highest C/N ratio, CLW and CUW, showed larger CEC values. The CEC increased during the composing process, as observed by comparison of the uncomposted material (UW0) and the final product (CUW) (Table 1). After the first cycle of composting, the CEC decreased (comparing UW0 and CUW15) and then increased gradually with composting time. This behaviour has been previously described by Harada and Inoko (1980) for urban-waste compost. A negative correlation between CEC and C/N ratio has been found with composting time (or composting maturity), similar to that found by Harada and Inoko (1980), although in their study the range for the C/N ratio was larger. This trend appears to indicate that after the first composting cycle, the carbon is mainly derived from humified organic matter, in which the acid sites are occupied by exchange cations. As the composting process advances, the formation of humic substances is favoured, increasing the content of carboxylic and phenolic groups.

3.1.2. *Equilibrium solutions*

The values of the parameters used to characterize the equilibrium solutions of the samples are given in Table 2. The concentrations of the metals determined in the equilibrium solutions are included in Table S3 (Supporting Information). The pH of the equilibrium solutions ranged between 6.3 and 8.5. The lowest values corresponded to the two vermicompost samples. Comparison of the compost and vermicompost samples made from the same feedstock showed that the pH of the equilibrium solution was lower after vermicomposting. During the composting process, the pH increased slightly in the first cycle and then increased sharply, by approximately one unit, in the second cycle, before remaining almost constant throughout the maturation stage.

The electrical conductivity (EC) was used as an indicator of salinity, in order to evaluate any potential problems on using the compost as soil amendment. The EC should remain below 4 mS/cm to be tolerated by plants (Lasaridi et al., 2006). This value was only exceeded in sample CLW, due to high concentrations of sodium and potassium. On

the other hand, this parameter did not follow any clear pattern throughout the composting process. However, the EC was much lower in the commercial product than in the uncomposted sample, and the largest decrease occurred during the first composting cycle. The pH and EC were correlated in mature compost samples, which may be explained by differences in pH-dependent occupation of acid sites in the organic matter by cations (Figure 2). The relationship was again different in the composted and vermicomposted samples. The pH of sample CSS was lower than expected from the EC, which can be explained by the high concentration of NH_4^+ in this sample (159.1 mg/L) relative to the other samples (between 4.6 and 21.2 mg/L). The high concentration of NH_4^+ in sample CSS may indicate that the compost was not mature (van der Wurff et al., 2016).

The percentages of the major elements that mobilize into the equilibrium solution are shown in Figure S2 (Supporting Information). The highest percentages generally corresponded to Na and K, except in sample CSS, in which Na, K and Mg occurred in almost the same proportion (~20%). The presence of a high concentration of Na in solution may be problematic regarding the use of compost as a soil amendment as this cation is rapidly absorbed by plants, thus preventing absorption of Ca and Mg (Kucbel et al., 2019). Comparison of the compost and vermicompost produced from the same waste material (CDDW and CVDW) revealed very similar results regarding the percentages of the major elements released to the solution, suggesting that the feedstock is the main factor responsible for the availability of the major elements. In general, the major cations tended to decrease in the equilibrium solution of the samples collected at the different stages of the composting process.

The sample with the highest concentration of dissolved organic carbon was that obtained from livestock waste (CLW), even though the carbon content of the compost was low. The percentage of oxidizable carbon was lowest in the solid sample of this compost, which led to the percentage of dissolved organic carbon being the highest. The equilibrium solution samples with the lowest content of dissolved organic carbon were CDDW and CVA. Comparison of the uncomposted sample (UW0) and the final composted sample (CUW) revealed a marked decrease in the dissolved organic carbon. Very similar findings have been reported by Huncce et al. (2020). The decrease is due to the fact that the water-soluble carbon is most readily degradable. The organic matter in the equilibrium solution of the uncomposted material is mainly composed of sugars, hemicellulose, phenols, organic acids and amino acids. However, the organic matter varies as the composting process advances, becoming enriched in HS-like components, which are resistant to degradation.

Semi-empirical relationships between parameters determined by UV-visible spectroscopy are often used to characterize organic matter in solution. These include the correlation between the coefficient of molar absorptivity at 280 nm (ϵ_{280}) and the percentage aromaticity ($\% \text{ aromaticity} = 0.05 \times \epsilon_{280} + 6.74$) and between this coefficient and the molar mass ($\text{molar mass} = 3.99 \times \epsilon_{280} + 490$) (Fuentes et al., 2006). The lowest values of ϵ_{280} corresponded to the uncomposted sample UW0 and sample CSS (Table S4, Supporting Information), which is representative of an immature compost. The highest value corresponded to sample CDDW, followed by the two vermicompost samples (CVA and CVDW). In the equilibrium solutions of the mature samples, except that of sample CSS, the percentage aromaticity was between 20 and 27 %. The parameters obtained with the semi-empirical relationships increased clearly throughout the composting process, which suggest that these are good indicators of the degree of humification of the organic matter in solution.

3.1.3. HS-like fractions

As in other studies with compost (Fuentes et al., 2018; Veecken et al., 2000), the protocol used for extracting the HS-like components is that recommended by the IHSS for extracting humic substances from soil. This protocol yields good results when applied to mature organic matter but is probably not as effective for immature organic matter. It is therefore possible that the method will quantify carbohydrates, proteins and amino acids as HS-like compounds, even though these are precursors of HS-like compounds and have not yet decomposed.

The percentages of major elements and the corresponding atomic ratios for the HS-like fractions extracted are shown in Table 3. Analysis of the extraction yields of HS-like components reveals that sample CLW had the highest content of HS-like compounds (42 g/kg of compost). The total carbon content was lowest in this compost, but the DOC was highest. Analysis of these data revealed a correlation between the amount of HS-like compounds extracted from the compost of different origin and the percentage of organic carbon in the equilibrium solution (Figure 3 a). Inclusion of the other samples in the same plot shows that the concentration of DOC present in the equilibrium solution provides a good estimate of the content of HS-like compounds. The only exception is sample CVA, for which a lower DOC was measured. Although organic matter humification should occur as the composting process advances, the extraction yield of HS-like compounds was very high for the fresh material at the beginning of the composting process. This can be explained by considering that the protocol used extracts both the HS-like compounds and their precursors, which are also present in the fresh material. After the first composting cycle, the percentage of HS-like compounds extracted increased and then remained almost constant throughout the maturation stage. A similar effect was described by Wang et al. (2019), although in most studies in which this topic is addressed the amount of HS-like compounds increases throughout the composting process (Amir et al., 2010). In addition to the previously mentioned correlation, the relationship between the yield of HS-like compounds and the pH of the equilibrium solution distinguished the two types of process: composting and vermicomposting (Figure 3 b).

Analysis of the HS-like fractions revealed a greater carbon content in the humic acid (HA) than in the fulvic acid (FA) fraction, whereas the latter is richer in oxygen. Regarding the content of other elements, generally the proportion of nitrogen was much higher in the HA extracted from mature compost. Comparison of the HS-like compounds extracted from compost and vermicompost from the same feedstock showed that the nitrogen content was higher in the vermicompost, which can be attributed to the fact that the solid sample also contained more nitrogen due to the direct action of earthworms. The high nitrogen contents in the FA fraction extracted from samples CSS and UW0 may indicate the presence of non-humified biomolecules (such as polysaccharides and polypeptides) or the incomplete hydrolysis of protein components (Sánchez-Monedero et al., 2002), again indicating the immaturity of both samples.

The atomic ratios in the FA and HA fractions obtained from the mature compost were plotted in a van Krevelen diagram, along with the ratios reported in other studies (Figure S3, Supporting Information). We can see that the samples are more widely dispersed in the FA fraction than in the HA fraction. Generally, the HAs are displaced towards lower values of the O/C ratio. This was expected as HA has higher carbon content and lower oxygen content than FA. This is usually attributed to the enrichment in oxygenated acidic groups for the FA fraction, and is therefore associated with the number of acidic sites, which is expected to be higher in this fraction. In general, the H/C ratios

are similar in both fractions, except in sample CUW. The variation in the atomic ratios during the different stages of composting is similar to that reported in previous studies (Amir et al., 2010; Baddi et al., 2004; Hanc et al., 2019b). Both fractions are displaced towards higher values of the O/C atomic ratio, which may be related to an increase in acidic groups throughout the composting process due to the gradual transformation of polysaccharides and other similar compounds into oxygenated compounds such as carboxylic and phenolic compounds (Hanc et al., 2019b). The H/C ratios remain almost constant, as previously described by different authors (Baddi et al., 2004; Riffaldi et al., 1983), and therefore will not indicate the level of decomposition of the organic matter. Comparison of the HS-like fractions extracted from the compost of different origin shows that the aromaticity is generally higher in the HA (Figure S4, Supporting Information).

3.2. Acid-base properties

3.2.1. Equilibrium solutions

Acid-base titrations were conducted to evaluate the concentration and type of acid groups present in the dissolved organic matter. The charge curve for sample CLW, resulting from the difference in the mmoles of acid or base consumed in the titration of the sample and those consumed by the electrolyte is shown in Figure 4 a. The figure includes the derivative obtained from the charge curve, in which three well-defined peaks were observed. The equilibrium solutions of all samples, except CSS, showed similar behaviour independently of the composting process and the feedstock material (Figure 4 b and Figure S5, Supporting Information). In almost all cases, the maximum values in the curves coincide at the same pKa values.

Despite the scarcity of information about the acid-base properties of DOC obtained from compost, Sciubba et al. (2015) also identified the presence of three types of acid positions and attributed these to carboxylic groups in amino acids (pKa between 2 and 3), carboxylic groups in organic acids (pKa between 4 and 6) and phenolic hydroxyl groups, hydroxyl in carbohydrate units and amino groups from amino acids (pKa between 8 and 11), respectively. The pH interval considered in the present study only enabled observation of positions type 2 and type 3 identified by Sciubba et al. (2015).

The 3-site Langmuir isotherm was used to obtain the acid-base parameters of the dissolved organic matter (equation 5):

$$Q = Q_0 + \frac{M_1}{1 + K_1[H^+]} + \frac{M_2}{1 + K_2[H^+]} + \frac{M_3}{1 + K_3[H^+]} \quad (5)$$

where K_i represents the protonation constant of the different groups, M_i represents the abundance of each type of group, and Q_0 is the charge due to occupation of the acid sites by cations that must be displaced by protonation. The model fitting results are shown in Table 4. For correct interpretation it should be noted that the charge generated on the organic matter is due to the deprotonation of acid sites and it is therefore negative. The Q_0 value shown in Table 4 can be interpreted as a net positive charge that is occupying acid sites, as positive charges are usually used to facilitate data handling.

The dissolved organic matter (DOM) containing the highest and lowest number of acid sites (M_T), estimated by measuring the consumption of acid or base, is that corresponding to samples CSS and CVA, respectively. The presence of some cations in

the solution may alter these results, as we attributed the consumption of acid/base to the protonation/deprotonation of the acid sites in the organic matter, without considering possible hydrolysis reactions and/or precipitation of cations. The possible effect of hydrolysis reactions may be important in samples such as CSS, due to the high concentration of NH_4^+ (Table 2). In all cases, the most acidic groups are the least abundant, whereas the abundance of the other two types of groups was very similar (Table 4). Again, sample CSS is the exception.

The log K values (Table 4) did not follow any trend in relation to the origin of the compost. However, in the vermicompost samples, the DOM contains more acid groups when the original material contains a high proportion of manure (i.e. sample CVA). The result of applying the different composting processes to similar types of material, as in samples CDDW and CVDW, demonstrates that vermicomposting generates DOM with higher content of functional groups, although these groups show a less acidic character. Sample CSS was once again an exception, as it yielded the highest log K value for the first type of sites (4.24 relative to a mean value of 3.43 ± 0.27 for the other samples) and the lowest values for sites 2 and 3 (6.27 and 9.65 relative to mean values for the other samples of 6.54 ± 0.08 and 10.17 ± 0.13 , respectively).

The same type of comparison can be carried out for the DOM in the equilibrium solution of the samples obtained at the different stages of composting (Figure S6, Supporting Information). There were notable differences in the curves of the fresh material and the material collected at the different stages of the process. This difference was also apparent in the acidity constants. The high abundance of the first type of sites in sample UW0 is attributed to the contribution of amino acids, which disappear as the organic matter matures. This, along with the fact that the DOM of sample CSS has the highest number of this type of site, appears to reinforce the idea of the incomplete degradation of the organic matter in the composting process of CSS.

The model described by equation 5 was also fitted to the charge curves of the DOM of the samples obtained throughout the composting process. The results (Table 4) show a clear decrease in the number of acid sites during the composting process, indicating notable differences between DOM at the beginning of the process, when non humified organic matter is also mobilized into the solution, and that after the first composting cycle. The most acidic groups were the most abundant in UW0 (almost 50% of the total), whereas they became the least abundant after the first composting cycle. The variation in the acidity of the groups as the process advanced indicated a clear difference between UW0 and the other samples, as it corresponded to the highest log K value for the first site (4.50 relative to a mean value for the other samples, of 3.75 ± 0.22) and lower values for sites 2 and 3 (6.05 and 9.73 compared with 6.57 ± 0.08 and 10.08 ± 0.03 , respectively).

3.2.2. *HS-like fractions*

As expected, the FA charge was larger than the HA charge over the entire pH range (Figure 5 and Figure S7, Supporting Information). The distribution of affinities for the HS-like fractions indicates the presence of two types of acid sites, while in the case of DOM three different acid sites were obtained. This indicates the existence of two types of acid sites, attributed to carboxylic and phenolic groups (Milne et al., 2001; Plaza et al., 2005). The charge generated due to the deprotonation of the acid groups can be described by equation 6:

$$Q = Q_0 + \frac{M_1}{1 + (K_1 [H^+])^{m_1}} + \frac{M_2}{1 + (K_2 [H^+])^{m_2}} \quad (6)$$

where K_i is the protonation constant of the carboxylic and phenolic groups, M_i is the abundance of each type of acid site, and m_i represents the width of the distributions.

The parameters obtained by fitting equation 6 are shown in Table S5 (Supporting Information). The Q_0 values are not shown as they are equal or close to zero. The results show that the bimodal model provided a good fit to the experimental data and demonstrate that the total number of acid sites (M_T) is greater in the FA fraction. The ratio between the total number of sites in both fractions ($M_T(\text{FA})/ M_T(\text{HA})$) was around 1.1 for the HS-like extracted from the compost samples, with the exception of sample CSS, which has a value slightly higher than 2, and around 1.5 for the corresponding extracts from vermicompost.

To enable comparison with the abundance of the acid sites in the DOM, the site abundance of the HS-like fractions is expressed in mmol of charge per g of compost, by using the extraction yield (Table 3). Considering the different solubility of the HS-like extracts, the DOM present in the equilibrium solution will be mainly constituted by the FA fraction. The comparison (Figure S8, Supporting Information) revealed similar trends, although the sites are more abundant in the DOM. This difference can be attributed to the low extraction yield for the HS-like fraction and to the presence in the equilibrium solution of non-humified organic matter and cations (hydrolysis of which may contribute to overestimation of acid sites in the DOM). The differences between the charge curves for the DOM and the HS-like extracts complicate comparisons. However, the mean values of the bimodal distributions that characterize the FA fraction are very similar to the pKa values of type 1 and 3 sites in the DOM (Table 4 and Table S5, Supporting Information). In all samples, the log K values of the most acidic sites in the DOM are lower than those obtained for the FA fraction. By contrast, the log K values for the weakest sites are lower in the FA fraction than in the DOM in the equilibrium solutions. Some of these differences could be attributed to the methodology followed for the potentiometric titrations (Figure S9, Supporting Information).

The charge curves of the HS-like fractions extracted from the samples collected throughout the composting process revealed clear differences in the FA and HA fractions (Figure 6). Regarding the FA fraction, a drastic change in the charge curves was observed between UW0 and CUW15. In both samples, the initial charges were similar, but then differed as the pH increased until reaching a value of 6, after which the values increased in parallel. After the first composting cycle, the differences were only observed at low pH values. By contrast, the HA fraction underwent a gradual increase of charge throughout the pH range as the composting process advanced, along with an increase in the total number of acid sites (Table S5).

A rigorous study of the acid-base properties of HS-like extracts requires determining the intrinsic protonation constants and considering the electrostatic effect of the organic polyelectrolytes. The Donnan model (Kinniburgh et al., 1996) was used in this work, which considers HS to form an electrically neutral gel-like phase. The Donnan volume (V_D) is related to the ionic strength through equation 7.

$$V_D = b (1 - \log \log I) - 1 \quad (7)$$

where b is an empiric parameter characteristic of a particular substance and describes V_D variation with the ionic strength. The optimization of b allows for calculating the electrostatic potential in the Donnan phase and the intrinsic protonation parameters (Table 5). These parameters can be compared to the generic parameters for aquatic and terrestrial FA and HA obtained by Milne et al. (2001) and to the intrinsic parameters specifically obtained for compost derived HS-like (Table S6, Supporting Information). The average values obtained for the parameter b , 0.62 for FA-like and 0.45 for HA-like, are higher for the FA-like fraction, similarly to the values reported by Milne et al. (2001) and by Plaza et al. (2005). An exception to this trend has been reported by Fernández et al. (2007) and by van Zomeren et al. (2009), who found very similar values for this parameter in the fulvic and humic fractions extracted from compost (Table S6, Supporting Information). Analysis of the results obtained for the HS-like fractions extracted at different stages of the composting process indicates that b decreases throughout the maturation process. The same variation was reported by Plaza et al. (2005). This trend is in agreement with the negative correlation that Christl and Kretzschmar (2001) established between the parameter b and the particle size and molecular weight. The higher b values observed for the HS-like in the first stages of composting suggest a high contribution of small molecular weight particles. This situation reverses as HS-like condensation occurs during the composting process.

The average content of total acid sites, M_T , obtained for all the samples, 5.58 and 4.00 mmol/g for FA-like and for HA-like, respectively, was small compared to the literature data (Table S6, Supporting Information). However, when the abundance of carboxylic and phenolic groups is expressed as a percentage of the total content of acid groups, the results are almost identical to the percentages reported by Plaza et al. (2005), with carboxylic groups representing approximately 65% of the total acid groups in both FA and HA fractions. Compared to generic aquatic and terrestrial HS (Milne et al., 2001) and to other HS-like derived from compost (Fernández et al. 2007, van Zomeren et al., 2009), our samples have lower percentage of carboxylic groups in both HS-like fractions. The distribution of acid groups, between carboxylic and phenolic type, in HS-like extracted during the maturation process is not very different from the distribution observed when comparing HS-like from different feedstock. An increase in the total content of acid groups, as opposed to the DOM of the equilibrium solutions, was observed during the composting process, which is in agreement with the behaviour displayed by the charge curves of the HS-like extracts (Figure 6). As previously reported by Plaza et al. (2005), a loss of aliphatic structures and carbohydrates, along with an increase in acid functional groups and groups containing N and S occur during the composting process. This increase in the site abundance was more evident in the HA-like extracts, from 3.57 to 4.74 mmol/g, than in the FA-like extracts, from 5.00 to 5.11 mmol/g, and it is reflected in the separation that exists between the charge curves with the composting time (Figure 6). The charge curves for FA-like extracts do not show a significant increase in the charge after the first composting cycle, thus indicating that no significant changes in chemical composition or molecular structure occur in this fraction after the 15-day decomposition phase. On the other hand, for the HA-like samples the separation between the charge curves is significant during the whole composting process.

The protonation constants of the acid groups indicate that these are more acidic in the FA-like fraction. Average values for $\log K$ of carboxylic and phenolic groups were 2.96 and 7.86 for FA-like, and 3.49 and 8.21 for HA-like, respectively. These are comparable to the generic values obtained by Milne et al. (2001) and to the values specifically obtained for HS-like extracted from compost found in the literature (Table

S6, Supporting Information). Although these values are similar, the log K for the carboxylic groups of generic HS are generally more acidic than those of HS-like extracted from compost. In the case of phenolic groups, a clear difference is observed in the FA fraction, being less acidic in the case of aquatic and terrestrial FA. As the composting process advances, log K experienced a shift toward a more acidic character (Table 5). In the case of FA-like extracts, log K for the carboxylic and phenolic groups after 15 days of composting were 3.53 and 8.00, respectively, while at the end of the process these values were 2.93 and 7.76. For the HA-like extracts, the values changed from 4.15 and 8.14, after 15 days, to 3.41 and 7.88, after 60 days, respectively. The same trend was described by Plaza et al. (2005), although the decrease in their log K values for the HA-like was more pronounced. This result indicates that, in general, the HS-like become more acidic during the composting process possibly due to the oxidation reactions in the lignin or to the microbial degradation of carbohydrates (Sánchez-Monedero et al., 2002).

3.3. Testing the compost as soil amendment

The physico-chemical characteristics of the Haplic Cambisol and the variations in these after amendment with compost of different feedstock (except compost CVDW, which was not applied due to limited sample availability) are shown in Table 6. Soil Ch2 showed an acidic character, characteristic of mine soils developed in areas rich in mineral sulphides. The carbon content was lower than 1% and the values of available phosphorus (Olsen P) and CEC were relatively low, reflecting the low productivity of the soil (Hazelton and Murphy, 2007).

Application of compost led to significant increase in the values of the different soil quality indicators, particularly pH, available phosphorus and CEC. This effect increased with the amount of compost included in the amendment. The pH of the compost samples ranged between 6.16 and 9.76, and application of the amendment therefore had a positive effect on the soil. In degraded or unproductive soils, increasing the pH is a key factor for *in situ* revegetation (Forjan et al., 2017). The only compost sample that did not yield pH values higher than 7 in the amended soil was CSS. However, this compost, together with compost CUW supplied the greatest amounts of organic matter to the soil. Application of amendments containing 10% of CUW or CSS increased the carbon content from 0.66% to 3.28% and 3.47%, respectively. The nitrogen content was below the limit of detection in the unamended soil but increased significantly after application of compost (Table 6). Compost CSS yielded the greatest increase in total P in the soil after application but did not supply as much available P as the other composts. Thus, application of composts CDDW and CLW produced an increase in the concentration of Olsen P in the soil from 4.6 mg/kg to 206.6 and 365 mg/kg, respectively. Both samples contain more than 10% of the total P as available P (Table 1), which may favour plant productivity. Finally, compost CLW also generated the greatest increase in the CEC relative to the control, from 1.27 to 37.3 cmol(+)/kg. The unamended soil contained low amounts of exchange cations, especially of Ca and K. The concentration of these ions increased with the percentage of compost in the soil, irrespective of the feedstock (Table S7, Supporting Information).

The parameters determined in the germination tests, i.e. normalized residual elongation (NRE), relative radicle growth (RRG), relative seed germination (RSG) and germination index (GI), are reported in Table 6. Seed germination was favoured by increasing the content of compost, as reflected by the increase in GI, although the intensity of the increase depended on the compost used. Thus, the GI values were highest in samples amended with composts CLW and CDDW. Application of compost CLW

generated the greatest increase in the concentration of available P and exchange cations, and yielded the optimal pH for plant growth. On the other hand, the smallest increase in GI occurred with compost CSS. Although this compost yielded the greatest increase in C, N and total P in the soil, it did not generate such a large increase in pH as the other compost. This indicates that pH is a key factor for selecting the compost used to amend this type of soil. Compost CSS was produced from sewage sludge, and therefore the concentration of species that are potentially toxic to *Lolium perenne* may be significant. Other factors that inhibit seed germination include high salinity and excess of ammonium (Hoekstra et al., 2002). The concentration of ammonium ion in the equilibrium solutions used in the germination assay was not determined, although it reached values higher than 150 mg/L in the original CSS sample (Table 2).

The lower positive effect of CSS was also reflected in the values of the NRE index (related to root growth), which reached a maximum value of 0.17 in the soil amended with 10% of compost. Root growth is a more sensitive parameter, although it is more difficult to quantify than seed germination. For the other samples, the values of NRE ranged between 0.33 and 0.53, and application of compost CLW had the most positive effect. The effects of the application of compost on shoot and root growth are shown in Figure S10 (Supporting Information). No clear trend was observed in the growth of the aerial part of the seedlings in relation to the different amendments, nor in the different compost percentages in each. However, the effect on root growth decreased in the order CLW > CVA > CUW > CDDW > CSS. A good correlation between the NRE index and the %N in the compost was observed (Figure S11, Supporting Information), although the variables were inversely related, so that as the N content increased, the root growth decreased. The correlations between total P and K, were lower (R^2 , 0.28-0.60). The increase in K led to an increase in the NRE index, whereas the opposite effect was observed in the case of total P. Thus, sample CSS generated a smaller increase in root growth because the NPK index (3.6-4.8-0.5) showed that the proportion of K was lower than that of the other two macronutrients (Figure S12, Supporting Information). The opposite was observed for sample CLW (NPK, 1.8-3.1-3.6), so that the use of this compost as an amendment favours growth.

4. Conclusions

Compost and vermicompost were studied to evaluate both the effect of the composting process and the effect of the feedstock material. The evolution of the organic matter was also monitored during the composting time. A decrease of the C/N ratio led to a CEC increase as the composting process progressed, whereas a linear increase of CEC as a function of C/N ratio was observed when samples from different feedstock were analysed. In both cases the correlation can be interpreted as an increase in the degree of maturity. The amount of C in the equilibrium solution can be used to quantify the HS-like fraction of the compost given the good correlation between these two parameters. This is particularly important as the classical extraction procedure for humic substances is time-consuming. The worms that participate in the vermicomposting process seem to perform a more efficient degradation of the original aliphatic materials, as it can be deduced by the higher abundance of acidic groups compared to traditionally composted samples.

The use of the compost as soil amendments improved the quality of acidic, unproductive mine soils. Depending on the feedstock material, different parameters were

improved, i.e. urban waste compost generated a greater increase in the organic carbon, while sewage sludge compost and livestock manure compost provided large amounts of major nutrients. Nevertheless, the nutrient content of the compost must be balanced to optimize improvements in soil fertility.

Acknowledgments

This work was financially supported by the Interreg VA Spain-Portugal Programme (EU) through the project Res2ValHum (0366_RES2VALHUM_1_P). R.L, J.A and S.F. are members of the CRETUS Institute (CRETUS, AGRUP2015/02, ref. 2018-PG10) and were also supported by the Group of Excellence GI-1245 financed by the Government of Galicia (ED431C 2018/12). A.C. Silva acknowledges receipt of a PhD grant (UMINHO/BD/40/2016) financed by the Operational Program Norte 2020 (through the Project NORTE-08-5369-FSE-000033). LIPOR and other companies are also acknowledged for the supply of the compost samples. The authors thank M. Santiso, F Casas, J. Maceira and D. Fiuza for their assistance in characterizing the compost and soil samples.

CRedit authorship contribution statement

Rocío López: Conceptualization, methodology, investigation, formal analysis, data curation, writing – original draft. **Juan Antelo:** Methodology, investigation, data curation, writing – review & editing. **Ana Catarina Silva:** Methodology, investigation, data curation, writing – review & editing. **Fátima Bento:** Conceptualization, funding acquisition, resources, validation, writing – review & editing. **Sarah Fiol:** Methodology, funding acquisition, resources, project administration, writing – review & editing.

References

- Adani, F., Genevini, P.L., Gasperi, F., Tambone, F., 1999. Composting and humification. *Compost Sci. Util.* 7, 24–33. <https://doi.org/10.1080/1065657X.1999.10701949>
- Amir, S., Hafidi, M., Bailly, J.R., Revel, J.C., 2003. Characterization of humic acids extracted from sewage sludge during composting and of their Sephadex R gel fractions. *Agronomie* 23, 269– 275. <https://doi.org/10.1051/agro:2002090>
- Amir, S., Jouraiphy, A., Meddich, A., El Gharous, M., Winterton, P., Hafidi, M., 2010. Structural study of humic acids during composting of activated sludge-green waste: Elemental analysis, FTIR and ¹³C NMR. *J. Hazar. Mater.* 177, 524–529. <https://doi.org/10.1016/j.jhazmat.2009.12.064>
- Arán, D., Verde, J.R., Antelo, J., Macías F., 2020. Environmental quality of soils and waters in Fé Mine: Initial situation and remediation alternatives. *Span. J. Soil Sci.* 10, 81–100. <https://doi.org/10.3232/SJSS.2020.V10.N1.06>
- Baddi G.A., Hafidi M., Cegarra J., Albuquerque J.A., González J., Gilard V., Revel J.-C., 2004. Characterization of fulvic acids by elemental and spectroscopic (FTIR

- and ^{13}C -NMR) analyses during composting of olive mill wastes plus straw. *Bioresour. Technol.* 93, 285–290. <https://doi.org/10.1016/j.biortech.2003.10.026>
- Bagur-González, M.G., Estepa-Molina, C., Martín-Peinado, F., Morales-Ruano, S., 2011. Toxicity assessment using *Lactuca sativa* L. bioassay of the metal(loid)s As, Cu, Mn, Pb and Zn in soluble-in-water saturated soil extracts from an abandoned mining site. *J. Soil. Sediment.* 11, 281–289. <https://doi.org/10.1007/s11368-010-0285-4>
- Bernal, M.P., Paredes, C., Sánchez-Monedero, A., Cegarra, J., 1998. Maturity and stability parameters of composts prepared with a wide range of organic wastes. *Bioresour. Technol.* 63, 91–99. [https://doi.org/10.1016/S0960-8524\(97\)00084-9](https://doi.org/10.1016/S0960-8524(97)00084-9)
- Bernal, M.P., Alburquerque, J.A., Moral, R., 2009. Composting of animal manures and chemical criteria for compost maturity assessment. A review. *Bioresour. Technol.* 100, 5444–5453. <https://doi.org/10.1016/j.biortech.2008.11.027>
- Cerdá, A., Artola, A., Font, X., Barrena, R., Gea, T., Sánchez, A., 2018. Composting of food wastes: Status and challenges. *Bioresour. Technol.* 248, Part A, 57–67. <https://doi.org/10.1016/j.biortech.2017.06.133>
- Christl, I., Kretzschmar, R., 2001. Relating ion binding by fulvic and humic acids to chemical composition and molecular size. 1. Proton binding. *Environ. Sci. Technol.* 35, 2505–2511. <https://doi.org/10.1021/es0002518>
- Di Salvatore, M., Carafa, A.M., Carratù, G., 2008. Assessment of heavy metals phytotoxicity using seed germination and root elongation tests: A comparison of two growth substrates. *Chemosphere* 73, 1461–1464. <https://doi.org/10.1016/j.chemosphere.2008.07.061>
- Droussi, Z., D’orazio, V., Provenzano, M.R., Hafidi, M., Ouattmane, A., 2009. Study of the biodegradation and transformation of olive-mill residues during composting using FTIR spectroscopy and differential scanning calorimetry. *J. Hazard. Mater.* 164, 1281–1285. <https://doi.org/10.1016/j.jhazmat.2008.09.081>
- Fernández, J.M., Plaza, C., Senesi, N., Polo, A., 2007. Acid-base properties of humic substances from composted and thermally-dried sewage sludges and amended soils as determined by potentiometric titration and the NICA-Donnan model. *Chemosphere* 69, 630–635. <https://doi.org/10.1016/j.chemosphere.2007.02.063>
- Forjan, R., Rodríguez-Vila, A., Cerqueira, B., Covelo, E.F., 2017. Comparison of the effects of compost versus compost and biochar on the recovery of a mine soil by improving the nutrient content. *J. Geochem. Explor.* 183, 46–57. <https://doi.org/10.1016/j.gexplo.2017.09.013>
- Fornes, F., Mendoza-Hernández, D., García-de-la-Fuente, R., Abad, M., Belda, R.M., 2012. Composting versus vermicomposting: A comparative study of organic matter evolution through straight and combined processes. *Bioresour. Technol.* 118, 296–305. <https://doi.org/10.1016/j.biortech.2012.05.028>

- Fuentes, M., Baigorri, R., González-Gaitano, G., García-Mina, J., 2018. New methodology to assess the quantity of humic substances in organic materials and commercial products for agriculture. *J. Soil. Sediment.* 18, 1389–1399. <https://doi.org/10.1007/s11368-016-1514-2>
- Fuentes, M., González-Gaitano, G., García-Mina, J.M., 2006. The usefulness of UV-visible and fluorescence spectroscopies to study the chemical nature of humic substances from soils and composts. *Org. Geochem.* 37, 1949–1959. <https://doi.org/10.1016/j.orggeochem.2006.07.024>
- Guo, X., Liu, H., Wu, S., 2019. Humic substances developed during organic waste composting: Formation mechanisms, structural properties, and agronomic functions. *Sci. Total Environ.* 662, 501–510. <https://doi.org/10.1016/j.scitotenv.2019.01.137>
- Hanc, A., Hrebeckova, T., Kuzel, S., 2019a. Vermicomposting of distillery residues in a vertical-flow windrow system. *Waste Biomass Valori.* 10, 3647–3657. <https://doi.org/10.1007/s12649-019-00671-4>
- Hanc, A., Enev, V., Hrebeckova, T., Klucakova, M., Pekar, M., 2019b. Characterization of humic acids in a continuous-feeding vermicomposting system with horse manure. *Waste Manage.* 99, 1–11. <https://doi.org/10.1016/j.wasman.2019.08.032>
- Hanc, A., Vasak, F., 2015. Processing separated digestate by vermicomposting technology using earthworms of the genus *Eisenia*. *Int. J. Environ. Sci. Technol.* 12, 1183–1190. <https://doi.org/10.1007/s13762-014-0500-8>
- Harada, Y., Inoko, A., 1980. Relationship between cation exchange capacity and degree of maturity of city refuse compost. *Soil Sci. Plant Nutr.* 26, 352–362. <https://doi.org/10.1080/00380768.1980.10431220>
- Hazelton, P., Murphy, B., 2007. *Interpreting Soils Test Resulting. What do All the Numbers Mean?* CSIRO Publishing, Australia.
- Hoekstra, N.J., Bosker, T., Lantinga, E.A., 2002. Effects of cattle dung from farms with different feeding strategies on germination and initial root growth of cress (*Lepidium sativum* L.). *Agric. Ecosyst. Environ.* 93, 189–196. [https://doi.org/10.1016/S0167-8809\(01\)00348-6](https://doi.org/10.1016/S0167-8809(01)00348-6)
- Hollins, O., Lee, P., Sims, E., Bertham, O., Symington, H., Bell, N., Pfaltzgraff, L., Sjögren, P., 2017. *Towards a Circular Economy - Waste Management in the EU, Study IP/G/STOA/FWC/2013-001/LOT 3/C3 EPRS, STOA.* Brussels, EU.
- Hunee, S.Y., Clemente, R., Bernal, M.P., 2020. Selection of Mediterranean plants biomass for the composting of pig slurry solids based on the heat production during aerobic degradation. *Waste Manage.* 104, 1–8. <https://doi.org/10.1016/j.wasman.2020.01.001>
- Jouraihy, A., Amir, S., Winterton, P., El Gharous, M., Revel, J-C., Hafidi, M., 2005. Chemical and spectroscopic analysis of organic matter transformation during

- composting of sewage sludge and green plant waste. *Int. Biodeterior. Biodegrad.* 56, 101–108. <https://doi.org/10.1016/j.ibiod.2005.06.002>
- Kaushik, P., Garg, V.K., 2003. Vermicomposting of mixed solid textile mill sludge and cow dung with the epigeic earthworm *Eisenia foetida*. *Bioresour. Technol.* 90, 311–316. [https://doi.org/10.1016/S0960-8524\(03\)00146-9](https://doi.org/10.1016/S0960-8524(03)00146-9)
- Kinniburgh, D.G., Milne, C.J., Benedetti, M.F., Pinheiro, J.P., Filius, J., Koopal, L.K., van Riemsdijk, W.H., 1996. Metal ion binding by humic acids: application of the NICA-Donnan model. *Environ. Sci. Technol.* 30, 1687–1698. <https://doi.org/10.1021/es950695h>
- Kucbel, M., Raclavská, H., Růžičková, J., Švédová, B., Sassmanová, V., Drozdová, J., Raclavský, K., Juchelková, D., 2019. Properties of composts from household food waste produced in automatic composters. *J. Environ. Manage.* 236, 657–666. <https://doi.org/10.1016/j.jenvman.2019.02.018>
- Kumar, M., Ou, Y.L., Lin, J.G., 2010. Co-composting of green waste and foodwaste at low C/N ratio. *Waste Manage.* 30, 602–609. <https://doi.org/10.1016/j.wasman.2009.11.023>
- Lasaridi, K., Protopapa, I., Kotsou, M., Pilidis, G., Manjos, T., Kyriacou, A., 2006. Quality assessment of composts in the Greek market: the need for standards and quality assurance. *J. Environ. Manage.* 80, 58–65. <https://doi.org/10.1016/j.jenvman.2005.08.011>
- López, R., Fiol, S., Antelo, J.M., Arce, F., 2003. Analysis of the effect of concentration on the acid-base properties of soil fulvic acid. Conformational changes. *Colloid. Surface A* 226, 1–8. [https://doi.org/10.1016/S0927-7757\(03\)00358-3](https://doi.org/10.1016/S0927-7757(03)00358-3)
- Lv, B., Xing, M., Yang, J., Qi, W., Lu, Y., 2013. Chemical and spectroscopic characterization of water extractable organic matter during vermicomposting of cattle dung. *Bioresour. Technol.* 132, 320–326. <https://doi.org/10.1016/j.biortech.2013.01.006>
- Milne, C.J., Kinniburgh, D.G., Tipping, E., 2001. Generic NICA-Donnan model parameters for proton binding by humic substances. *Environ. Sci. Technol.* 35, 2049–2059. <https://doi.org/10.1021/es000123j>
- Murphy, J., Riley, J.P., 1962. A modified single solution method for the determination of phosphate in natural waters. *Anal. Chim. Acta* 27, 31–36. [https://doi.org/10.1016/S0003-2670\(00\)88444-5](https://doi.org/10.1016/S0003-2670(00)88444-5)
- Palansooriya, K.N., Shaheen, S.M., Chen, S.S., Tsang, D.C.W., Hashimoto, Y., Hou, D., Bolan, N.S., Rinklebe, J., Ok, Y.S., 2020. Soil amendments for immobilization of potentially toxic elements in contaminated soils: A critical review. *Environ. Int.* 134, 105046. <https://doi.org/10.1016/j.envint.2019.105046>

- Plaza, C., García-Gil, J.C., Polo, A., Senesi, N., Brunetti, G., 2005. Proton binding by humic and fulvic acids from pig slurry and amended soils. *J. Environ. Qual.* 34, 1131–1137. <https://doi.org/10.2134/jeq2004.0378>
- Riffaldi, R., Levi-Minzi, R., Saviozzi, A., 1983. Humic fractions of organic waste. *Agr. Ecosyst. Environ.* 10, 353–359. [https://doi.org/10.1016/0167-8809\(83\)90086-5](https://doi.org/10.1016/0167-8809(83)90086-5)
- Ritchie, J.D., Perdue, E.M., 2008. Analytical constraints on acidic functional groups of humic substances. *Org. Geochem.* 39, 783–799. <https://doi.org/10.1016/j.orggeochem.2008.03.003>
- Roe, N., Stoffella, P., Graetz, D., 1997. Compost from various municipal solid waste feedstocks affect vegetable crops. I. Emergence and seedling growth. *J. Am. Soc. Horticult. Sci.* 122, 427–432. <https://doi.org/10.21273/JASHS.122.3.433>
- Sánchez-Monedero, M.A., Cegarra, J., García, D., Roig, A., 2002. Chemical and structural evolution of humic acids during organic waste composting. *Biodegrad.* 13, 361–371. <https://doi.org/10.1023/A:1022888231982>
- Sciubba, L., Cavani, L., Grigatti, M., Ciavatta, C., Marzadori, C., 2015. Relationships between stability, maturity, water-extractable organic matter of municipal sewage sludge composts and soil functionality. *Environ. Sci. Pollut. Res.* 22, 13393–13403. <https://doi.org/10.1007/s11356-015-4611-7>
- Silva, M.E.F., Lemos, L.T., Bastos, M.M.S.M., Nunes, O.C., Cunha-Queda, A.C., 2013. Recovery of humic-like substances from low quality composts. *Bioresour. Technol.* 128, 624–632. <https://doi.org/10.1016/j.biortech.2012.11.013>
- Soobhany, N., Gunasee, S., Rago, Y.P., Joyram, H., Raghoo, P., Mohee, R., Garg, V.K., 2017. Spectroscopic, thermogravimetric and structural characterization analyses for comparing Municipal Solid Waste composts and vermicomposts stability and maturity. *Bioresour. Technol.* 236, 11–19. <https://doi.org/10.1016/j.biortech.2017.03.161>
- Spaccini, R., Piccolo, A., 2009. Molecular characteristics of humic acids extracted from compost at increasing maturity stages. *Soil Biol. Biochem.* 41, 1164–1172. <https://doi.org/10.1016/j.soilbio.2009.02.026>
- Sparks, D.L., Page, A.L., Helmke, P.A., Loeppert, R.H., 1996. *Methods of Soil Analysis. Part 3 - Chemical Methods.* SSSA Book Ser. 5. Soil Science Society of America, America Society of Agronomy, Madison, Wisconsin, USA.
- Suthar, S., 2010. Recycling of agro-industrial sludge through vermitechnology. *Ecol. Eng.* 36, 1028–1036. <https://doi.org/10.1016/j.ecoleng.2010.04.015>
- Swift, R.S., 1996. Organic matter characterization, in: Sparks, D.L., Page, A.L., Helmke, P.A., Loeppert, R.H., Soltanpour, P.N., Tabatabai, M.A., Johnston, C.T., Summer, M.E. (Eds.), *Methods of Soil Analysis: Part 3. Chemical Methods.* Soil Science Society of America, Madison, WI, pp. 1011–1069.

- Teodoro, M., Trakal, L., Gallagher, B.N., Simek, P., Soudek, P., Pohorelý, Beesley, L., Jacka, K., Kova, M., Seyedsadr, S., Mohan, D., 2020. Application of co-composted biochar significantly improved plant-growth relevant physical/chemical properties of a metal contaminated soil. *Chemosphere* 242, 125255. <https://doi.org/10.1016/j.chemosphere.2019.125255>
- van der Wurff, A.W.G., Fuchs, J.G., Raviv, M., Termorshuizen, A.J., 2016. Handbook for Composting and Compost Use in Organic Horticulture. BioGreenhouse COST Action FA 1105, www.biogreenhouse.org. <http://dx.doi.org/10.18174/375218>
- van Zomeren, A., Costa, A., Pinheiro, J.P., Comans, R.N.J., 2009. Proton binding properties of humic substances originating from natural and contaminated materials. *Environ. Sci. Technol.*, 43, 1393–1399. <https://doi.org/10.1021/es0709223>
- Veeken, A., Nierop, K., de Wilde, V., Hamelers, B., 2000. Characterisation of NaOH-extracted humic acids during composting of a biowaste. *Bioresour. Technol.* 72, 33–41. [https://doi.org/10.1016/S0960-8524\(99\)90096-2](https://doi.org/10.1016/S0960-8524(99)90096-2)
- Walkley, A., 1947. A critical examination of a rapid method for determining organic carbon in soils - effect of variations in digestion conditions and of inorganic soil constituents. *Soil Sci.* 63, 251–264. <http://dx.doi.org/10.1097/00010694-194704000-00001>
- Walter, I., Martínez, F., Cala, V., 2006. Heavy metal speciation and phytotoxic effects of three representative sewage sludges for agricultural uses. *Environ. Pollut.* 139, 507–514. <https://doi.org/10.1016/j.envpol.2005.05.020>
- Wang, L., Li, X., Prasher, S.O., Yan, B., Ou, Y., Cui, H., Cui, Y., 2019. Organic matter, a critical factor to immobilize phosphorus, copper, and zinc during composting under various initial C/N ratios. *Bioresour. Technol.*, 289, 121745. <https://doi.org/10.1016/j.biortech.2019.121745>

Figure captions

Figure 1: Correlation between CEC and C/N ratio of the compost. The solid line corresponds to the positive correlation observed for compost of different feedstock (filled symbols) and the dashed line shows a negative correlation during the composting process (empty symbols).

Figure 2. Variation of the electric conductivity (EC) as a function of the pH in the equilibrium solution of the compost samples.

Figure 3. Relationship of the yield in the HS-like extraction and a) the C_{oxi} content in the equilibrium solution and b) the pH of the equilibrium solution.

Figure 4. a) Charge curve and proton affinity distributions in the DOM of the equilibrium solution of sample CLW as a function of pH. b) Proton affinity distributions in the DOM of the equilibrium solution of compost samples of different origin as a function of pH.

Figure 5. Charge curves of the HS-like of sample CLW as a function of pH. Symbols represent experimental data obtained from potentiometric titrations and lines represent fits with equation 6.

Figure 6. Charge curves of the a) FA-like and b) HA-like extracted at different stages of the composting process of sample CUW. Symbols represent experimental data from potentiometric titrations and lines represent fits with equation 6.

Table captions

Table 1. Chemical composition of the compost samples. Percentages in parentheses are calculated based on the total content.

Table 2. Chemical characterization of the equilibrium solutions.

Table 3. Chemical characterization and extraction yield of the HS-like obtained with the extraction protocol of the IHSS.

Table 4: Acid-base parameters of the dissolved organic matter in the equilibrium solutions from compost samples of different feedstock and different composting stages. Errors associated to the fitting are also shown.

Table 5. Intrinsic acid-base parameters of the HS-like extracted from the compost. Errors associated to the fitting are also shown.

Table 6. Variation of the soil properties and the parameters obtained in the germination assays following the application of compost as a soil amendment.

Table 1.

	%C	%C _{org}	%N	C/N	%S	P _{total} (mg/kg)	P _{soiln} (mg/kg)	Ca (mg/kg)	Mg (mg/kg)	Na (mg/kg)	K (mg/kg)	CEC cmol(+)/Kg
Compost	CLW	22.8 (17.8%) (77.8%)	1.8	12.8	0.7	13505	3452 (25.6%)	78040	18230	11410	30187	149.7
	CUW	34.0 (29.3%) (86.1%)	2.7	12.6	0.2	5002	279 (5.6%)	77470	3739	4830	12237	167.0
	CDDW	23.3 (18.8%) (81.0%)	2.3	10.3	0.2	2513	251 (10.0%)	16100	3621	618.1	7597	95.3
	CSS	31.3 (30.9%) (98.6%)	3.6	8.6	1.1	21021	508 (2.4%)	19880	5200	2150	4269	51.5
	CVA	29.7 (27.5%) (92.4%)	2.4	12.5	0.4	5512	1386 (25.1%)	13100	4516	2807	10247	136.7
	CVDW	43.9 (33.1%) (75.4%)	5.2	8.5	0.4	4152	297 (7.2%)	9632	4430	1089	14577	98.8
Maturation	UW0	40.4 (38.6%) (95.4%)	2.5	15.9	0.2	3129	295 (9.4%)	25670	2520	4300	10063	127.6
	CUW15	38.9 (36.9%) (95.0%)	2.8	13.8	0.3	3950	255 (9.0%)	28420	2637	4531	8878	96.3
	CUW30	35.2 (34.8%) (98.9%)	2.7	12.9	0.3	4717	474 (10.0%)	43100	3451	5190	10737	108.7

Table 2.

		DOC (mg/L)	%C ^a	NH ₄ ⁺ (mg/L)	Ca (mg/L)	Mg (mg/L)	Na (mg/L)	K (mg/L)	PO ₄ ³⁻ (mg/L)	pH	EC (μS/cm)
Compost	CLW	861	9.7	21.2	22.5	38.9	552.0	1363.6	124.1	8.5	6020
	CUW	548	3.7	7.8	28.0	8.8	175.4	349.2	29.5	7.7	2840
	CDDW	115	1.2	4.6	28.6	12.3	17.3	149.5	45.2	6.9	883
	CSS	698	4.5	159.1	95.7	55.8	21.1	55.0	29.0	7.6	3850
Vermicompost	CVA	111	0.8	14.1	80.7	66.1	114.2	431.0	241.4	6.7	3280
	CVDW	717	4.3	11.1	17.8	15.1	26.7	333.2	70.3	6.3	1370
Maturation	UW0	1187	6.2	10.8	197.0	62.3	207.9	349.9	110.2	6.4	3870
	CUW15	429	2.3	13.2	133.1	37.4	194.3	322.0	80.4	6.8	3030
	CUW30	430	2.5	10.1	64.1	24.9	225.6	533.1	60.0	7.9	3300

^a%C oxidizable released to the solution

Table 3

		FA							HA								
		yield (g/kg)	%C	%N	%H	%O	C/N	O/C	H/C	yield (g/kg)	%C	%N	%H	%O	C/N	O/C	H/C
Compost	CLW	7.0	35.4	3.4	3.4	56.1	12.3	1.2	1.1	34.9	55.2	6.0	5.7	33.4	11.4	0.4	1.2
	CUW	3.3	40.8	4.4	6.0	47.8	10.9	0.9	1.8	21.1	54.5	6.7	6.1	31.7	9.4	0.4	1.3
	CDDW	4.2	38.4	2.6	3.1	55.5	17.1	1.1	1.0	19.0	50.6	4.9	4.5	39.5	12.0	0.6	1.1
	CSS	4.8	40.9	6.1	5.8	45.0	7.9	0.8	1.7	23.8	58.5	5.8	8.3	25.5	11.8	0.3	1.7
Vermicompost	CVA	2.3	49.7	3.7	5.4	40.1	15.6	0.6	1.3	33.4	53.5	3.7	5.5	36.2	16.8	0.5	1.2
	CVDW	4.7	39.8	3.2	3.6	52.9	14.5	1.0	1.1	26.4	52.7	6.1	4.6	36.1	10.0	0.5	1.1
	UW0	2.2	52.2	6.2	6.2	34.6	9.9	0.5	1.4	33.1	59.8	5.4	8.0	26.1	12.8	0.3	1.6
Maturation	CUW15	2.7	52.3	5.2	6.1	35.7	11.8	0.5	1.4	19.8	54.4	5.4	6.9	32.4	11.7	0.4	1.5
	CUW30	2.4	50.3	4.3	5.7	38.8	13.8	0.6	1.3	21.9	54.6	5.6	5.7	33.1	11.3	0.5	1.3

Table 4

	CLW	CUW	CDDW	CSS	CVA	CVDW	UW0	CUW15	CUW30
log K₁	3.67	3.69	3.06	4.24	3.27	3.45	4.50	3.99	3.57
(*)M ₁	0.09	0.06	0.06	0.18	0.04	0.08	0.42	0.05	0.03
	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01
log K₂	6.56	6.51	6.42	6.27	6.55	6.65	6.05	6.54	6.65
(*)M ₂	0.43	0.18	0.10	0.25	0.08	0.33	0.16	0.24	0.25
	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01
log K₃	10.25	10.11	10.17	9.65	10.00	10.33	9.73	10.06	10.08
(*)M ₃	0.45	0.19	0.10	0.74	0.10	0.21	0.27	0.22	0.21
	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01
(*)Q ₀	-0.48	-0.23	-0.14	-0.40	-0.10	-0.36	-0.52	-0.26	-0.26
	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01
R ²	0.9997	0.9997	0.9999	0.9998	0.9996	0.9993	0.9998	0.9994	0.9991
RMSE	0.0055	0.0024	0.0009	0.0062	0.0013	0.0052	0.0035	0.0041	0.0050
(*)M _T	0.96	0.44	0.25	1.17	0.22	0.62	0.85	0.52	0.50

(*) Parameters in mmol/g_{composit}. M_T corresponds to the total abundance of acid positions

Table 5

	b	log K ₁	M ₁ (mmol/g ^{sH})	m ₁	log K ₂	M ₂ (mmol/g ^{sH})	m ₂	R ²	RMSE	M _T (mmol/g ^{sH})
FA CLW	0.57	2.72 ± 0.02	3.02 ± 0.02	0.44 ± 0.02	8.31 ± 0.02	1.56 ± 0.02	0.43 ± 0.02	0.9997	0.0154	4.58
FA CUW	0.63	2.93 ± 0.01	3.49 ± 0.08	0.60 ± 0.01	7.76 ± 0.16	1.62 ± 0.23	0.34 ± 0.05	0.9998	0.0142	5.11
FA CDDW	0.66	2.76 ± 0.01	3.46 ± 0.07	0.54 ± 0.01	7.58 ± 0.08	1.51 ± 0.16	0.30 ± 0.03	0.9999	0.0073	4.96
FA CSS	0.65	3.38 ± 0.04	4.77 ± 0.16	0.45 ± 0.01	8.07 ± 0.17	2.81 ± 0.49	0.39 ± 0.06	0.9998	0.0260	7.57
FA CVA	0.60	3.31 ± 0.01	3.01 ± 0.15	0.88 ± 0.03	7.42 ± 0.14	2.61 ± 0.39	0.32 ± 0.05	0.9996	0.0270	5.63
FA CVDW	0.63	2.63 ± 0.01	3.89 ± 0.04	0.54 ± 0.01	8.03 ± 0.06	1.72 ± 0.11	0.40 ± 0.03	0.9997	0.0190	5.61
Average FA ^a	0.62 ± 0.03	2.96 ± 0.32	3.61 ± 0.60	0.58 ± 0.15	7.86 ± 0.33	1.97 ± 0.53	0.36 ± 0.05			5.58 ± 0.97
HA CLW	0.47	3.64 ± 0.02	2.64 ± 0.03	0.39 ± 0.01	8.43 ± 0.10	1.31 ± 0.13	0.57 ± 0.05	0.9999	0.0070	3.95
HA CUW	0.46	3.41 ± 0.04	2.80 ± 0.19	0.60 ± 0.03	7.88 ± 0.22	1.94 ± 0.48	0.33 ± 0.08	0.9994	0.0272	4.74
HA CDDW	0.48	3.15 ± 0.04	2.93 ± 0.11	0.47 ± 0.01	7.96 ± 0.22	1.49 ± 0.31	0.36 ± 0.07	0.9998	0.0132	4.42
HA CSS	0.43	3.78 ± 0.07	2.60 ± 0.07	0.30 ± 0.01	8.46 ± 0.21	1.11 ± 0.24	0.55 ± 0.09	0.9999	0.0081	3.70
HACVA	0.45	3.46 ± 0.03	2.47 ± 0.03	0.44 ± 0.01	8.37 ± 0.11	1.09 ± 0.13	0.57 ± 0.06	0.9998	0.0124	3.56
HA CVDW	0.44	3.47 ± 0.02	2.43 ± 0.04	0.60 ± 0.01	8.15 ± 0.19	1.32 ± 0.20	0.50 ± 0.07	0.9998	0.0138	3.75
Average HA ^a	0.46 ± 0.02	3.49 ± 0.21	2.62 ± 0.19	0.47 ± 0.11	8.21 ± 0.25	1.38 ± 0.29	0.48 ± 0.10			4.00 ± 0.44

^aCalculated average value for the model parameters of HS-like extracts

	b	log K ₁	M ₁ (mmol/g ^{sH})	m ₁	log K ₂	M ₂ (mmol/g ^{sH})	m ₂	R ²	RMSE	M _T (mmol/g ^{sH})
FA CUW15	0.74	3.53 ± 0.01	3.38 ± 0.09	0.87 ± 0.02	8.00 ± 0.27	1.62 ± 0.33	0.35 ± 0.08	0.9995	0.0260	5.00
FA CUW30	0.68	3.15 ± 0.01	3.45 ± 0.03	0.71 ± 0.01	7.96 ± 0.07	1.69 ± 0.09	0.34 ± 0.02	1.0000	0.0060	5.14
HA CUW15	0.50	4.15 ± 0.02	2.15 ± 0.08	0.69 ± 0.02	8.14 ± 0.21	1.42 ± 0.28	0.41 ± 0.08	0.9998	0.0132	3.57
HA CUW30	0.48	3.93 ± 0.03	2.50 ± 0.09	0.59 ± 0.01	8.12 ± 0.28	1.37 ± 0.34	0.41 ± 0.09	0.9998	0.0133	3.88

Table 6

	pH	%C	%N	P-total (mg/kg)	P-Olsen (mg/kg)	CEC (cmol_c/kg)	Root length (cm)	NRE	RGS(%)	RRG(%)	GI(%)
<i>Ch2 (Control)</i>	4.53	0.66	n.d.	258.0	4.6	1.27	4.32	0.0	100.0	100.0	100.0
<i>Ch2 + 2%CLW</i>	7.13	0.96	0.05	509.8	53.6	12.3	6.39	0.48	100.0	147.7	147.7
<i>Ch2 +5%CLW</i>	7.77	1.59	0.09	640.4	105.7	27.1	6.62*	0.53*	109.4	153.1*	167.5*
<i>Ch2 + 10%CLW</i>	8.19*	1.90	0.13	846.6	365.0*	37.3*	6.56	0.52	103.8	151.8	157.5
<i>Ch2 + 2%CUW</i>	6.63	1.06	0.02	347.4	3.8	4.4	5.75	0.34	113.2	134.1	151.8
<i>Ch2 +5%CUW</i>	7.59	1.38	0.06	429.5	9.9	13.2	5.80	0.47	105.7	147.4	155.8
<i>Ch2 + 10%CUW</i>	7.77	3.28	0.32	603.5	22.0	20.0	6.37	0.33	107.5	133.0	143.0
<i>Ch2 + 2%CSS</i>	5.34	1.29	n.d.	704.5	28.1	2.8	3.97	-0.08	103.8	91.9	95.4
<i>Ch2 +5%CSS</i>	5.98	2.59	0.53	1338.2	42.5	6.7	5.01	0.16	103.8	115.8	120.2
<i>Ch2 + 10%CSS</i>	6.15	3.47*	0.39*	1956.2*	101.6	10.8	5.08	0.17	113.2*	117.5	133.0
<i>Ch2 + 2%CVA</i>	6.63	0.98	n.d.	478.8	29.6	6.2	5.97	0.38	100.0	138.0	138.0
<i>Ch2 +5%CVA</i>	7.59	1.36	0.06	733.6	39.9	11.5	6.24	0.44	107.5	144.4	155.3
<i>Ch2 + 10%CVA</i>	7.77	2.04	0.16	1128.3	78.4	23.4	6.47	0.50	100.0	149.6	149.6
<i>Ch2 + 2%CDDW</i>	7.21	0.93	0.06	534.7	48.1	12.1	6.55	0.52	103.8	151.5	157.2
<i>Ch2 +5%CDDW</i>	7.49	1.56	0.48	760.1	69.2	16.1	6.14	0.42	111.3	141.9	158.0
<i>Ch2 + 10%CDDW</i>	7.79	2.29	0.27	1002.4	206.6	22.3	6.26	0.45	110.4	144.7	159.7

n.d. – non detected

*Indicates the case with the higher increase in the corresponding parameter after the amendment application.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

